



International Journal of
**Agricultural
Research**

ISSN 1816-4897



Academic
Journals Inc.

www.academicjournals.com

Impact of Plant Growth Promoting Rhizobacteria on Crop Production

¹Jay Prakash Verma, ²Janardan Yadav, ¹Kavindra Nath Tiwari,

²Lavakush and ²Vimal Singh

¹Department of Botany, MMV, Banaras Hindu University,
Varanasi-221005, Uttar Pradesh, India

²Department of Soil Science and Agricultural Chemistry,
Institute of Agricultural Sciences, Banaras Hindu University,
Varanasi-221005, Uttar Pradesh, India

Abstract: Rhizospheric soils of crop plants have more flora and fauna due to availability of more organic compound, macronutrient and micronutrient. Rhizobacteria that exert beneficial effects on plant growth and development are referred to as Plant Growth Promoting Rhizobacteria (PGPR). Plant growth promoting rhizobacteria is a group of free living soil bacteria, which have ability to promote growth and yield of crop plant by direct and indirect mechanism. PGPR is generally two type, one is colonies inside plant cells that called intracellular PGPR (iPGPR) and other colonies out side plant in rhizosphere that called extracellular PGPR (ePGPR). This review generally focused on direct and indirect mechanism of PGPR. Direct mechanism of plant growth promotion may involve the synthesis of substances by the bacterium or facilitation of the uptake of nutrients from the environment. The indirect mechanism of plant growth occurs when PGPR lessen or prevent the deleterious effects of plant pathogens on plants by production of inhibitory substances or by increasing the natural resistance of the host. The search for PGPR and investigation of their mode of action are increasing at a rapid use as commercial biofertilizers. The mode of action and practical application of PGPR in crop production are the major focus of this review.

Key words: PGPR, rhizosphere, IAA, phosphate solubilization, growth, yield

INTRODUCTION

Bacteria able to colonize plant root systems and promote plant growth are referred to as Plant Growth Promoting Rhizobacteria (PGPR) (Kloepper and Schroth, 1978). PGPR can affect plant growth either indirectly or directly; indirect promotion of plant growth occurs when PGPR lessen or prevent the deleterious effects of one or more phytopathogenic organisms; while direct promotion of plant growth by PGPR involves either providing plants with a compound synthesized by the bacterium or facilitating the uptake of certain nutrients from the environment (Glick, 1995). Thus, in the broadest sense, PGPR include the N₂-fixing rhizobacteria that colonize the rhizosphere, providing N to plants, including the rhizobia of the well-characterized legume-rhizobia symbiosis. Regardless of the mechanisms of plant growth promotion, PGPR must colonize the rhizosphere around the roots, the rhizoplane

Corresponding Author: Jay Prakash Verma, Department of Botany, MMV,
Banaras Hindu University, Varanasi-221005, Uttar Pradesh, India
Tel: 0542-2307120, +91-9452762725 Fax: 0542-2368465

(root surface) or the root itself (within root tissues) (Glick, 1995). Plant growth promoting bacteria (PGPR) associations range in degree of bacterial proximity to the root and intimacy of association. In general, these can be separated into extracellular PGPR (ePGPR), existing in the rhizosphere, on the rhizoplane or in the spaces between cells of the root cortex and intracellular PGPR (iPGPR), which exist inside root cells, generally in specialized nodular structures. The latter includes rhizobia and Frankia species, both of which fix nitrogen in symbiosis with higher plants. There has been considerable development in understanding signaling mechanisms of rhizobia (iPGPR) during the establishment of the rhizobia-legume symbiosis and this may serve as a model of knowledge regarding cross-talk and plant growth promoting mechanisms (Gray and Smith, 2005). The inoculation of seeds with PGPRs is known to increase nodulation, nitrogen uptake and growth and yield response of crop plants (Dorosinsky *et al.*, 1975; Hernandez *et al.*, 1983; Johri *et al.*, 2003; Yadav and Verma, 2009; Verma *et al.*, 2010).

PGPR and Their Hosts: Rhizospheric and Endophytic Region

For PGPR to have a beneficial effect on plant growth via an enhancement of the nutrient status of their host, there obviously needs to be an intimate relationship between the PGPR and the host plant. However, the degree of intimacy between the PGPR and the host plant can vary depending on where and how the PGPR colonizes the host plant. Relationships between PGPR and their hosts can be categorized into two levels of complexity: (1) Rhizospheric and (2) Endophytic.

Rhizospheric Region

The rhizosphere can be defined as any volume of soil specifically influenced by plant roots and/or in association with roots and hairs and plant-produced material (Andrade *et al.*, 1997; Mahafee and Kloepper, 1997; Bringhurst *et al.*, 2001). This space includes soil bound by plant roots, often extending a few mm from the root surface (Bringhurst *et al.*, 2001) and can include the plant root epidermal layer (Mahafee and Kloepper, 1997). Plant exudates in the rhizosphere, such as amino acids and sugars, provide a rich source of energy and nutrients for bacteria, resulting in bacterial populations greater in this area than outside the rhizosphere. Extracellular PGPR (ePGPR) which is existing in the rhizosphere. ePGPR increase plant growth through a variety of mechanisms; these include genera such as *Bacillus*, *Pseudomonas*, *Burkholderia*, *Erwinia*, *Caulobacter*, *Serratia*, *Arthrobacter*, *Micrococcus*, *Flavobacterium*, *Chromobacterium*, *Agrobacterium*, *Hyphomycrobium* and free-living nitrogen-fixing bacteria (*Azotobacter* and *Azospirillum*) (Foster *et al.*, 1983; Prithiviraj *et al.*, 2003; Gray and Smith, 2005). Most rhizosphere organisms occur within 50 mm of root surface and populations within 10 mm of root surface may reach 1.2×10^8 cells cmK^{-3} or 10^9 - 10^{12} microbial cells g k^{-1} soil. Despite large numbers of bacteria in the rhizosphere, only 7-15% of the total root surface is generally occupied by microbial cells (Foster *et al.*, 1983; Pinton *et al.*, 2001; Gray and Smith, 2005).

Endophytic Region

Rhizobacteria that establish inside plant roots, forming more intimate associations, are endophytes. To aid in this conceptualization, simple terms have been adopted: intracellular PGPR-(iPGPR) bacteria residing inside plant cells, producing nodules and being localized inside those specialized structures. These include a wide range of soil bacteria forming less formal associations than the rhizobia-legume symbiosis; endophytes may stimulate plant growth, directly or indirectly and include the rhizobia. Soil bacteria in the genera *Rhizobium*,

Bradyrhizobium, *Sinorhizobium*, *Azorhizobium*, *Mesorhizobium* and *Allorhizobium*, belonging to the family Rhizobiaceae, invade plant root systems and form root nodules (Martinez-Romero and Wang, 2000). Collectively they are often referred to as rhizobia. These iPGPR are mostly Gram-negative and rod-shaped, with a lower proportion being Gram-positive rods, cocci and pleomorphic forms. The primary mechanism by which rhizobia increase plant growth is N₂-fixation.

Root Colonization by PGPR

Successful colonization and persistence in the plant rhizosphere are required for PGPR to exert their beneficial effect on plants (Elliott and Lynch, 1984). Rhizosphere colonization is also considered to be a crucial step in the application of microorganisms for beneficial purposes such as biofertilization, phytostimulation, biocontrol and phytoremediation (Lugtenberg *et al.*, 2001). Root colonization, which is a complex process, is under the influence of various parameters such as bacterial traits, root exudates, biotic and abiotic factors (Benizri *et al.*, 2001). The method chosen for studying the traits associated with root colonization depends on the objective and different approaches and techniques have been used to quantify and identify inoculated strains on the host plant (Lugtenberg *et al.*, 2001). The major rhizobacterial genera include free living and symbiotic bacterial species belonging to *Acetobacter*, *Arthrobacter*, *Azospirillum*, *Azotobacter*, *Bacillus*, *flavobacter*, *Pseudomonas*, *Proteus*, *Rhizobium*, *Serratia*, *Xanthomonas* and other (Glick, 1995).

MODE OF ACTION OF PGPR AS BIOFERTILIZERS

PGPR inoculant currently commercialized as novel solution for plant growth enhancements through direct, indirect mechanism and combination of mode of action.

The direct mechanisms of plant growth promotion may involve the synthesis of substances by the bacterium or facilitation of the uptake of nutrients from the environment (Glick *et al.*, 1999). The direct growth promoting mechanisms are as follows (1) Biological nitrogen fixation (2) Increasing the availability of nutrients in the rhizosphere (solubilization of phosphorus, facilitated absorption of iron by production of siderophores) and (3) inducing of phytohormones production such as auxins, cytokinins, gibberellins (Kloepper *et al.*, 1989; Glick, 1995; Glick *et al.*, 1999; Vessey, 2003; Patten and Glick, 2002). The indirect mechanism of plant growth occurs when PGPR lessen or prevent the deleterious effects of plant pathogens on plants by production of inhibitory substances or by increasing the natural resistance of the host (Handelsman and Stabb, 1996; Nehl *et al.*, 1996; Cartieaux *et al.*, 2003). The indirect mechanisms of plant growth promotion by PGPR include (1) antibiotic production (2) depletion of iron from the rhizosphere (3) synthesis of antifungal metabolites (4) production of fungal cell wall lysing enzymes (5) competition for sites on roots and (6) induced systemic resistance (Weller and Cook, 1986; Dunne *et al.*, 1993; Kloepper *et al.*, 1988; Liu *et al.*, 1995; Glick *et al.*, 1999).

Direct Mechanism

Biological Nitrogen Fixation

Symbiotic Nitrogen Fixation

Legume and their rhizobia are often introduced to agricultural ecosystems to improve soil fertility and farming systems flexibility (Brockwell and Bottomley, 1995; Sessitsch *et al.*, 2002). Nitrogen is known to be an essential nutrient for plant growth and development. Intensive farming practices that achieve high yield require chemical fertilizers, which are not

costly but may also create environmental problems. The extensive use of chemical fertilizers in agriculture is currently under debate due to environmental concern and fear for consumer health. Consequently, there has recently been growing level of interest in environmentally and friendly sustainable agricultural practices and organic farming system (Rigby and Caceres, 2001; Lee and Song, 2007). Increasing and extending the role of biofertilizers such as rhizobium would reduce the need for chemical fertilizers and decrease adverse environmental effects. Therefore, in the development and implementations of sustainable agriculture techniques, biofertilization is great importance in alleviating environmental pollution and the deterioration of nature (Elkoca *et al.*, 2008).

Rhizobium symbiosis with legumes species is of special importance, producing 50% of 175 million tons of total biological nitrogen fixation annually worldwide (Sarioglu *et al.*, 1993). Chickpea and *Rhizobium leguminosarum* subsp. *ciceri* association annually produce up to 176 kg N ha⁻¹ depending on cultivar, bacterial strain and environmental factors (Rupela and Saxena, 1987; Beck *et al.*, 1991, Yadav *et al.*, 2010). However, Rhizobium species producing nodules in chickpea are specific only to this species and thus inoculation with effective strains is advised in soils with no or weak bacterial presence (Rupela and Saxena, 1987).

Nitrogen is one of the most common nutrients required for plant growth and productivity as it forms an integral part of proteins, nucleic acids and other essential biomolecules (Böckman, 1997). More than 80% of nitrogen is present in the atmosphere, but is unavailable to plants. It needs to be converted into ammonia, a form available to plants and other eukaryotes. Biological nitrogen fixation involves the conversion of nitrogen to ammonia by microorganisms using a complex enzyme system identified as nitrogenase (Kim and Rees, 1994). Biological nitrogen fixation fixes about 60% of the earth's available nitrogen and represents an economically beneficial and environmentally sound alternative to chemical fertilizers (Ladha and Kundu, 1997). Plant growth-promoting rhizobacteria that fix nitrogen in non-leguminous plants are diazotrophs that form a non-obligate interaction with the host (Glick *et al.*, 1999). The nitrogen fixation in chickpea is performed by two species of genus *Mesorhizobium*, *M. ciceri* (Nour *et al.*, 1994) and *M. mediterraneum* (Nour *et al.*, 1995). The symbiosis between chickpea and its specific rhizobia has been recently studied in several countries-Tunisia (Aouani *et al.*, 2001), Morocco (Maatallah *et al.*, 2002), Canada (Kyei-Boahen *et al.*, 2002), Turkey (Icgen *et al.*, 2002) and Portugal (Laranjo *et al.*, 2001, 2002) because of the promising agricultural usefulness of this crop as a grain legume for human and animal nutrition and because of the interesting extreme host specificity of its rhizobia. The most studied and longest exploited PGPR are the rhizobia (including the *Allorhizobium*, *Azorhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Rhizobium* and *Sinorhizobium*) for their ability to fix N₂ in their legume hosts. The root nodules bacteria in the α -Proteobacteria are contained in five families: Rhizobiaceae, Phyllobacteriaceae, Bradyrhizobiaceae, Hyphomicrobiaceae and Methylobacteriaceae as defined by their 16S rDNA sequence analysis (Sawada *et al.*, 2003). Commercial rhizobia inoculants for use on legume crops were first introduced in the 1890s (Fred *et al.*, 1932).

The rhizobial species involved in symbiotic association are *Bradyrhizobium japonicum* (Guerinot and Chelm, 1984), *Rhizobium ciceri* (Nour *et al.*, 1994), *Rhizobium etli* (Segovia *et al.*, 1993), *Rhizobium galegae* (Lindstrom, 1989), *Rhizobium gallicum* (Amarger *et al.*, 1997), *Rhizobium fredii* (Scholla and Elkan, 1984), *Sinorhizobium fredii* (Chen *et al.*, 1988), *Sinorhizobium medicae* (Rome *et al.*, 1996), *Sinorhizobium arboris* (Nick *et al.*, 1999), *Mesorhizobium chacoense* (Velazquez *et al.*, 2001), *Mesorhizobium pluriflorum* (De Lajudie *et al.*, 1998a), *Azorhizobium caulinodans* (Dreyfus *et al.*, 1988) and *Allorhizobium undicola* (De Lajudie *et al.*, 1998b). Understanding of the *Frankia* sp.

symbioses expanded rapidly in the 1970s and 1980s and their ability to produce nodules on sea buckthorn (*Hippophae rhamnoides*) (Oremus, 1980) solidified a new line of dynamic research. Jarvis *et al.* (1997) described the genus *Mesorhizobium* (rhizobia phylogenetically intermediate between the genera *Bradyrhizobium* and *Rhizobium*) to include root nodule bacteria that had considerable phenotypic and genotypic differences to the other root nodules bacteria genera. In case of lentil, seed treatment with *Rhizobium leguminosarum*, increased seedling height, root nodules mass and shoot biomass. Similarly, nodulation was found to be improved in peanut by an array of *Rhizobium* species (Dey *et al.*, 2004). Similarly data were reported by Yanni (1992), the response of lentil, chickpea and pea upon with *Rhizobium*.

Non-Symbiotic Nitrogen Fixation

Azotobacter species (*Azotobacter chroococcum* and *Azotobacter vinelandii*) and *Azospirillum* species (Mirza *et al.*, 2001) are free living, hetrotrophic diazotrophs that fix nitrogen in non leguminous plant such as wheat, rice and others crops. Yield of rice in field trials increased significantly 20% with applications of *Azotobacter* (Yanni and El-Fattah, 1999). One of the best studied diazotrophs for nitrogen fixation is *Azospirillum* sp. isolated from nitrogen-poor soils by Beijernick in 1925 (Holguin *et al.*, 1999). Members of this bacterial genus are capable of fixing atmospheric nitrogen and of promoting plant growth. A mixed inoculum of *Staphylococcus* and *Azospirillum* promoted the nitrogen fixing activity of *Azospirillum* (Holguin and Bashan, 1997). Combined inoculation of *A. brasilense* with *Pseudomonas striata* significantly increased grain yield, nitrogen and phosphorus uptake of sorghum (Alagawadi and Gaur, 1992). Oliveira *et al.* (1997) demonstrated that co-inoculation of clover with a mixture of *A. brasilense* Sp7 and *Rhizobium* sp. resulted in increased acetylene reduction, nodulation and shoot dry weight of plants. Endophytic bacteria that colonize the interiors of plant tissues such as roots, stem and leaves and are able to fix nitrogen are also found to be beneficial for plant growth (James *et al.*, 1997). Some experiments have involved the colonization of para-nodules on associations involving wheat and various diazotrophs (Chen *et al.*, 1993; Gantar and Elhai, 1999; Tchan *et al.*, 1991; Youssef *et al.*, 1998), *Azorhizobium caulinodans* and rice (Christiansen-Weniger, 1996), *Azotobacter nigricans* and rape (Koval'skaya *et al.*, 2001) and *Azospirillum* sp. and maize (Christiansen-Weniger and Vanderleyden, 1994).

Increasing the Availability of Nutrients in the Rhizosphere

There is ample evidence that the mode of action of many PGPR is by increasing the availability of nutrients for the plant in the rhizosphere (Glick, 1995; Vessey, 2003). The method by which these increases take place involve solubilization of unavailable forms of nutrients, siderophore production and ammonia production which helps facilitate the transport of certain nutrients.

Solubilization of Phosphorus

Phosphorus is second only to nitrogen in mineral nutrients most commonly limiting the growth of terrestrial plants. Ironically, soils may have large reserves of total P, but the amounts available to plants is usually a tiny proportion of this total (Stevenson and Cole, 1999). Indian soils are normally deficient in available phosphorus even though the bound component may be sufficient abundant (Johri *et al.*, 2003). Therefore, the use of PSM (phosphate solubilizing microorganism) is very common. However, immobilization of added




Fig. 1: Phosphate solubilizing (a) bacteria *Pseudomonas aeruginosa* strain BHUPSB01 (accession number GU124822) and (b) *Burkholderia cepacia* strain BHUPSB03 (accession number GU124830) strain showing halo zone around bacterial colony on Pikovskaya agar media

phosphorus in soils is a serious problem (Gaind *et al.*, 2000). The low availability of P to plants is because the vast majority of soil P is found in insoluble forms and plants can only absorb P in two soluble forms, the monobasic (H_2PO_4^-) and the dibasic (HPO_4^{2-}) ions. *In vitro* phosphate solubilizing bacteria *Pseudomonas aeruginosa* strain BHUPSB01 (accession number GU124822), *P. aeruginosa* strain BHUPSB02 (GU124826) and *Pseudomonas putida* strain BHUPSB04 (accession number GU124834), *Burkholderia cepacia* BHUPSB03 (GU124830), *Paenibacillus polymyxa* BHUPSB16 (GU124833), *P. polymyxa* BHUPSB17 (GU124838) and *Bacillus subtilis* BHUPSB13 (GU124812 showed halo zone around bacterial colony on Pikovskaya agar media (Gaur, 1990). This halo zone showed to solubilization of tricalcium phosphate in soluble form due to production of organic acid (Fig. 1a, b).

The solubilization of P in the rhizosphere is the most common mode of action implicated in PGPR that increase nutrient availability to host plants (Richardson, 2001). Examples of recently studied associations include *Azotobacter chroococcum* and wheat (Kumar and Narula, 1999), *Bacillus circulans* and *Cladosporium herbarum* and wheat (Singh and Kapoor, 1999), *Bacillus* sp. and five crop species (Pal, 1998), *Enterobacter agglomerans* and tomato (Kim and McDonald, 1998), *Pseudomonas chlororaphis* and *P. putida* and soybean (Cattelan *et al.*, 1999), *Rhizobium* sp. and *Bradyrhizobium japonicum* and radish (Antoun *et al.*, 1998) and *Rhizobium leguminosarum* bv. *Phaseoli* and maize (Chabot *et al.*, 1998). Phosphate-solubilizing bacteria are common in rhizospheres (e.g., Nautiyal *et al.*, 2000; Vazquez *et al.*, 2000). However, the ability to solubilize P by no means indicates that a rhizospheric bacterium will constitute a PGPR. For example, Cattelan *et al.* (1999) found only two of five rhizospheric isolates positive for P solubilization actually had a positive effect on soybean seedling growth. Likewise, not all P-solubilizing PGPR increase plant growth by increasing P availability to the hosts. For example, Freitas *et al.* (1997) found a number of P-solubilizing *Bacillus* sp. isolates and a *Xanthomonas maltophilia* isolate from canola (*Brassica napus* L.) rhizosphere which had positive effects on plant growth, but no effects on P content of the host plants.

Phosphate solubilizing bacteria are also known to increase phosphorus uptake resulting in better growth and higher yield of crop plants (Alagawadi and Gaur, 1988; Gaur *et al.*, 2004; Verma *et al.*, 2010). These microorganisms are bacteria and fungi that inhabitant the rhizosphere (Azcon *et al.*, 1976; Bowen and Rovira, 1999). Most soil bacteria can solubilize insoluble phosphates; particularly active are those that belong to the genera *Pseudomonas*, *Enterobacter* and *Bacillus* as well as some soil fungi, *Penicillium* and *Aspergillus* (Domey and Lipmann, 1988; Pargiri and Bezbaruah, 1990; Rao, 1992; Rokade and Patil, 1993; Whitelaw, 2000; Yadav *et al.*, 2010). Some researchers prefer to use fungal P-solubilizers arguing that bacterial strains can lost their ability to solubilize P after several cycles of *in vitro* culture (Whitelaw, 2000), but this point is quite controversial.

Facilitated Absorption of Iron by Production of Siderophores

Siderophores are low-molecular weight iron-binding molecules that are synthesized by many microorganisms under low-iron conditions (Neilands, 1981). Microbial siderophores may stimulate plant growth directly by increasing the availability of iron in the soil surrounding the roots (Kloepper *et al.*, 1980). Marschner and Römheld (1994) reported that plants may also utilize siderophores synthesized by microorganisms colonizing the rhizosphere; this would be a source of soluble iron for the host plant. Plants such as sorghum, oats, peanut, cotton, cucumber and sunflower demonstrated the ability to use radiolabelled microbial siderophores as a sole source of iron (Crowley *et al.*, 1988; Bar-Ness *et al.*, 1991; Cline *et al.*, 1984; Jurkevitch *et al.*, 1986; Wang *et al.*, 1993). Growth of cucumber in the presence of microbial siderophores resulted in increased plant biomass and chlorophyll content (Ismande, 1998). Uptake of microbial siderophores by plants has been attributed to microorganisms living.

Iron is an essential nutrient of plants, but it is relatively insoluble in soil solutions. Plant roots prefer to absorb iron as the more reduced ferrous (Fe^{2+}) ion, but the ferric (Fe^{3+}) ion is more common in well aerated soil although it is easily precipitated in iron-oxide forms (Salisbury and Ross, 1992). However, there is a controversy as the significance of bacterial Fe^{3+} siderophore uptake to the iron nutrition of plants. While some believe that there contribution of these siderophores to the overall iron requirements of plants is small (Glick, 1995), other suggest an important role (Duijff *et al.*, 1994), even a vital role, especially in calcareous soils (Masalha *et al.*, 2000). Bar-Ness *et al.* (1992), who had earlier supported the concept of bacterial siderophore uptake by plants (Bar-Ness *et al.*, 1991), concluded that two bacterial siderophores (pseudobactin and ferrioxamine B) were inefficient as iron sources for plants and that rhizospheric siderophore-producing bacteria can be in competition with the plant for iron. In fact, the vast majority of research on microbial siderophores in the rhizosphere is associated with their biocontrol activities due to their competitive effects with plant pathogens (Hiifte *et al.*, 1994).

Production of Phytohormones

One of the direct mechanisms by which PGPR promote plant growth is by production of plant growth regulators or phytohormones (Glick, 1995). Frankenberger and Arshad (1995) have discussed in detail the role of auxins, cytokinins, gibberellins, ethylene and Abscisic Acids (ABA) which, when applied to plants, help in increasing plant yield and growth. Microbial production of individual phytohormones such as auxins and cytokinins has been reviewed by various authors over the last 20 years (Pilet *et al.*, 1979; Hartmann *et al.*, 1983; Fallik *et al.*, 1988; Barbieri and Galli, 1993; Patten and Glick, 1996, 2002).

Auxins

The production of an active substance by the fungi *Rhizopus suinus* and *Absidia ramosa* was identified to be auxin by Thimann (1935), following induction of curvature in Avena as demonstrated by Nielsen (Frankenberger and Arshad, 1995). Various authors have identified the production of indole-3-acetic acid by microorganisms in the presence of the precursor tryptophan or peptone. Some of the plant responses to auxin are as follows: (1) cell enlargement (2) cell division (3) root initiation (4) root growth inhibition (5) increased growth rate (6) phototropism (7) geotropism and (8) apical dominance (Frankenberger and Arshad, 1995). Eighty percent of microorganisms isolated from the rhizosphere of various crops have the ability to produce auxins as secondary metabolites (Kampert *et al.*, 1975; Loper and Scroth, 1986). Bacteria belonging to the genera *Azospirillum*, *Pseudomonas*, *Xanthomonas* and *Rhizobium* as well as *Alcaligenes faecalis*, *Enterobacter cloacae*, *Acetobacter diazotrophicus* and *Bradyrhizobium japonicum* have been shown to produce auxins which help in stimulating plant growth (Patten and Glick, 1996). Various metabolic pathways such as (1) indole-3-acetamide pathway (2) indole-3-pyruvic acid pathway (3) tryptophan side chain pathway (4) tryptamine pathway and (5) indole-3-acetonitrile pathway are involved in the production of IAA. Phytopathogens such as *Agrobacterium tumefaciens*, *A. rhizogenes* and *P. syringae* pv. *savastanoi* synthesize IAA via the indole-3-acetamide pathway (Liu *et al.*, 1982; Offringa *et al.*, 1986). Synthesis of IAA by *Rhizobium* sp. in the presence and absence of tryptophan has also been demonstrated (Kittell *et al.*, 1989; Verma *et al.*, 2010). Bent *et al.* (2001) reported that the production of indole compounds by three different strains, *Paenibacillus polymyxa* L6, *P. polymyxa* Pw-2 and *Pseudomonas fluorescens* M20 increased in concentration with increasing concentrations of tryptophan (0-200 mg mL⁻¹) at different times. Reports by Asghar *et al.* (2002) showed that PGPR strains produced 24.6 µg mL⁻¹ of auxins in the presence of the precursor L-tryptophan in the medium, which was 184-fold more than that without L-tryptophan. Example of phytohormones producing bacteria under *in vitro* condition (Table 1). The isolates of *Bacillus*, *Pseudomonas* and *Azotobacter* produced IAA, whereas only 85.7% of *Rhizobium* was able to produce IAA. Production of ammonia was commonly detected in the isolates of *Bacillus* (95.0%) followed by *Pseudomonas* (94.2%), *Rhizobium* (74.2%) and *Azotobacter* (45.0%) (Joseph *et al.*, 2007). Higher level of IAA production by *Pseudomonas* was recorded by other workers (Xie *et al.*, 1996). Ahmad *et al.* (2005) has reported that bacterial isolates were tested for the production of indole acetic acid (IAA) in a medium with 0, 1, 2 and 5 mg mL⁻¹ of tryptophan. Production of IAA in fluorescent *Pseudomonas* isolates increased with an increase in tryptophan concentration from 1 to 5 mg mL⁻¹ in the majority of isolates. IAA production in *Azotobacter* isolates has been reported by other researchers (Gonzalez Lopez and Vela, 1981; Nieto and Frankenberger, 1989; Khalid *et al.*, 2001). IAA production has reported in *Azospirillum* sp. under *in vitro* condition (Dobbelaere *et al.*, 2001; Lambrecht *et al.*, 2000) (Table 1).

Other Phytohormones

Production of other phytohormones by biofertilizing-PGPR has been identified, but not nearly to the same extent as bacteria which produce IAA. Cytokinins are a class of phytohormones which are known to promote cell divisions, cell enlargement and tissue expansion in certain plant parts (Salisbury, 1994). Researchers have recently begun to identify cytokinin production by PGPR. Gibberellins (gibberellic acid; GA) are a class of phytohormones most commonly associated with modifying plant morphology by the extension of plant tissue, particularly stem tissue (Salisbury, 1994). Evidence of GA production by PGPR is rare, however Gutierrez-Manero *et al.* (2001) provide evidence that

Table 1: *In vitro* production of plant growth regulators by rhizobacteria

PGPR	PGR	References
<i>Arthrobacter mysoarens</i> 7, <i>Flavobacterium</i> sp. L30, <i>Klebsiella</i> CIAM 880 <i>Azotobacter</i> sp.,	Indole-3-acetic acid, ethylene	Pishchik <i>et al.</i> (2002)
<i>Azotobacter chroococcum</i> <i>Azospirillum</i> sp. <i>Azospirillum</i> sp. <i>Azospirillum</i> sp. <i>Bacillus licheniformis</i> <i>B. licheniformis</i> <i>B. pumilus</i> <i>B. subtilis</i> <i>Bacillus megaterium</i> BHUPSB14 <i>Pseudomonas</i> sp. <i>Pseudomonas aeruginosa</i> <i>Pseudomonas fluorescens</i> <i>Pseudomonas fluorescens</i> G20-18	Indole-3-acetic acid Gibberellic acid Indole-3-acetic acid Indole-3-acetic acid Ethylene Physiologically active gibberellins Physiologically active gibberellins Ethylene Indole-3-acetic acid Auxins. Ethylene Ethylene Isopentyl adenosine, <i>t</i> -zeatin ribose, dihydrozeatin riboside	Zahir <i>et al.</i> (1998a, b 2000), Khalid <i>et al.</i> (2001) Verma <i>et al.</i> (2010) Lucangeli and Botlini (1997) Dobbelaere <i>et al.</i> (2001) Lambrecht <i>et al.</i> (2000) Fukudah <i>et al.</i> (1989) Gutierrez-Manero <i>et al.</i> (2001) Gutierrez-Manero <i>et al.</i> (2001) Mansouri and Bunch 1989) Verma <i>et al.</i> (2010) Pal <i>et al.</i> (2000) Mansouri and Buncli (1989) Swanson <i>et al.</i> (1979) Garcia de Salamone <i>et al.</i> (2001)
<i>Pseudomonas fluorescens</i> BHUPSB06 <i>Pseudomonas putida</i> <i>Pseudomonas syringae</i> <i>Pseudomonas tabaci</i> <i>Rhizobacterial isolates</i> <i>Rhizobium</i> sp. <i>R. leguminosarum</i> <i>R. leguminosarum</i> <i>R. leguminosarum</i> <i>Rhizobium</i> sp.	Indole-3-acetic acid Ethylene Ethylene Ethylene Auxins Isopentyl adenine, isopentyl adenosine Indole-3-acetic acid Indole-3-acetic acid Isopentyl adenosine, zeatin, methylthiozeatin Indole-3-acetic acid	Verma <i>et al.</i> (2010) Pazout <i>et al.</i> (1981), Fukudah <i>et al.</i> (1989) Sato <i>et al.</i> (1997), Swanson <i>et al.</i> (1979) Swanson <i>et al.</i> (1979) Asghar <i>et al.</i> (2000, 2002) Sturtevant and Taller (1989) Badnoch-jones <i>et al.</i> (1982a), Dazzo <i>et al.</i> (2000) Wang <i>et al.</i> (1993) Taller and Slunevant (1991) Verma <i>et al.</i> (2010)

four different forms of GA are produced by *Bacillus pumilus* and *Bacillus licheniformis* (Table 1). Inoculation of alder (*Alnus glutinosa*) with these PGPR could effectively reverse a chemically induced inhibition of stem growth. Although this is not strong evidence of GA production being a common method of growth promotion by PGPR, it does suggest that it may have a role and indicates that more research in this area is warranted. Ethylene is the only gaseous phytohormone. It is also known as the 'wounding hormone' because its production in the plant can be induced by physical or chemical perturbation of plant tissues (Salisbury, 1994). Among its myriad of effects on plant growth and development, ethylene production can cause an inhibition of root growth. Glick *et al.* (1998) put forward the theory that the mode of action of some PGPR was the production of 1-aminocyclopropane-1-carboxylate (ACC) deaminase, an enzyme which could cleave ACC, the immediate precursor to ethylene in the biosynthetic pathway for ethylene in plants. They submitted that ACC deaminase activity would decrease ethylene production in the roots of host plants and result in root lengthening. Subsequently, Glick's research program has provided numerous empirical studies with wild-type and genetically modified PGPR to support their theory (Holguin and Glick, 2001; Li *et al.*, 2000; Mayak *et al.*, 1999; Saleh and Glick, 2001; Shah *et al.*, 1998; Wang *et al.*, 2000). In some cases, the growth promotion effects of ACC deaminase-producing PGPR appear to be best expressed in stressful situations such as flooded (Grichko and Glick, 2001) or heavy metal-contaminated soils (Burd *et al.*, 1998).

The Indirect Mechanism

The indirect mechanism of plant growth occurs when PGPR lessen or prevent the deleterious effects of plant pathogens on plants by production of inhibitory substances or

Table 2: Biological control by PGPR against certain diseases, pathogens and insects in different crops

Crop	Disease/pathogen/insect	PGPR	References
Barley	Powdery mildew	<i>B. subtilis</i>	Schobeck <i>et al.</i> (1980)
Beans	Halo blight	<i>Pseudomonas fluorescens</i> strain 97	Alstrom (1991)
	<i>Sclerotium rolfsii</i>	<i>Pseudomonas cepacia</i>	Fridlender <i>et al.</i> (1993)
Carnation	<i>Fusarium</i> wilt	<i>Pseudomonas</i> sp. (WCS 417r)	Van Peer <i>et al.</i> (1991)
Cotton	Damping off	<i>Pseudomonas fluorescens</i>	Howell and Stipanovic (1979, 1980)
	<i>Meloidogyne incognita</i> , <i>M. Arenaria</i>	<i>B. subtilis</i>	Sikoraa (1998)
	<i>Rhizoctonia solani</i>	<i>Pseudomonas cepacia</i>	Fridlender <i>et al.</i> (1993)
Cucumber	<i>Helicoverpa armigera</i>	<i>Pseudomonas gladioli</i>	Quingwen <i>et al.</i> (1998)
	Cucumber anthracnose	<i>Pseudomonas putida</i> (89B-27), <i>Serratia marcescens</i> (90-166)	Wei <i>et al.</i> (1991, 1996)
	<i>Pythium ultimum</i>	<i>Pseudomonas cepacia</i>	Fridlender <i>et al.</i> (1993)
	Bacterial wilt	<i>Pseudomonas putida</i> (89B-27), <i>S. marcescens</i> (90-166)	Kloepper <i>et al.</i> (1993)
	Bacterial angular leaf spot	<i>P. putida</i> (89B-27), <i>Flavomonas oryzihabitans</i> INR-5, <i>S. marcescens</i> (90-166), <i>Bacillus pumilus</i> (INR-7)	Kloepper <i>et al.</i> (1993)
	Cucumber mosaic virus	<i>P. putida</i> (89B-27), <i>S. marcescens</i> (90-166)	Raupach <i>et al.</i> (1996)
	Striped Cucumber beetle	<i>P. putida</i> (89B-27), <i>Flavomonas</i> <i>oryzihabitans</i> INR-5, <i>Pseudomonas</i> sp.	Zehnder <i>et al.</i> (1997)
Green gram	<i>Aspergillus</i> sp., <i>Curvularia</i> sp., <i>Fusarium oxysporum</i> , <i>Rhizoctonia solani</i>		Sindhu <i>et al.</i> (1999)
	Mung bean Root rot. Root knot	<i>P. aeruginosa</i> , <i>B. subtilis</i>	Siddiqui <i>et al.</i> (2001a, b)
	Rice Rice sheath blight	<i>Streptomyces</i> spp. and <i>Bacillus cereus</i> in combination with <i>Ps. fluorescens</i> and <i>Burkholderia</i>	Sung and Chung (1997)
Rice	Rice sheath blight	Combination of <i>P. fluorescens</i> strains Pfl and Fp7	Nandakumar (1998)
	<i>Rhizoctonia solani</i> (sheath blight pathogen)	<i>P. fluorescens</i> Strains Pf 1 and Fp7	Vidhayasekaran and Muthamilan (1999)
	Blue mold	<i>S. marcescens</i> 90-1 16, <i>B. pumilus</i> SE 34, <i>Ps. fluorescens</i> 898-6 1, <i>B. pumilus</i> T4, <i>B. pasteurii</i> C-9	Zhang <i>et al.</i> (2002)
	Tomato	<i>B. subtilis</i> strain 1N 937b	
Tomato	Tomato mottle virus	<i>B. amylioliquifaciens</i> strain 1N 937a, <i>B. subtilis</i> strain 1N 937b	Murphy <i>et al.</i> (2000)
	Wheat Take till disease	<i>Bacillus pseudomonas</i> , <i>Penicillium</i> , <i>Beauveria</i> , <i>Rhodococcus</i> Mixture of <i>Pseudomonas</i> sp.	Renwick <i>et al.</i> (1991) Pierson and Weller (1994)

by increasing the natural resistance of the host. Control of phytopathogenic microorganism by releasing siderophores, B-1,3-glucanase, chitinases, antibiotics, fluorescent pigment and cynide (Scher and Baker, 1982; Voisard *et al.*, 1989; Catellan *et al.*, 1999; Pal *et al.*, 2001).

Many rhizobacteria including fluorescent *Pseudomonas* secrete a variety of antifungal molecule under in vitro and in situ conditions (Cook *et al.*, 1995). There are several reports in vitro antagonism of pathogenic fungi and field performance by bacteria recovered from the rhizosphere of plants in India (Johri *et al.*, 1997). Some important plant disease controlling bacteria have been depicted in Table 2. *Pseudomonas aeruginosa* strain BHUPSB01 (accession number GU124822), *P. aeruginosa* strain BHUPSB02 (GU124826) and *Pseudomonas putida* strain BHUPSB04 (accession number GU124834), *Burkholderia cepacia* BHUPSB03 (GU124830), *Paenibacillus polymyxa* BHUPSB16 (GU124833), *P. polymyxa* BHUPSB17 (GU124838) and *Bacillus subtilis* BHUPSB13 (GU124812) showed antagonistic against phytopathogenic fungi *Fusarium oxysporum* and *Rhizoctonia solani* (Fig. 2, 3).

The rhizosphere and root zone of tea (*Camellia sinensis*) is a good habitat for PGPR strains represented by *Bacillus*, *Proteus* and *Pseudomonas*, inhibitory to phytopathogenic fungi *in vitro*, viz., *Fusarium oxysporum* f. sp. *lycospersici*, *Fusarium oxysporum* f. sp.



Fig. 2: Antagonistic activity of *Pseudomonas aeruginosa* strain BHUPSB01 (accession number GU124822) against *Fusarium oxysporum*



Fig. 3: Antagonistic activity of *Pseudomonas putida* strain BHUPSB04 (accession number GU124834) against *Rhizoctonia solani*

ciceri, *F. udum*, *F. solani*, *F. moniliformae*, *F. semitectum*, *Fomes lamiensis* and *Ustilina zonata* (Kumar-Dileep and Bezbaruah, 1996; Kumar and Bezbaruah, 1997). Suppression of soil borne plant pathogens by siderophore producing pseudomonads was observed (Bakker *et al.*, 1986, 1987; Loper, 1988) and the wild type strain was more effective in suppressing disease compared to non-siderophore-producing mutants. *Pseudomonas* sp. strains MRS23 and CRP55b showed varying diameters of inhibition zones of the four pathogenic fungi, *Aspergillus* sp. *F. oxysporum*, *P. aphanidermatum* and *R. solani* in the plating media (Kings B, PDA and NB) (Goel *et al.*, 2002). *Pseudomonas* strains MRS23 and CRP55b inhibited the growth of pathogenic fungi, i.e., *Aspergillus* sp. *Fusarium oxysporum* f. sp. *ciceri* and *Rhizoctonia solani* under culture condition (Goel *et al.*, 2002). Like this, several species of pseudomonas, viz. *P. fluorescens*, *P. putida*, *P. cepacia* (*Burkholderia cepacia*) and *P. aeruginosa* isolated from plant tissue or soil have been reported as potential biocontrol agents of several phytopathogenic fungi (Duffy *et al.*, 1996; Goel *et al.*, 2000). Saikia *et al.* (2003, 2004, 2005, 2009) reported that *Pseudomonas fluorescens* I-94 (Pf1-94) and

P. fluorescens (Pf4-92) strains that were isolated from rhizosphere soil of chickpea showed Induced Systemic Resistance (ISR) against fusarium wilt of chickpea and charcoal rot (Srivastava *et al.*, 2001). Earlier studies indicated that shoot and root lengths were significantly increased in *P. fluorescens* treated chickpea plants and the reduction in disease severity was more pronounced when chemical inducers were applied with *P. fluorescens* (Saikia *et al.*, 2003). Root rot disease in mung bean has controlled by *Pseudomonas aeruginosa*, *Bacillus subtilis* (Siddidui *et al.*, 2001) (Table 2).

Siderophore production is an important feature for the suppression of plant pathogens and promotion of plant growth. Fluorescent siderophore production was observed as a mechanism of biocontrol of bacterial wilt disease in the fluorescent pseudomonads RBL 101 and RSI 125. Press *et al.* (2001) reported the catechol siderophore biosynthesis gene in *Serratia marcescens* 90-166 and associated with induced resistance in cucumber against anthracnose. Pseudomonads also produce two other siderophores, pyochelin and its precursor salicylic acid and pyochelin is thought to contribute to the protection of tomato plants from *Pythium* by *Pseudomonas aeruginosa* 7NSK2 (Buysens *et al.*, 1996). Howell and Stipanovic (1979, 1980) has been reported damping off disease in cotton controlled by *Pseudomonas fluorescens*. Different environmental factors can also influence the quantity of siderophores produced (Duffy and Défago, 1999).

EFFECT OF PGPR ON PLANT GROWTH AND YIELDS

Table 3 lists many of the PGPR which have been implicated as helper bacteria in various legume-rhizobia symbioses and co-inoculation with other plant growth promoting rhizobacteria for enhanced crop production. There is evidence for a number of modes of action for PGPR stimulation of legume-rhizobia symbioses, but the most commonly implicated mode is phytohormone-induced (usually IAA) stimulations of root growth (e.g., Molla *et al.*, 2001; Srinivasan *et al.*, 1996; Vessey and Buss, 2002). In this way, the stimulation of nodulation is most commonly an indirect effect; the PGPR stimulates root growth, which provides more sites for infection and nodulation. However, this is not always the case. Cattelan *et al.* (1999) found that a number of putative PGPR had positive effects on shoot and/or root growth in soybean and were positive for production of IAA or ACC deaminase, but these putative PGPR had no positive effects on nodulation. In fact, Cattelan *et al.* (1999) found several rhizospheric isolates which stimulated aspects of the soybean-bradyrhizobia symbiosis and which had β -glucoanase or cyanide production.

Phosphate-solubilizing *Bacillus* sp. stimulates plant growth through enhanced P nutrition (Elkoca *et al.*, 2008; Whitelaw *et al.*, 1997; Verma *et al.*, 2010) and increasing the uptake of N, P, K and Fe (Biswas *et al.*, 2000). Some PGPR strains, from a range of genera, enhance legume growth, nodulation and nitrogen fixation, root and shoot biomass, nodule dry matter, nitrogenase activity and grain yield in chickpea (Parmar and Dadarwal, 1999; Hamaoui *et al.*, 2001; Sindhu *et al.*, 2002a; Zaidi *et al.*, 2003; Gull *et al.*, 2004). Combined inoculation of *Rhizobium* with *Pseudomonas striata* or *Bacillus* and with *Bacillus megaterium* (Elkoca *et al.*, 2008) have shown increased dry matter, grain yield and phosphorus uptake significantly over the uninoculated control in legumes. Interactions between these PGPR with *Rhizobium* may be antagonistic or synergistic and the beneficial effects of such interactions could be exploited for economic gain (Dubey, 1996; Glick, 1995). Many different N₂-fixing bacteria, including symbionts, such as root nodulating *Rhizobium* sp. (Bohlool *et al.*, 1992) and different free-living rhizobacteria, such as *Azospirillum*, *Azotobacter*, *Bacillus*, *Enterobacter*, *Pseudomonas*, *Serratia* and *Streptomyces* (Kloepper

Table 3: Co-inoculation of Legumes with PGPR and *Rhizobium*

Crop	Rhizobium	Coinoculating PGPR	Plant responses to inoculation	Reference
Alfalfa	<i>R. meliloti</i>	<i>Pseudomonas</i>	Upon co-inoculation, a significant increase in plant growth, nitrogenase activity, nodule total nodule weight and total plant nitrogen was reported	Knight and Langston-Unkefer (1988)
Bean	<i>R. phaseoli</i>	<i>Pseudomonas putida</i>	Inoculation with <i>P. putida</i> markedly increased nodulation compared to a <i>R. phaseoli</i> control. Also, 2-ketogluconic acid, a pflO5pbate-solubilizing compound, was detected in <i>P. putida</i>	Grimes and Mount (1984)
Chickpea	Indigenous <i>Rhizobium</i>	<i>A. brasilense</i>	Inoculation with <i>A. brasilense</i> has the potential to prevent or at least diminish the effects: caused by salinity stress in greenhouse experiments with chickpeas irrigated with saline water	Hamaoui <i>et al.</i> (2001)
	<i>Mesorhizobium</i>	<i>Pseudomonas</i>	A significant increase in nodule weight and shoot biomass was observed when co-inoculated with <i>Mesorhizobium</i> and <i>Pseudomonas</i> in sterilized chillum jar conditions. In pot experiment, co-inoculation significantly increased root and shoot biomass	Sindhu <i>et al.</i> (2002a, b)
	<i>Rhizobium</i>	<i>Pseudomonas</i> , <i>Bacillus</i>	Upon co-inoculation, reduction in wilt incidence and an increase in nodulation were observed	
	<i>Rhizobium</i>	<i>Pseudomonas</i> , <i>Bacillus</i>	A significant increase in nodule weight, root and shoot biomass and total plant nitrogen were reported due to co-inoculation	Parmar and Dadarwal (1999)
Chickpeas, common beans	<i>Rhizobium</i> sp.	<i>P. fluorescens</i> <i>B. megaterium</i> <i>A. chroococcum</i>	A significant increase in nodule weight, root and shoot biomass and total plant nitrogen were reported due to co-inoculation	Verma <i>et al.</i> (2010)
	<i>Rhizobium</i>	<i>A. brasilense</i>	Inoculation with <i>A. brasilense</i> increased nodule dry weight, various plant growth parameters and seed yield of nodulated chickpeas in the field. In common bean, inoculation with <i>Rhizobium meliloti</i> and <i>R. tropici</i> increased seed yield while combined inoculation with <i>Rhizobium</i> and <i>A. brasilense</i> resulted in a further increase. Plants inoculated with <i>A. brasilense</i> alone did not differ in yield from the non-inoculated controls, despite a relative increase in shoot dry weight	Burdman <i>et al.</i> (1996)
	<i>Rhizobium</i>	<i>A. brasilense</i>	Inoculation with <i>A. brasilense</i> increased nodule dry weight, various plant growth parameters and seed yield of nodulated chickpeas in the field. In common bean, inoculation with <i>Rhizobium meliloti</i> and <i>R. tropici</i> increased seed yield while combined inoculation with <i>Rhizobium</i> and <i>A. brasilense</i> resulted in a further increase. Plants inoculated with <i>A. brasilense</i> alone did not differ in yield from the non-inoculated controls, despite a relative increase in shoot dry weight	Burdman <i>et al.</i> (1996)
Clover	<i>R. leguminosarum</i> sp. Strain bv. <i>trifolii</i> 2A	<i>Pseudomonas</i> 267 and R.	Co-inoculation of clover plants significantly increased shoot weight and nodule weight in comparison with control plants infected only with <i>R. leguminosarum</i> bv. <i>trifolii</i>	Derylo and Skorupska (1993)
Common bean	<i>R. phaseoli</i>	<i>P. putida</i>	Co-inoculation caused a marked increase in nodulation. Co-inoculation promoted root hair formation and an increase in secretion of the nod gene induced flavonoids resulting in greater numbers of nodules	Grimes and Mount (1984), Burdman <i>et al.</i> (1996)
	<i>Rhizobium</i>	<i>A. brasilense</i>		
Green gram	<i>Bradyrhizobium</i> sp. (Vigna) strains Cog 15 and S24	<i>Bacillus</i>	Co-inoculants of the PGPR. <i>Bacillus</i> and <i>Bradyrhizobium</i> strains failed to show any conclusive influence on nodulation and ARA at 50 days of plant growth. <i>Bacillus</i> had a direct effect on shoot biomass development, N-content and grain yield of green gram when co-inoculated with <i>Bradyrhizobium</i> strain S24 at 50 and 80 days of plant growth. Also, single inoculation of <i>Bacillus</i> isolates significantly increased grain yield over the non-inoculated control	Gupta <i>et al.</i> (1998)
	<i>Bradyrhizobium</i>	<i>Bacillus</i>	Co-inoculation enhanced nodulation and plant growth of green gram.	Sindhu <i>et al.</i> (2002b)
	<i>Bradyrhizobium</i>	<i>Pseudomonas</i>	Co-inoculation resulted in significant increases in nodule weight, plant dry weight and total plant N compared to inoculation with <i>Bradyrhizobium</i> alone	Sindhu <i>et al.</i> (1999)
Natural pasture	Indigenous <i>Rhizobium</i>	<i>A. brasilense</i>	Inoculation with <i>A. brasilense</i> showed a similar effect to that of phosphate fertilization.	Itzigsohn <i>et al.</i> (2000)

Table 3: Continued

Crop	Rhizobium	Coinoculating		Reference
		PGRP	Plant responses to inoculation	
Pea	<i>R. leguminosarum</i>	<i>Pseudomonas</i>	Numbers of nodules were greater in co-inoculated plants than with <i>R. leguminosarum</i> inoculation alone	Bolton <i>et al.</i> (1990)
Peas, Chick peas, Vetch	<i>Rhizobium</i>	<i>A. brasilense</i>	Co-inoculation significantly increased seed yield, but did not affect dry matter yield of garden peas and chickpeas. Vetch co-inoculation significantly increased N ₂ -fixation, nitrogen content and dry matter yield	Sarig <i>et al.</i> (1986)
Soybean (1998)	<i>B. japonicum</i>	<i>S. liquefaciens</i>	Nodulation, nitrogen fixation, protein and N yield were increased by co- inoculation	Dashti <i>et al.</i>
		2-68, <i>S. Protea maculans</i> I-102		
	<i>B. japonicum</i>	<i>P. fluorescens</i>	Co-inoculation increased colonization of <i>Bradyrhizobium japonicum</i> on soybean roots, nodule number and the acetylene reduction assay	Chebotar <i>et al.</i> (2001)
	<i>B. japonicum</i>	<i>Azospirillum</i>	Co-inoculation promoted nodulation, nitrogenase activity and plant growth	Iruthayathas <i>et al.</i> (1983), Polonenko <i>et al.</i> (1987)
	<i>B. japonicum</i>	<i>S. liquefaciens</i>	Co-inoculation increased grain yield, grain protein	Dashti <i>et al.</i>
(1997)		2-68, <i>S. protea maculans</i> I-102	and total plant protein production under short season conditions	
	<i>B. japonicum</i>	<i>P. putida</i> , <i>P. fluorescens</i> , <i>Aeromonas hydrophila</i>	Co-inoculation promoted significant increase in the weight and number of nodules. Several strains increased the dry weight of shoots and roots when inoculated with <i>B. japonicum</i> , but these effects did not correlate with changes in number and weight of nodules	Polonenko <i>et al.</i> (1987)
White clover	<i>R. leguminosarum</i> bv. <i>trifolii</i>	<i>Azospirillum lipoferum</i>	Co-inoculation enhanced the number of nodules by 2-3 times, from 5 to 20 days after inoculation. Acetylene reduction activity was also increased by 2.3- to 2.7-fold at 20 days after inoculation	Tchebotar <i>et al.</i> (1998)

and Beauchamp, 1992; Hiipte *et al.*, 1994; Cakmakci *et al.*, 1999), have already been described. When used as seed inoculants, some of these free-living N₂-fixing bacteria show beneficial effects on plant growth and hence they are called Plant Growth-Promoting Rhizobacteria (PGPR) (Davison, 1988). *Bacillus* species used as biofertilizers may have direct effects on plant growth through the synthesis of plant growth hormones (Amer and Utkhede, 2000), N₂-fixation and solubilization of phosphate (Sahin *et al.*, 2004). Phosphate-solubilizing *Bacillus* sp. stimulates plant growth through enhanced P-nutrition (Whitelaw *et al.*, 1997), increasing the uptake of N, P, potassium (K) and iron (Fe) (Biswas *et al.*, 2000). Phosphorus biofertilizers could help increase the availability of phosphates accumulated in the soil and could enhance plant growth by increasing the efficiency of biological nitrogen fixation and the availability of Fe and zinc (Zn) through production of plant growth promoting substances (Kucey *et al.*, 1989). Indeed, trials with *Bacillus* species have shown yield increases in many crops (Greenough and Batson, 1989; Ozturk *et al.*, 2003).

Rhizobium or *Bradyrhizobium* are also widely used in agriculture for crop improvement because of their ability to fix atmospheric nitrogen. Some PGPR strains, from a range of genera, enhance legume growth, nodulation and nitrogen fixation when co-inoculated with rhizobia (Grimes and Mount, 1984; Petersen *et al.*, 1996). Co-inoculation studies with PGPR and *Rhizobium/Bradyrhizobium* sp. have been shown to increase root and shoot biomass, nodule dry matter, nitrogenase activity, N₂-fixation and grain yield in chickpea (Parmar and Dadarwal, 1999; Hamaoui *et al.*, 2001; Sindhu *et al.*, 2002a; Zaidi *et al.*, 2003; Gull *et al.*, 2004) and various legumes such as common bean (Grimes and Mount, 1984), green gram

(Sindhu *et al.*, 1999) and soybean (Dashti *et al.*, 1998). Furthermore, combined inoculations with N₂-fixing and P-solubilizing bacteria were more effective than single inoculation possibly by providing a more balanced nutrition for plants (Belimov *et al.*, 1995). Dual inoculations increased yields in many crops compared to single inoculations with N₂-fixing or P-solubilizing bacteria (Alagawadi and Gaur, 1992; Belimov *et al.*, 1995; Verma *et al.*, 2009).

Anjair *et al.* (2003) has been reported that *Pseudomonas aeruginosa* PNA1, an isolate from chickpea rhizosphere in India, protected pigeonpea and chickpea plants from *fusarium* wilt disease, which is caused by *Fusarium oxysporum* f.sp. *ciceri* and *Fusarium udum*. Inoculation with strain PNA1 significantly reduced the incidence of *fusarium* wilt in pigeonpea and chickpea on both susceptible and moderately tolerant genotypes. Valverde *et al.* (2006) has been reported effects of coinoculations with *Pseudomonas jessenii* PS06 (a) phosphate-solubilizing bacteria) and *Mesorhizobium ciceri* C-2/2 strains enhanced the growth and seed yield of chickpea under greenhouse and field conditions. Co-inoculation studies with PGPR and *Rhizobium/Bradyrhizobium* sp. have been shown to increased root and shoot biomass, nodule dry matter, nitrogenase activity, N₂-fixation and grain yield in chickpea (Sindhu *et al.*, 2002a, b; Zaidi *et al.*, 2003; Gull *et al.*, 2004; Elkoca *et al.*, 2008) and various legume such as green gram (Sindhu *et al.*, 1999) and pigeonpea (Tilak *et al.*, 2006). Mittal *et al.* (2008) has been reported the effect of six phosphate-solubilizing fungi (PSF, two strains of *Aspergillus awamori* and four of *Penicillium citrinum*) isolated from rhizosphere of various crops, was observed on the growth and seed production of chickpea plants (*Cicer arietinum* L. cv. GPF2) in pot experiments. Akhtar and Siddiqui (2009) has been reported that effects of Phosphate solubilizing microorganisms (*Glomus intraradices*, *Pseudomonas putida*, *P. alcaligenes*, *P. aeruginosa* (Pa28), *A. awamori*) and *Rhizobium* sp. was observed on the growth, nodulation yield and root-rot disease complex of chickpea under field condition. Inoculation of *Rhizobium* sp. caused a greater increase in growth and yield than *P. putida*, *P. aeruginosa* or *G. intraradices*. Hamaoui *et al.* (2001) also reported that inoculation with PGPR significantly enhanced nodulation by native rhizobia in chickpea and faba bean. Favorable effects of inoculation with N₂-fixing and P-solubilizing microorganisms and significant increase in nodulation and N₂-fixation, N% and total plant nitrogen of legume crops have been reported by many workers (Manjunatha and Devi, 1990; Petersen *et al.*, 1996; Khan *et al.*, 1997; Sindhu *et al.*, 2002a, b). Zaidi and Khan (2007) reported that the stimulatory effect of rhizotrophic microorganisms on growth, yield and nutrient uptake of chickpea was determined in a pot experiment using sterilised soil deficient in available phosphorus (P). Plant vigour, yield and nutrient uptake were significantly enhanced following inoculation with *Mesorhizobium ciceri* and the phosphate solubilising bacterium *Serratia* (T1) or phosphate solubilising fungus *Penicillium* (WF6). Akhtar and Siddiqui (2008) have been reported that *Glomus intraradices*, *Pseudomonas alcaligenes* and *Bacillus pumilus*: effective agents for the control of root-rot disease complex of chickpea (*Cicer arietinum* L.). Sivaramaiah *et al.* (2007) reported that Rhizobacteria belonging to *Bacillus* sp. were isolated from the rhizosphere of chickpea (*Cicer arietinum*). Two *Bacillus* strains CBS127 and CBS155 inhibited the growth of all the four pathogenic fungi (*Alternaria* sp., *Fusarium oxysporum*, *Pythium aphanidermatum* and *Rhizoctonia solani*) tested on nutrient agar medium plates *in vitro*. Ben-Romdhane *et al.* (2008) reported that selection of high nitrogen-fixing rhizobia (*Mesorhizobium ciceri* strain CMG 6 and *Mesorhizobium ciceri* strain CTM 226) nodulating chickpea (*Cicer arietinum* L.) for semi-arid Tunisia. Siddiqui and Mahmood (2001 b) have been reported the effects of rhizobacteria, i.e. *Pseudomonas fluorescens*, *Azotobacter chroococcum* and *Azospirillum brasilense*, alone and in combination with root symbionts; *Rhizobium* sp. and *Glomus mosseae*, on the growth of chickpea, *Cicer arietinum* and reproduction of *Meloidogyne javanica* were studied.

Rhizobium sp. with *P. fluorescens* was better for nodulation than the use of *Rhizobium* sp. alone or its use with *A. brasilense*. Root colonization by *G. mosseae* was better in the presence of *P. fluorescens* or *A. chroococcum* than the use of *G. mosseae* alone or its use with *A. brasilense*. *P. fluorescens* caused greater root colonization followed by *A. chroococcum* and *A. brasilense*. Verma *et al.* (2009) have reported that the maximum significant increase in nodule number, dry weight of nodule, root and shoot were recorded in coinoculation of *Mesorhizobium* sp. and *Pseudomonas fluorescens* followed by co-inoculation of *Mesorhizobium* sp., *Azotobacter chroococcum* and *Bacillus megaterium* over uninoculated control in both year of field study while nitrogen and phosphorus content increase in nodules, grain and straw. Verma *et al.* (2010) have reported that application of *Rhizobium* sp. BHURC01 and plant growth promoting rhizobacteria on nodulation, plant biomass and yields of chickpea (*Cicer arietinum* L.).

The higher grain yield 37.26 and 23.64% and straw yield 27.95 and 21.70% in first and second year of field study, respectively was increased significantly in co-inoculation of *Rhizobium* sp. BHURC01 with *P. fluorescens* followed by *B. megaterium* and *A. chroococcum* over uninoculated control.

CONCLUSION

The rhizosphere of a plant is a zone of intense microbial activity. Rhizobacteria that exert beneficial effect on plant growth and development or referred to as plant growth promoting rhizobacteria (PGPR). PGPR can affect plant growth directly or indirectly. Indirect promotion of plant growth occurs when introduced PGPR lessens or prevents deleterious effect of one and more phytopathogenic organisms in the rhizosphere. The direct promotion of plant growth by PGPR may include the production of plant growth regulators or facilitating the uptake of certain nutrient from the root environment. There is increasing evidence that rhizobia can promote the growth of non-legumes via direct and indirect mechanism. Further more, co-inoculation of legumes with rhizobia and PGPR is even more effective for improving nodulation and growth of legumes. Application of PGPR strains can provide an effective, economical and practical way of plant protection via disease suppression. PGPR strains mixture often show synergistic action in plant protection and growth promotion involving many mechanisms. Effective plant growth promoting rhizobacteria may be enhancing plant growth and grain yield in multiple crop production.

ACKNOWLEDGMENTS

We thank the Centre of Organic Farming, Ministry of Agriculture, Government of India, Ghaziabad, for providing a project fund to carry out research work on PGPR at Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh, India.

REFERENCES

- Ahmad, F., I. Ahmad and M.S. Khan, 2005. Indole acetic acid production by the indigenous isolates of *Azotobacter* and fluorescent *Pseudomonas* in the presence and absence of tryptophan. Turk. J. Biol., 29: 29-34.
- Akhtar, M.S. and Z.A. Siddiqui, 2008. Biocontrol of a root-rot disease complex of chickpea by *Glomus intraradices*, *Rhizobium* sp. and *Pseudomonas striata*. Crop Prot., 23: 410-417.
- Akhtar, M.S. and Z.A. Siddiqui, 2009. Effects of phosphate solubilizing microorganisms and *Rhizobium* sp. on the growth, nodulation, yield and root-rot disease complex of chickpea under field condition. African J. Biotechnol., 8: 3489-3496.

- Alagawadi, A.R. and A.C. Gaur, 1988. Associative effect of *Rhizobium* and phosphate solubilizing bacteria on the yield and nutrient uptake of chickpea. *Plant Soil*, 105: 241-246.
- Alagawadi, A.R. and A.C. Gaur, 1992. Inoculation of *Azospirillum brasilense* and phosphate solubilizing bacteria on yield of sorghum (*Sorghum bicolor* L.) Moench in dry land. *Trop. Agric.*, 69: 347-350.
- Alstrom, S., 1991. Induction of disease resistance in common bean susceptible to halo blight bacterial pathogen after seed bacterization with rhizosphere *Pseudomonads*. *J. Gen. Applied Microbiol.*, 37: 495-501.
- Amarger, N., V. Macheret and G. Laguerre, 1997. *Rhizobium gallicum* sp. nov. and *Rhizobium giardinii* sp. nov., from *Phaseolus vulgaris* nodules. *Int. J. Syst. Bacteriol.*, 47: 996-1006.
- Amer, G.A. and R.S. Utkhede, 2000. Development of formulations of biological agents for management of root rots of lettuce and cucumber. *Can. J. Microbiol.*, 46: 809-816.
- Andrade, G., E. Esteban, L. Velasco, J.L. Maria and E.J. Bedmar, 1997. Isolation and identification of N₂-fixing microorganisms from the rhizosphere of *Capparis spinosa* (L.). *Plant Soil*, 197: 19-23.
- Anjair, V., P. Cornelis and N. Koedom, 2003. Effect of genotype and root colonization in biological control of *Fusarium* wilts in pigeonpea and chickpea by *Pseudomonas aeruginosa* PNA1. *Can. J. Microbiol.*, 49: 85-91.
- Antoun, H., C.J. Beauchamp, N. Goussard, R. Chabot and R. Lalonde, 1998. Potential of *Rhizobium* and *Bradyrhizobium* species as plant growth promoting rhizobacteria on non-legumes: Effects of radishes (*Rhaphanus sativus* L.). *Plant Soil*, 204: 57-67.
- Aouani, M.E., R. Mhamdi, M. Jebara and N. Amarger, 2001. Characterization of rhizobia nodulating chickpea in Tunisia. *Agronomie*, 21: 577-581.
- Asgar, H.N., Z.A. Zahir, A. Khalid and M. Arshad, 2000. Assessment of auxine production from rhizobacteria isolated from different varieties of rapeseed. *Pak. J. Agric. Sci.*, 37: 101-104.
- Asgar, H.N., Z.A. Zahir, M. Arshad and A. Khaliq, 2002. Relationship between *in vitro* production of auxins by rhizobacteria and their growth promoting activities in *Brassica juncea* L. *Bio. Fertility Soil*, 35: 231-237.
- Azcon, R., J.M. Barea and D.S. Hayman, 1976. Utilization of rock phosphate in alkaline soils by plants inoculated with mycorrhizal fungi and phosphate solubilizing bacteria. *Soil Biol. Biochem.*, 8: 135-138.
- Bakker, P.A.H.M., J.G. Lamers, A.W. Bakker, J.D. Marugg, P.J. Weisbeek and B. Schippers, 1986. The role of siderophores in potato tuber yield increase by *Pseudomonas putida* in a short rotation of potato. *Eur. J. Plant Pathol.*, 92: 249-256.
- Bakker, P.A.H.M., A.W. Bakker, J.D. Marugg, P.J. Weisbeek and B. Schippers, 1987. Bioassay for studying the role of siderophores in potato growth stimulation by *Pseudomonas* spp. in short potato rotations. *Soil Biol. Biochem.*, 19: 443-449.
- Bar-Ness, E., Y. Chen, Y. Hader, H. Marschner and V. Romheld, 1991. Siderophores of *Pseudomonas putida* as an Iron Source for Dicot and Monocot Plants. In: *Iron Nutrition and Interaction in Plants*, Chen, Y. and Y. Hader (Eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands, pp: 271-281.
- Bar-Ness, E., Y. Hadar, Y. Chen, V. Romheld and H. Marschner, 1992. Short-term effects of rhizosphere microorganisms on Fe uptake from microbial siderophores by maize and oat. *Plant Physiol.*, 100: 451-456.
- Barbieri, P. and E. Galli, 1993. Effect on wheat root development of inoculation with an *Azospirillum brasilense* mutant with altered indole-3-acetic acid production. *Res. Microbiol.*, 144: 69-75.

- Beck, D.P., J. Wery, M.C. Saxena and A. Ayadi, 1991. Dinitrogen fixation and nitrogen balance in cool-season food legumes. *Agron. J.*, 83: 334-341.
- Belimov, A.A., A.P. Kojemiakor and C.V. Chuvarliyeva, 1995. Interactions between barley and mixed cultures of nitrogen fixing and phosphate solubilizing bacteria. *Plant Soil*, 173: 29-37.
- Ben Romdhane, S., M.E. Aouani, M. Trabelsi, P. de Lajudie and R. Mhamdi, 2008. Selection of high nitrogen-fixing rhizobia nodulating chickpea (*Cicer arietinum*) for semi-arid Tunisia. *J. Agron. Crop Sci.*, 194: 413-420.
- Benizri, E., E. Baudoin and A. Guckert, 2001. Root colonization by inoculated plant growth promoting rhizobacteria. *Biocontrol. Sci. Technol.*, 11: 557-574.
- Bent, E., S. Tuzun, C.P. Chanway and S. Enebak, 2001. Alterations in plant growth and in root hormone levels of lodgepole pines inoculated with rhizobacteria. *Can. J. Microbiol.*, 47: 793-800.
- Biswas, J.C., J.K. Ladha, F.B. Dazzo, Y.G. Yanni and B.G. Rolfe, 2000. Rhizobial inoculation influences seedling vigor and yield of rice. *Agron. J.*, 92: 880-886.
- Bohlool, B.B., J.K. Ladha, D.P. Garrity and T. George, 1992. Biological nitrogen fixation for sustainable agriculture. *Plant Soil*, 141: 1-11.
- Bolton, H.Jr., L.F. Elliott, R.F. Turco and A.C. Kennedy, 1990. Rhizoplane colonization of pea seedlings by *Rhizobium leguminosarum* and a deleterious root colonizing *Pseudomonas* sp. and effects on plant growth. *Plant Soil*, 123: 121-124.
- Bowen, G.D. and A.D. Rovira, 1999. The rhizosphere and its management to improve plant growth. *Adv. Agron.*, 66: 1-102.
- Bringhurst, R.M., Z.G. Cardon and D.J. Gage, 2001. Galactosides in the rhizosphere: Utilization by *Sinorhizobium meliloti* and development of a biosensor. *PNAS*, 98: 4540-4545.
- Brockwell, J. and P.J. Bottomley, 1995. Recent advances in inoculant technology and prospects for the future. *Soil Biol. Biochem.*, 27: 683-697.
- Burd, G.I., D.G. Dixon and B.R. Glick, 1998. A plant growth promoting bacterium that decreases nickel toxicity in plant seedlings. *Applied Environ. Microbiol.*, 64: 3663-3668.
- Burdman, S., H. Volpin, Y. Kapulnik and Y. Okon, 1996. Promotion of *nod* gene inducers and nodulation in common bean (*Phaseolus vulgaris*) root inoculated with *Azospirillum brasilense* Cd. *Applied Environ. Microbiol.*, 62: 3030-3033.
- Buysens, S., K. Heungens, J. Poppe and M. Höfte, 1996. Involvement of pyochelin and pyoverdine in suppression of *Pythium*-induced damping-off of tomato by *Pseudomonas aeruginosa* 7 NSK2. *Applied Environ. Microbiol.*, 62: 865-871.
- Bøckman, O.C., 1997. Fertilizers and biological nitrogen fixation as sources of plant nutrients: Perspectives for future agriculture. *Plant Soil*, 194: 11-14.
- Cakmakci, R., F. Kantar and O.F. Algur, 1999. Sugar beet and barley yields in relation to *Bacillus polymyxa* and *Bacillus megaterium* var. *Phosphaticum* inoculation. *J. Plant Nutr. Soil Sci.*, 162: 437-442.
- Cartieaux, F., M.C. Thibaud, L. Zimmerli, P. Lessard and C. Sarrobert *et al.*, 2003. Transcriptome analysis of *Arabidopsis* colonized by a plantgrowth promoting rhizobacterium reveals a general effect on disease resistance. *Plant J.*, 36: 177-188.
- Cattelan, A.J., P.G. Hartel and J.J. Fuhrmann, 1999. Screening of plant growth promoting rhizobacteria to promote early soybean growth. *J. Amer. Soil Scie. Soc.*, 63: 1670-1680.
- Chabot, R., C.J. Beauchamp, J.W. Kloepper and H. Antoun, 1998. Effect of phosphorus on root colonization and growth promotion of maize by bioluminescent mutants of phosphate-solubilizing *Rhizobium leguminosarum* biovar *Phaseoli*. *Soil Biol. Biochem.*, 30: 1615-1618.

- Chebotar, V.K., C.A. Jr. Asis and S. Akao, 2001. Production of plant growth-promoting substances and high colonization ability of rhizobacteria enhance the nitrogen fixation of soybean when coinoculation with *Bradyrhizobium japonicum*. Biol. Fertil. Soils, 34: 427-432.
- Chen, W.X., G.H. Yan and J.L. Li, 1988. Numerical taxonomic study of fast-growing soybean rhizobia and a proposal that *Rhizobium fredii* be assigned to *Sinorhizobium* gen. nov. Int. J. Syst. Bacteriol., 38: 392-397.
- Chen, T.W., S. Scherer and P. Boger, 1993. Nitrogen fixation of *Azorhizobium* in artificially induced root para-nodules in wheat. Curr. Plant Sci. Biotechnol. Agric., 14: 593-606.
- Christiansen-Weniger, C. and J. Vanderleyden, 1994. Ammonium excreting *Azospirillum* sp. become intracellularly established in maize (*Zea mays*) paranodules. Biol. Fertil. Soils, 17: 1-8.
- Christiansen-Weniger, C., 1996. Endophytic establishment of *Azorhizobium caulinodans* through auxin-induced root tumors of rice (*Oryza sativa* L.). Biol. Fertil. Soils, 21: 293-302.
- Cline, G.R., C.P.P. Reid and P.J. Szanislo, 1984. Effects of hydroxamate siderophore on iron absorption by sunflower and sorghum. Plant Physiol., 76: 36-39.
- Cook, D., D. Dreyer, D. Bonnet, M. Howell, E. Nony and K. van den Bosch, 1995. Transient induction of a peroxidase gene in *Medicago truncatula* precedes infection by *Rhizobium meliloti*. Plant Cell, 7: 43-55.
- Crowley, D.E., C.P.P. Reid and P.J. Szanislo, 1988. Utilization of microbial siderophores in iron acquisition by oat. Plant Physiol., 87: 680-685.
- Dashti, H., T.C. Mathew, M.M. Jadaun and E. Ashkanani, 1997. Zinc and liver cirrhosis: Biochemical and histopathologic assessment. Nutrition, 13: 206-218.
- Dashti, N., F. Zhang, R. Hynes and D.L. Smith, 1998. Plant growth promoting rhizobacteria accelerate nodulation and increase nitrogen fixation activity by field grown soybean (*Glycine max* (L.) Merr.) under short season conditions. Plant Soil, 200: 205-213.
- Davison, J., 1988. Plant beneficial bacteria. BioTechnology, 6: 282-286.
- Dazzo, F.B., Y.G. Yanni, R. Rizk, F.J. de Bruijn and J. Rademaker *et al.*, 2000. Progress in Multinational Collaborative Studies on the Beneficial Association Between *Rhizobium leguminosarum* bv. *Trifolii* and Rice. In: The Quest for Nitrogen Fixation in Rice, Ladha, J.K. and P.M. Reddy (Eds.). IRRI, Los Banos, Philippines.
- De Lajudie, P., A. Willems, G. Nick, F. Moreira and F. Molouba *et al.*, 1998a. Characterization of tropical tree rhizobia and description of *Mesorhizobium plurifarium* sp. Nov. Int. J. Syst. Bacteriol., 48: 369-382.
- De Lajudie, P., E. Laurent-Fulele, A. Willems, U. Torck and R. Coopman *et al.*, 1998b. *Allorhizobium undicola* gen. nov., sp. nov., nitrogen-fixing bacteria that efficiently nodulate *Neptunia natans* in Senegal. Int. J. Syst. Bacteriol., 48: 1277-1290.
- Derylo, M. and A. Skorupska, 1993. Enhancement of symbiotic nitrogen fixation by vitamin-secreting fluorescent *Pseudomonas*. Plant Soil, 154: 211-217.
- Dey, R., K.K. Pal, D.M. Bhatt and S.M. Chauhan, 2004. Growth promotion and yield enhancement of peanut (*Arachis hypogaea* L.) by application of plant growth-promoting rhizobacteria. Microbiol. Res., 159: 371-394.
- Dobbelaere, S., A. Croonenboghs, A. Thys, D. Ptacek and J. Vanderleyden *et al.*, 2001. Responses of agronomically important crops to inoculation with *Azospirillum*. Aust. J. Plant Physiol., 28: 871-879.
- Domey, S. and G. Lipmann, 1988. Stimulation of Plant Growth by Phosphate, Solubilizing Bacteria. In: Interrelationships between Microorganisms and Plants in Soil, Vancura, V. and F. Kunc (Eds.). Academic Press, San Diego, CA., San Diego, CA.

- Dorosinsky, L.M. and A.A. Kadyrov, 1975. Effect of inoculation on nitrogen fixation by chickpea, its crop and content of protein. *Mikrobiologiya*, 44: 1103-1106.
- Dreyfus, B., J.L. Garcia and M. Gillis, 1988. Characterization of *Azorhizobium caulinodans* gen. nov., sp. nov., a stem-nodulating nitrogen-fixing bacterium isolated from *Sesbania rostrata*. *Int. J. Syst. Bacteriol.*, 38: 89-98.
- Dubey, S.K., 1996. Combined effect of *Bradyrhizobium japonicum* and phosphate solubilizing *Pseudomonas striata* on nodulation, yield attributes and yield of rainfed soybean (*Glycine max*) under different sources of phosphorus in vertisols. *Ind. J. Agric. Sci.*, 66: 28-32.
- Duffy, B.K., A. Simon and D.M. Weller, 1996. Combination of *Trichoderma koningii* with fluorescent *Pseudomonads* for control of take-all on wheat. *Phytopathology*, 86: 188-194.
- Duffy, B.K. and G. Defago, 1999. Environmental factors modulating antibiotic and siderophore biosynthesis by *Pseudomonas fluorescens* biocontrol strains. *Applied Environ. Microbiol.*, 65: 2429-2438.
- Duijff, B.J., W.J. de Kogel, P.A.H.M. Bakker and B. Schippers, 1994. Influence of pseudobactin 358 on the iron nutrition of barley. *Soil Biol. Biochem.*, 26: 1681-1688.
- Dunne, C., J.J. Crowley, Y.M. Loccoz, D.N. Dowling, F. de Bruijn and F. O'Gara, 1993. Biological control of *Pythium ultimum* by *Stenotrophomonas maltophilia* W81 is mediated by an extracellular proteolytic activity. *Microbiology*, 143: 3921-3931.
- Elkoca, E., F. Kantar and F. Sahin, 2008. Influence of nitrogen fixing and phosphate solubilizing bacteria on nodulation, plant growth and yield of chickpea. *J. Plant Nutr.*, 33: 157-171.
- Elliott, L.F. and J.M. Lynch, 1984. Pseudomonads as a factor in the growth of winter wheat (*Triticum aestivum* L.). *Soil Biol. Biochem.*, 16: 69-71.
- Fallik, E., Y. Okon and M. Fischer, 1988. The effect of *Azospirillum brasilense* inoculation on metabolic enzyme activity in maize root seedlings. *Symbiosis*, 6: 17-28.
- Foster, R.C., A.D. Rovira and T.W. Cock, 1983. Ultrastructure of the Root-Soil Interface. The American Phytopathological Society, St Paul, MN., pp: 157.
- Frankenberger, W.T. Jr. and M. Arshad, 1995. *Phytohormones in Soil: Microbial Production and Function*. Marcel Dekker Inc., NY., USA., ISBN: 0824794427, pp: 503.
- Fred, E.B., I.L. Baldwin and E. McCoy, 1932. *Root Nodule Bacteria and Leguminous Plants*. University of Wisconsin, Madison, WI., USA., pp: 343.
- Freitas, J.R., M.R. Banerjee and J.J. Germida, 1997. Phosphate solubilizing *Rhizobacteria* enhance the growth and yield but not phosphorus uptake of canola (*Brassica napus* L.). *Biol. Fert. Soil*, 24: 358-364.
- Fridlender, M., J. Inbar and I. Chet, 1993. Biological control of soil borne plant pathogens by a b-1, 3-glucanase-producing *Pseudomonas cepacia*. *Soil Biol. Biochem.*, 25: 1211-1221.
- Fukudah, H., M. Takahashi, T. Fujiit and T. Ogawat, 1989. Ethylene production from L-methionine by *Cryptococcus albidus*. *J. Fermentation Bioeng.*, 67: 173-175.
- Gaind, S., R.S. Mathur and K.V.B.R. Tilak, 2000. Phosphate Solubilizing Microorganisms. In: *Recent Advances in Biofertilizer Technology*, Yadav, A.K., M.R. Motsara and S.R. Chaudhury (Eds.). Society for Promotion and Utilization Resources and Technology, New Delhi, pp: 190-206.
- Gantar, M. and J. Elhai, 1999. Colonization of wheat para-nodules by the N₂-fixing cyanobacterium *Nostoc* sp. strain 2S9B. *New Phytol.*, 141: 373-379.
- Garcia de Salamone, I.E., R.K. Hynes and L.N. Nelson, 2001. Cytokinin production by plant growth promoting rhizobacteria and selected mutants. *Can. J. Microbiol.*, 47: 103-113.

- Gaur, A.C., 1990. Physiological Functions of Phosphate Solubilizing Micro-Organisms. In: Phosphate Solubilizing Micro-Organisms as Biofertilizers, Gaur, A.C. (Eds.). Omega Scientific Publisher, New Delhi, pp: 16-72.
- Gaur, R., N. Shani, Kawaljeet, B.N. Johri, P. Rossi and M. Aragno, 2004. Diacetyl phloroglucinol-producing *Pseudomonas* do not influence AM fungi in wheat rhizosphere. *Curr. Sci.*, 86: 453-457.
- Glick, B.R., 1995. The enhancement of plant growth by free living bacteria. *Can. J. Microbiol.*, 41: 109-117.
- Glick, B.R., D.M. Penrose and J. Li, 1998. A model for the lowering of plant ethylene concentrations by plant growth promoting bacteria. *J. Theor. Biol.*, 190: 63-68.
- Glick, B.R., C.L. Patten, G. Holguin and D.M. Penrose, 1999. Biochemical and Genetic Mechanisms Used by Plant Growth Promoting Bacteria. Imperial College Press, London, UK.
- Goel, A.K., S.S. Sindhu and K.R. Dadarwal, 2000. Pigment diverse mutants of *Pseudomonas* sp.: Inhibition of fungal growth and stimulation of growth of *Cicer arietinum*. *Biol. Plant.*, 43: 563-569.
- Goel, A.K., S.S. Sindhu and K.R. Dadarwal, 2002. Stimulation of nodulation and plant growth of chickpea (*Cicer arietinum* L.) by *Pseudomonas* sp. antagonistic to fungal pathogens. *Biol. Fertil. Soils*, 36: 391-396.
- Gonzalez Lopez, J. and G.R. Vela, 1981. True morphology of the Azotobacteraceae- filterable bacteria. *Nature*, 289: 588-590.
- Gray, E.J. and D.L. Smith, 2005. Intracellular and extracellular PGPR: Commonalities and distinctions in the plant-bacterium signaling processes. *Soil Biol. Biochem.*, 37: 395-412.
- Greenough, D.R. and W.E. Batson, 1989. Potential for *Bacillus subtilis* as a bioprotectant of cotton seedlings in soil infested with *Rhizoctonia solani*. *Phytopathology*, 79: 373-373.
- Grichko, V.P. and B.R. Glick, 2001. Amelioration of flooding stress by ACC deaminase-containing plant growth-promoting bacteria. *Plant Physiol. Biochem.*, 39: 11-17.
- Grimes, H.D. and M.S. Mount, 1984. Influence of *Pseudomonas putida* on nodulation of *Phaseolus vulgaris*. *Soil Biol. Biochem.*, 16: 27-30.
- Guerinot, M.L. and B.K. Chelm, 1984. Isolation and expression of the Bradyrhizobium japonicum adenylate cyclase gene (*cya*) in *Escherichia coli*. *J. Bacteriol.*, 159: 1068-1071.
- Gull, M., F.Y. Hafeez, M. Saleem and K.A. Malik, 2004. Phosphorus uptake and growth promotion of chickpea by co-inoculation of mineral phosphate solubilizing bacteria and a mixed rhizobial culture. *Aust. J. Exp. Agric.*, 44: 623-628.
- Gupta, A., A.K. Sexena, M. Gopal and K.V.B.R. Tilak, 1998. Effect of plant growth promoting rhizobacteria on competitive ability of introduced *Bradyrhizobium* (*Vigna*) for *Nodulation* sp. *Microbial. Res.*, 153: 113-117.
- Gutierrez-Manero, F.J., B. Ramos-Solano, A. Probanza, J. Mehouchi, F.R. Tadeo and M. Talon, 2001. The plant-growth-promoting rhizobacteria *Bacillus pumilus* and *Bacillus licheniformis* produce high amounts of physiologically active gibberellins. *Physiol. Plant.*, 111: 206-211.
- Hamaoui, B., J.M. Abbadi, S. Burdman, A. Rashid. S. Sarig and Y. Okon, 2001. Effects of inoculation with *Azospirillum brasilense* on chickpeas (*Cicer arietinum*) and faba beans (*Vicia faba*) under different growth conditions. *Agronomie*, 21: 553-560.
- Handelsman, J. and E.V. Stabb, 1996. Biocontrol of soil borne plant pathogens. *Plant Cell*, 8: 1855-1869.
- Hartmann, A., M. Singh and W. Klingmuller, 1983. Isolation and characterization of *Azospirillum mutants* excreting high amounts of indolacetic acid. *Can. J. Microbiol.*, 29: 916-923.

- Herandez, L.G. and G.S. Hill, 1983. Effect of plant population and inoculation on yield and yield components of chickpea (*Cicer arietinum*). Proc. Agron. Soc. N.Z., 13: 75-79.
- Hiiifte, M., M.V. Woestyne and W. Verstraete, 1994. Role of Siderophores in Plant Growth Promotion and Plant Protection by Fluorescent Pseudomonads. In: Biochemistry of Metal Micronutrients in the Rhizosphere, Manthey, J.A., D.E. Crowley and D.G. Luster (Eds.). Lewis Publishers, Boca Raton, FL., USA., pp: 81-92.
- Holguin, G. and Y. Bashan, 1997. Nitrogen fixation by *Azospirillum brasilense* Cd is promoted when co-cultured with mangrove rhizobacteria *Staphylococcus* sp. Soil Biol. Biochem., 28: 1651-1660.
- Holguin, G., C.L. Patten and B.R. Glick, 1999. Genetics and molecular biology of *Azospirillum*. Biol. Fert. Soils, 29: 10-23.
- Holguin, G. and B.R. Glick, 2001. Expression of the ACC deaminase gene from *Enterobacter cloacae* UW4 in *Azospirillum brasilense*. Microbiol. Ecol., 41: 281-288.
- Howell, C.R. and R.D. Stipanovic, 1979. Control of rhizoctonia solani on cotton seedlings by *Pseudomonas fluorescens* and with an antibiotic producing bacteria. Phytopathology, 69: 480-482.
- Howell, C.R. and R.D. Stipanovic, 1980. Suppression of *Pythium ultimum*-induced damping-off of cotton seedling by *Pseudomonas fluorescens* and its antibiotic, Pyoluteorin. Phytopathology, 70: 712-715.
- Icgen, B., G.C. Gurakan and G. Ozcengiz, 2002. Characterization of *Salmonella enteritidis* isolates of chicken, egg and human origin from Turkey. Food Microbiol., 19: 375-382.
- Iruthayathas, E.E., S. Gunasekaran and K. Vlassak, 1983. Effect of combined inoculation of *Azospirillum* and *Rhizobium* on nodulation and N₂ fixation of winged bean and soybean. Scientia Hortic., 20: 231-240.
- Ismande, J., 1998. Iron, sulfur and chlorophyll deficiencies: A need for an integrative Physiol. Plant, 103: 139-144.
- Itzigsohn, R., S. Burdman, Y. Okon, E. Zaady, R. Yonatan and A. Perevolotsky, 2000. Plant-growth promotion in natural pastures by inoculation with *Azospirillum brasilense* under suboptimal growth conditions. Arid Soil Res. Manage., 14: 151-158.
- James, E.K, F.L. Olivares, J.I. Baldani and J. Dobereiner, 1997. *Herbaspirillum* an endophytic diazotroph colonizing vascular tissue in leaves of *Sorghum bicolor* L. Moench. J. Exp. Bot., 48: 785-797.
- Jarvis, B.D.W., P. van Berkum, W.X Chen, S.M. Nour, M.P. Fernandez, J.C Cleyet-Marel and M. Gillis, 1997. Transfer of *Rhizobium loti*, *Rhizobium huakuii*, *Rhizobium ciceri*, *Rhizobium mediterraneum* and *Rhizobium tianshanense* to *Mesorhizobium* gen. nov. Int. J. Syst. Bacteriol., 47: 895-898.
- Johri, B.N., C.V.S. Rao and R. Goel, 1997. Fluorescent Pseudomonads in Plant Disease Management. In: Biotechnological Approaches in Soil Microorganisms for Sustainable Crop Production, Dadarwal, K.R. (Eds.). Scientific Publishers, Jodhpur.
- Johri, B.N., A. Sharma and J.S. Viridi, 2003. Rhizobacterial diversity in India and its influence on soil and plant health. Adv. Biochem. Eng. Biotechnol., 84: 49-89.
- Joseph, B., R.R. Patra and R. Lawrence, 2007. Characterization of plant growth promoting rhizobacteria associated with chickpea (*Cicer arietinum* L.). Int. J. Plant Prod., 2: 141-152.
- Jurkevitch, E., Y. Hadar and Y. Chen, 1986. The remedy of lime-induced chlorosis in peanuts by *Pseudomonas* sp. siderophores. J. Plant Nutr., 9: 535-545.
- Kampert, M., E. Strzelczyk and A. Pokojska, 1975. Production of auxins by bacteria isolated from the roots of pine seedlings (*Pinus silvestris* L.) and from soil. Acta Microbiol. Polonica, 7: 135-143.

- Khalid, M., M. Arshad, A. Khalid and Z.A. Zahir, 2001. Biosynthesis of auxins by *Azotobacter*. Pak. J. Soil Sci., 20: 1-10.
- Khan, M., S. Aamil and A. Zaidi, 1997. Associative effect of *Bradyrhizobium* sp. (vigna) and M.d phosphate solubilizing bacteria on moongbean (*Vigna radiata* (L.) wilczek). Biojournal, 10: 101-106.
- Kim, J. and D.C. Rees, 1994. Nitrogenase and biological nitrogen fixation. Biochemistry, 33: 389-397.
- Kim, K.Y. and G.A. McDonald, 1998. Effect of phosphate solubilizing bacteria and vesicular-arbuscular mycorrhizae on tomato growth and soil microbial activity. Fert. Soils, 26: 79-87.
- Kittell, B.L., D.R. Helinski and G.S. Ditta, 1989. Aromatic aminotransferase activity and indoleacetic acid production in *Rhizobium meliloti*. J. Bacteriol., 171: 5458-5466.
- Kloepper, J.W. and M.N. Schroth, 1978. Plant growth-promoting rhizobacteria on radishes. 4th Int. Conf. Plant Pathogen Bacteria Angers France, 2: 879-882.
- Kloepper, J.W., J. Leong, M. Teintze and M.N. Schroth, 1980. Enhanced plant growth by siderophores produced by plant growth promoting rhizobacteria. Nature, 286: 885-886.
- Kloepper, J.W., R. Lifshitz and M.N. Scgroth, 1988. *Pseudomonas* Inoculants to Benefit Plant Production. ISI Atlas of Science, Animal and Plant Science Institute for Public Information, Philadelphia, USA., pp: 60-64.
- Kloepper, J.W., R. Lifshitz and R.M. Zablotowicz, 1989. Free-living bacteria inocula for enhancing crop productivity. Trends Biotechnol., 7: 39-44.
- Kloepper, J.W. and C.J. Beauchamp, 1992. A review of issues related to measuring of plant roots by bacteria. Can. J. Microbiol., 38: 1219-1232.
- Kloepper, J.W., S. Tuzun, L. Liu and G. Wei, 1993. Plant Growth-Promoting Rhizobacteria as Inducers of Systemic Disease Resistance. In: Pest Management: Biologically based Technologies, Lumsden, R.D. and J.L. Waughn (Eds.). American Chemical Society Publication, Washington, DC.
- Knight, T.J. and P.J. Langston-Unkefer, 1988. Enhancement of symbiotic dinitrogen fixation by a toxin-releasing plant pathogen. Science, 241: 951-954.
- Koval'skaya, N.Y., E.S. Lobakova and M.M. Umarov, 2001. The formation of artificial nitrogen-fixing symbioses with rape (*Brassica napus* var. *Napus*) plants in nonsterile soil. Microbiology, 70: 606-612.
- Kucey, R.M.N., H.H. Janzen and M.E. Legget, 1989. Microbial mediated increases in plant available phosphorus. Adv. Agron., 42: 199-228.
- Kumar, B.S.D. and B. Bezbaruah, 1997. Plant growth promotion and fungal pest control through an antibiotic and siderophore producing fuorescent *Pseudomonas* strain from tea (*Camellia sinensis* L.) O Kuntze plantations. Ind. J. Exp. Biol., 35: 289-292.
- Kumar, V. and N. Narula, 1999. Solubilization of inorganic phosphates and growth emergence of wheat as affected by *Azotobocter chroococcum* mutants. Biol. Fert. Soils, 28: 301-305.
- Kumar-Dileep, B.S. and B. Bezbaruah, 1996. Antibiosis and plant growth promotion by a *Pseudomonas strain* isolated from soil under tea cultivation. Indian J. Microbiol., 36: 45-48.
- Kyei-Boahen, S., A.E. Slinkard and F.L. Walley, 2002. Evaluation of rhizobial inoculation method for chickpea. Agron. J., 94: 851-859.
- Ladha, J.K. and D.K. Kundu, 1997. Legumes for Sustaining Soil Fertility in Lowland Rice. In: Extending Nitrogen Fixation Research, Rupela, O.P., C. Johansen and D.F. Herridge (Eds.). International Crops Research Institute for the Semi-Arid Tropics, Patancheru Andra Pradesh, India, pp: 76-102.

- Lambrecht, M., Y. Okon, A.V. Broek and J. Vanderleyden, 2000. Indole-3-acetic acid: A reciprocal signaling molecule in bacteria-plant interaction. Trends Microbiol., 8: 298-300.
- Laranjo, M., R. Rodrigues L. Alho and S. Oliveira, 2001. Rhizobia of chickpea from southern Portugal: Symbiotic efficiency and genetic diversity. J. Applied Microbiol., 90: 662-667.
- Laranjo, M., C. Branco, R. Soares, L. Alho, M.D. Carvalho and S. Oliveira, 2002. Comparison of *Chickpea rhizobia* isolates from diverse portuguese natural populations based on symbiotic effectiveness and DNA fingerprint. J. Applied Microbiol., 92: 1043-1050.
- Lee, J.Y. and S.H. Song, 2007. Evaluation of groundwater quality in coastal areas: Implications for sustainable agriculture. Environ. Geol., 52: 1231-1242.
- Li, J., D.H. Ovakim, T.C. Charles and B.R. Glick, 2000. An ACC deaminase minus mutant of *Enterobacter cloacae* UW4 no longer promotes root elongation. Curr. Microbiol., 41: 101-105.
- Lindstrom K., 1989. *Rhizobium galegae* a new species of legume root nodule bacteria. Int. J. Syst. Bacteriol., 39: 365-367.
- Liu, S.T., K.L. Perry, C.L. Schardl and C.I. Kado, 1982. *Agrobacterium* Ti plasmid indoleacetic acid gene is required for crown gall oncogenesis. Proc. Nat. Acad. Sci. USA., 79: 2812-2816.
- Liu, L., J.W. Kloepper and S. Tuzun, 1995. Induction of systemic resistance in cucumber against fusarium wilt by plant growth-promoting rhizobacteria. Phytopathology, 85: 695-698.
- Loper, J. E. and M.N. Scroth, 1986. Influence of bacterial sources of indole-3-acetic acid on root elongation of sugar beet. Phytopathol., 76: 386-389.
- Loper, J.E., 1988. Role of fluorescent siderophore production in biological control of *Pythium ultimum* by a *Pseudomonas fluorescens* strain. Phytopathology, 78: 166-172.
- Lucangeli, C. and R. Bottini, 1997. Effect of *Azospirillum* sp. On endogenous gibberellin content and growth and growth of maize (*Zea mays* L.) treated with uniconazole. Symbiosis, 23: 63-71.
- Lugtenberg, B.J.J., L. Dekkers and G.V. Bloemberg, 2001. Molecular determinants of rhizosphere colonization by *Pseudomonas*. Ann. Rev. Phytopathol., 38: 461-490.
- Maatallah, J., E.B. Berraho, S. Munoz, J. Sanjuan and C. Lluch, 2002. Phenotypic and molecular characterization of chickpea rhizobia isolated from different areas of Morocco. J. Applied Microbiol., 93: 531-540.
- Mahafee, W.F. and J.W. Kloepper, 1997. Temporal changes in the bacterial communities of soil, rhizosphere and endorhiza associated with field-grown cucumber (*Cucumis sativus* L.). Microbial. Ecol., 34: 210-223.
- Manjunatha, H. and L.S. Devi, 1990. Effect of phosphate solubilizing bacteria on P availability to groundnut from rock phosphate. Curr. Res., 19: 56-57.
- Mansouri, S. and A.W. Bunch, 1989. Bacterial ethylene synthesis from 2-oxo-4-thyobutyric acid and from methionine. J. Gen. Microbiol., 135: 2819-2827.
- Marschner, H. and V. Rohmheld, 1994. Strategies of plants for acquisition of iron. Plant Soil, 165: 261-274.
- Martinez-Romero, E. and E.T. Wang, 2000. *Sesbania herbacea*-*Rhizobium huautlense* nodulation in flooded soils and comparative characterization of *S. herbacea* nodulating rhizobia in different environments. Microbe Ecol., 41: 25-32.
- Masalha, J., H. Kosegarten, O. Elmaci and K. Mengel, 2000. The central role of microbial activity for iron acquisition in maize and sunflower. Biol. Fert. Soils, 30: 433-439.
- Mayak, S., T. Tivosh and B.R. Glick, 1999. Effect of wild type and mutant plant growth-promoting rhizobacteria on the rooting of mung bean cuttings. J. Plant Growth Regul., 18: 49-53.

- Mirza, M.S., W. Ahmad, F. Latif, J. Haurat, R. Bally, P. Normand and K.A. Malik, 2001. Isolation, partial characterization and the effect of Plant Growth-Promoting Bacteria (PGPB) on micro-propagated sugarcane *in vitro*. *Plant Soil*, 237: 47-54.
- Mittal, V., O. Singh, H. Nayyar, J. Kaur and R. Tewari, 2008. Stimulatory effect of phosphate-solubilizing fungal strains (*Aspergillus awamori* and *Penicillium citrinum*) on the yield of chickpea (*Cicer arietinum* L. cv. GPF2). *Soil Biol. Biochem.*, 40: 718-727.
- Molla, A.H., Z.H. Shamsuddin, M.S. Halimi, M. Marziah and A.B. Puteh, 2001. Potential for enhancement of root growth and nodulation of soybean co-inoculated with *Azospirillum* and *Bradyrhizobium* in laboratory systems. *Soil Biol. Biochem.*, 33: 457-463.
- Murphy, J.F., G.W. Zehnder, D.J. Schuster, E.J. Sikora, J.E. Polston and J.W. Kloepper, 2000. Plant growth-promoting rhizobacterial mediated protection in tomato against Tomato mottle virus. *Plant Dis.*, 84: 779-784.
- Nandakumar, R., 1998. Induction of systemic resistance in rice with fluorescent *Pseudomonas* for the management of sheat blight disease. M.Sc. Thesis, Tnua, Coimbatore, India
- Nautiyal, C.S., S. Bhadauria, P. Kumar, H. Lal, R. Mondal and D. Verma, 2000. Stress induced phosphate solubilization in bacteria isolated from alkaline soils. *FEMS Microbiol. Lett.*, 182: 291-296.
- Nehl, D.B., S.J. Allen and J.F. Brown, 1996. Deleterious rhizosphere bacteria: An integrating perspective. *Applied Soil Ecol.*, 5: 1-20.
- Neilands, J.B., 1981. Iron absorption and transport in microorganisms. *Annual. Rev. Nutr.*, 1: 27-46.
- Nick, G., P. de Lajudie, B.D. Eardly, S. Soumalainen and L. Paulin *et al.*, 1999. *Sinorhizobium arboris* sp. nov. and *Sinorhizobium kostiense* sp. nov. isolated from leguminous trees in Sudan and Kenya. *Int. J. Syst. Bacteriol.*, 49: 1359-1368.
- Nieto, K.F. and W.T.Jr. Frankenberger, 1989. Biosynthesis of cytokinins produced by *Azotobacter chroococcum*. *Soil Biol. Biochem.*, 21: 967-972.
- Nour, S.M., M.P. Fernandez, P. Normand and J.C. Cleyet-Marel, 1994. *Rhizobium ciceri* sp. nov. consisting of strains that nodulate chickpeas (*Cicer arietinum* L.). *Int. J. Syst. Bacteriol.*, 44: 511-522.
- Nour, S.M., J.C. Cleyet-Marel, P. Normand and M.P. Fernandez, 1995. Genomic heterogeneity of strains nodulating chickpeas (*Cicer arietinum* L.) and description of *Rhizobium mediterraneum* sp. nov. *Int. J. Syst. Bacteriol.*, 45: 640-648.
- Offringa, I.A., L.S. Melchers, A.J.G. Regenburt-Tuink, P. Costantino, R.A. Schilperoord and P.J.J. Haoykaas, 1986. Complementation of *Agrobacterium tumefaciens* tumor-inducing aux mutant by genes from the TR-region of the Ri-plasmid of *Agrobacterium rhizogenes*. *Proc. Nat. Acad. Sci. USA.*, 83: 6935-6939.
- Oliveira, F.L., E.M. Ferreira and M.E. Pampulha, 1997. Nitrogen fixation, nodulation and yield of clover plants co-inoculated with root-colonizing bacteria. *Symbiosis*, 23: 35-42.
- Oremus, P.A.I., 1980. Occurrence and infective potential of the endophyte *Frankia* of *Hippophae rhamnoides* L. sp. *Rhamnoides* in coastal sand-dune areas. *Plant Soil*, 56: 123-139.
- Ozturk, A., O. Caglar and F. Sahin, 2003. Yield response of wheat and barley to inoculation of Plant growth promoting rhizobacteria at various levels of nitrogen fertilizers. *J. Plant Nutr. Soil Sci.*, 166: 262-266.
- Pal, S.S., 1998. Interactions of an acid tolerant strain of phosphate solubilizing bacteria with a few acid tolerant crops. *Plant Soil*, 198: 169-177.
- Pal, K.K., K.V.B.R. Tilak, A.K. Saxena, R. Dey and C.S. Singh, 2001. Suppression of maize root diseases caused by *Macrophomina phaseolina*, *Fusarium moniliforme* and *Fusarium graminearum* by plant growth promoting rhizobacteria. *Microbiol. Res.*, 156: 209-223.

- Pargiri, I. and B. Bezbaruah, 1990. Strains contributing to phosphorus mobilization in acid soils. *Ind. J. Agric. Sci.*, 60: 197-200.
- Parmar, N. and K.R. Dadarwal, 1999. Stimulation of nitrogen fixation and induction of flavonoid like compounds by rhizobacteria. *J. Applied Microbiol.*, 86: 36-44.
- Patten, C.L. and B.R. Glick, 1996. Bacterial biosynthesis of indole-3-acetic acid. *Can. J. Microbiol.*, 42: 207-220.
- Patten, C.L. and B.R. Glick, 2002. The role of bacterial indoleacetic acid in the development of the host plant root system. *Applied Environ. Microbiol.*, 68: 3795-3801.
- Pazout, J., M. Wurst and V. Vancura, 1981. Effect of aeration on ethylene production by soil bacteria samples cultivated in a closed system. *Plant Soil*, 62: 431-437.
- Petersen, D.J., M. Srinivasan and C.P. Chanway, 1996. *Bacillus polymyxa* stimulates increased *Rhizobium etli* populations and nodulation when co-resident in the rhizosphere of *Phaseolus vulgaris*. *FEMS Microbiol. Lett.*, 142: 271-276.
- Pierson, E.A. and D.M. Weller, 1994. Use of mixtures of fluorescent pseudomonada to suppress take-all and improve the growth of wheat. *Phytopathology*, 84: 940-947.
- Pilet, P.E., M.C. Elliott and M.M. Moloney, 1979. Endogenous and exogenous auxin in the control of root growth. *Planta*, 146: 405-408.
- Pinton, R., Z. Varanini and P. Nannipieri, 2001. The Rhizosphere as a Site of Biochemical Interactions Among Soil Components, Plants and Microorganisms. In: *The Rhizosphere*, Pinton, R., Z. Varanini and P. Nannipieri (Eds.). Marcel Dekker, Inc., New York, USA., pp: 1-18.
- Pishchik, V.N., N.I. Vorobyev, I.I. Chernyaeva, S.V. Timofeeva, A.P. Kozhemyakov, Y.V. Alexeev and S.M. Lukin, 2002. Experimental and mathematical simulation of plant growth promoting rhizobacteria and plant interaction under cadmium stress. *Plant Soil*, 243: 173-186.
- Polonenko, D.R., F.M. Scher, J.W. Kloepper, C.A. Singleton, M. Laliberte and I. Zalerka, 1987. Effect of root colonizing bacteria on nodulation of soybean roots by *Bradyrhizobium japonicum*. *Can. J. Microbiol.*, 33: 498-503.
- Press, C.M., J.E. Loper and J.W. Kloepper, 2001. Role of iron in rhizobacteria mediated induced systemic resistance of cucumber. *Phytopathology*, 91: 593-598.
- Prithiviraj, B., X. Zhou, A. Souleimanov, W.M. Kahn and D.L. Smith, 2003. A host-specific bacteria-to-plant signal molecule (Nod factor) enhances germination and early growth of diverse crop plants. *Planta*, 21: 437-445.
- Quingwen, Z., L. Ping, W. Gang and C. Qingnian, 1998. On the biochemical mechanism of induced resistance of cotton to cotton bollworm by cutting off young seedling at plumular axis. *Acta Phytophylocica Sinica*, 25: 209-212.
- Rao, N.S., 1992. *Biofertilizers in Agriculture*. AA Balkema, Rotterdam, pp: 188.
- Raupach, G.S., L. Liu, J.F. Murphy, S. Tuzun and J.W. Kloepper, 1996. Induced systemic resistance in cucumber and tomato against *Cucumber mosaic virus* using plant growth-promoting rhizobacteria (PGPR). *Plant Dis.*, 80: 891-894.
- Renwick, A., R. Campbell and S. Coe, 1991. Assessment of in vivo screening systems for potential biocontrol agents of *Gaeumannomyces graminis*. *Plant Pathol.*, 40: 524-532.
- Richardson, A.E., 2001. Prospects for using soil microorganisms to improve the acquisition of phosphorus by plants. *Aust. J. Plant Physiol.*, 28: 897-906.
- Rigby, D. and D. Caceres, 2001. Organic farming and the sustainability of agricultural systems. *Agric. Syst.*, 68: 21-40.
- Rokade, S.M. and P. Patil, 1993. Phosphate solubilizing microorganisms: A review. *J. Maharashtra Agric. Univ.*, 18: 93-101.
- Rome, S., M.P. Fernandez, B. Brunel, P. Normand and J.C. Cleyet-Marel, 1996. *Sinorhizobium medicae* sp. nov. isolated from annual *Medicago* spp. *Int. J. Syst. Bacteriol.*, 46: 972-980.

- Rupela, O.P. and M.C. Saxena, 1987. Nodulation and Nitrogen Fixation in Chickpea. In: The Chickpea, Saxena, M.C. and K.B. Singh (Eds.). CAB International, Wallingford, Oxon, pp: 191-206.
- Sahin, F., R. Cakmakci and F. Kantar, 2004. Sugar beet and barley yields in relation to inoculation with N₂-fixing and phosphate solubilizing bacteria. *Plant Soil*, 265: 123-129.
- Saikia, R., T. Singh, R. Kumar, J. Srivastava, A.K. Srivastava, K. Singh and D.K. Arora, 2003. Role of salicylic acid in systemic resistance induced by *Pseudomonas fluorescens* against *Fusarium oxysporum* f. sp. *ciceri* in chickpea. *Microbiol. Res.*, 158: 203-213.
- Saikia, R., R. Kumar, T. Singh, A.K. Srivastava, D.K. Arora and M.W. Lee, 2004. Induction of defense related enzymes and pathogenesis related proteins in *Pseudomonas fluorescens*-treated chickpea in response to infection by *Fusarium oxysporum* f.sp. *Ciceri*. *Mycobiology*, 32: 47-52.
- Saikia, R., B.P. Singh, R. Kumar and D.K. Arora, 2005. Detection of pathogenesis related proteins chitinase and 1, 3-glucanase in induced chickpea. *Curr. Sci.*, 89: 659-663.
- Saikia, R., S. Varghese, B.P. Singh and D.K. Arora, 2009. Influence of mineral amendment on disease suppressive activity of *Pseudomonas fluorescens* to *Fusarium* wilt of chickpea. *Microbiol. Res.*, 164: 365-373.
- Saleh, S.S. and B.R. Glick, 2001. Involvement of *gacS* and *rpoS* in enhancement of the plant growth-promoting capabilities of *Enterobacter cloacae* CAL2 and UW4. *Can. J. Microbiol.*, 47: 698-705.
- Salisbury, F.B. and C.W. Ross, 1992. *Plant Physiology*. Wadsworth, California, pp: 451-497.
- Salisbury, F.B., 1994. The Role of Plant Hormones. In: *Plant-Environment Interactions*, Wilkinson, R.E. (Ed.). Marcel Dekker, New York, USA., pp: 39-81.
- Sarig, S., Y. Kapulnik and Y. Okon, 1986. Effect of *Azospirillum* inoculation on nitrogen fixation and growth of several winter legumes. *Plant Soil*, 90: 335-342.
- Sarioglu, G., S. Ozcelik and S. Kaymaz, 1993. Selection of effective nodosity bacteria (*Rhizobium leguminosarum* biovar. *viceae*) from lentil grown in Elazığ. *Turk. J. Agric. For.*, 17: 569-573.
- Sato, M., K. Watanabe, M. Yazawa, Y. Takikawa and K. Nishiyama, 1997. Detection of new ethylene production bacteria *Pseudomonas syringae* pvs. *Cannabina* and *Sesami* by PCR amplification of genes for the ethylene forming enzyme. *Phytopathology*, 87: 1192-1196.
- Sawada, H., L.D. Kuykendall and J.M. Young, 2003. Changing concepts in the systematics of bacterial nitrogen fixing legume symbionts. *J. Gen. Applied Microbiol.*, 49: 155-179.
- Scher, F.M. and R. Baker, 1982. Effect of *Pseudomonas putida* and a synthetic iron chelator on induction of soil suppressiveness to *Fusarium* wilt pathogens. *Phytopathology*, 72: 1567-1573.
- Schobeck, F., H.W. Dehne and W. Beicht, 1980. Untersuchungen zur Aktivierung unspezifischer Resistenzmechanismen in Pflanzen. *Z. Pflk. Pflschutz*, 87: 654-666.
- Scholla, M.H. and G.H. Elkan, 1984. *Rhizobium fredii* sp. nov., a fast-growing species that effectively nodulates soybeans. *Int. J. Syst. Bacteriol.*, 34: 484-486.
- Segovia, L., J.P.W. Young and E. Martinez-Romero, 1993. Reclassification of American *Rhizobium leguminosarum* biovar *phaseoli* type I strain as *Rhizobium etli* sp. nov. *Int. J. Syst. Bacteriol.*, 43: 374-377.
- Sessitsch, A., J.G. Howieson, X. Perret, H. Antoun and E. Martinez-Romero, 2002. Advances in *Rhizobium* research. *Crit. Rev. Plant Sci.*, 21: 323-378.
- Shah, S., J. Li, B.A. Moffat and B.R. Glick, 1998. Isolation and Characterization of ACC deaminase genes from two different plant growth promoting rhizobacteria. *Can. J. Microbiol.*, 44: 833-843.

- Siddiqui, I.A. and S. Ehteshamul-Haque, 2001a. Suppression of the root rot-root knot disease complex by *Pseudomonas aeruginosa* in tomato: The influence of inoculum density, nematode populations, moisture and other plant associated bacteria. *Plant Soil*, 237: 81-89.
- Siddiqui, Z.A. and I. Mahmood, 2001b. Effects of rhizobacteria and root symbionts on the reproduction of *Meloidogyne javanica* and growth of chickpea. *Bioresour. Technol.*, 79: 41-45.
- Sikoraa, E.J., R.T. Gudauskas, J.F. Murphy, D.W. Porch and M. Andianifahanana *et al.*, 1998. A multivirus epidemic of tomatoes in Alabama. *Plant Dis.*, 82: 117-120.
- Sindhu, S.S., S.K. Gupta and K.R. Dadarwal, 1999. Antagonistic effect of *Pseudomonas* sp. on pathogenic fungi and enhancement of growth of green gram (*Vigna radiata*). *Biol. Fertil. Soils*, 29: 62-68.
- Sindhu, S.S., S. K. Gupta, S. Suneja and K.R. Dadarwal, 2002a. Enhancement of green gram nodulation and growth by *Bacillus* species. *Biologia Plantarum*, 45: 117-120.
- Sindhu, S.S., S. Suneja, A.K. Goel, N. Paramar and K.R. Dadarwal, 2002b. Plant growth promoting effects of *Pseudomonas* sp on coinoculation with *Mesorhizobium* sp. cicer strain under sterile and wilt sick soil conditions. *Applied Soil Ecol.*, 19: 57-64.
- Singh, S. and K.K. Kapoor, 1999. Inoculation with phosphate solubilizing microorganisms and a vesicular arbuscular mycorrhizal fungus improves dry matter yield and nutrient uptake by wheat grown in a sandy soil. *Biol. Fert. Soils*, 28: 139-144.
- Sivaramaiah, N, D.K. Malik and S.S. Sindhu, 2007. Improvement in symbiotic efficiency of chickpea (*Cicer arietinum*) by coinoculation of *Bacillus* strains with *Mesorhizobium* sp *Cicer*. *Indian J. Microbiol.*, 47: 51-56.
- Srinivasan, M., D.J. Petersen and F.B. Holl, 1996. Influence of indoleacetic-acid-producing *Bacillus* isolates on the nodulation of *Phaseolus vulgaris* by *Rhizobium etli* under gnotobiotic conditions. *Can. J. Microbiol.*, 42: 1006-1014.
- Srivastava, A.K., T. Singh, T.K. Jana and D.K. Arora, 2001. Induced resistance and control of charcoal rot in *Cicer arietinum* (chickpea) by *Pseudomonas fluorescens*. *Can. J. Bot.*, 79: 787-795.
- Stevenson, F.J. and M.A. Cole, 1999. *Cycles of Soil: Carbon, Nitrogen, Phosphorous, Sulfur and Micronutrients*. 2nd Edn., Wiley and Sons Inc., Tronto, Canada, ISBN-10: 0471320714.
- Sturtevant, D. and B.J. Taller, 1989. Cytokinin production by *Parasponia nodule bacterium*. Proceedings of the 89th Annual Meeting of American Society of Microbiology, (AAMASM'89), American Society of Microbiology, Washington, DC., pp: 300-300.
- Sung, K.C. and Y.R. Chung, 1997. Enhanced Suppression of Rice Sheath Blight Using Combination of Bacteria Which Produced Chitinases or Antibiotics. In: *Plant Growth Promoting Rhizobacteria Present Status and Future Prospects*, Ogoshi, A., K. Kobayashim, Y. Homma, F. Kodama, N. Kondo and S. Akino (Eds.). Nakanishi Printing, Sapporo, Japan.
- Swanson, B.T., H.F. Wilkins and B.W. Kennedy, 1979. Factor affecting ethylene production by some plant pathogenic bacteria. *Plant Soil*, 51: 19-26.
- Taller, B.J. and D.B. Sturtevant, 1991. Cytokinin Production by Rhizobia: Current Plant Science and Biotechnology in Agriculture, In: *Advances in Molecular Genetics of Plant Microbes Interaction*, Hennecke, H. and D.P.S. Verma (Eds.). Kluwer Academic Publication, Dordrecht, Netherlands, pp: 215-221.
- Tchan, Y.T., A.M.M. Zeman and I.R. Kennedy, 1991. Nitrogen fixation in para-nodules of wheat roots by introduced free-living diazotrophs. *Plant Soil*, 137: 43-47.

- Tchebotar, V.K., U.G. Kang, C.A. Jr. Asis and S. Akao, 1998. The use of GUS-reporter gene to study the effect of *Azospirillum-rhizobium* coinoculation on nodulation of white clover. *Biol. Fertil. Soils*, 27: 349-352.
- Thimann, K.V., 1935. On the plant growth hormone produced by *Rhizopus suinus*. *J. Biol. Chem.*, 109: 279-291.
- Tilak, K.V.B.R., N. Ranganayaki and C. Manoharachari, 2006. Synergistic effect of plant growth promoting rhizobacteria and rhizobium on nodulation and nitrogen fixation by pigeonpea (*Cajanus cajan*). *Eur. J. Soil Sci.*, 57: 67-71.
- Valverde, A., A. Burgos, T. Fiscella, R. Rivas and E.V. Quez *et al.*, 2006. Differential effects of coinoculations with *Pseudomonas jessenii* PS06 (a phosphate-solubilizing bacterium) and *Mesorhizobium ciceri* C-2/2 strains on the growth and seed yield of chickpea under greenhouse and field conditions. *Plant Soil*, 287: 43-50.
- Van Peer, R., G.J. Niemann and B. Schippers, 1991. Induced resistance and phytoalexin accumulation in biological control of fusarium wilt of carnation by *Pseudomonas* sp. strain WCS 417r. *Phytopathology*, 81: 728-734.
- Vazquez, P., G. Holguin, M.E. Puente, A. Lopez-Cortes and Y. Bashan, 2000. Phosphate solubilizing microorganisms associated with the rhizosphere of mangroves in a semiarid coastal lagoon. *Biol. Fert. Soils*, 30: 460-468.
- Velazquez, E., J.M. Igual, A. Willems, M.P. Fernandez and E. Munoz *et al.*, 2001. *Mesorhizobium chacoense* sp. nov., a novel species that nodulates *Prosopis alba* in the Chaco Arido region (Argentina). *Int. J. Syst. Evol. Microbiol.*, 51: 1011-1021.
- Verma, J.P., J. Yadav and K.N. Tiwari, 2009. Effect of *Mesorhizobium* and plant growth promoting rhizobacteria on nodulation and yields of chickpea. *Biol. Forum-An Int. J.*, 1: 11-14.
- Verma, J.P., J. Yadav and K.N. Tiwari, 2010. Application of *Rhizobium* sp. BHURC01 and plant growth promoting rhizobacteria on nodulation, plant biomass and yields of Chickpea (*Cicer arietinum* L.). *Int. J. Agric. Res.*, 5: 148-156.
- Vessey, J.K. and T.J. Buss, 2002. *Bacillus cereus* UW85 inoculation effects on growth, nodulation, and N accumulation in grain legumes. Controlled-environment studies. *Can. J. Plant Sci.*, 82: 282-290.
- Vessey, J.K., 2003. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil*, 255: 571-586.
- Vidhayasekaran, P. and M. Muthamilan, 1999. Evaluation of powder formulation of *Pseudomonas fluorescens* Pfl for control of rice sheath blight. *Biocontrol Sci. Technol.*, 9: 67-74.
- Voisard, C., C. Keel, D. Haas and G. Defago, 1989. Cyanide production by *Pseudomonas fluorescens* helps suppress black root rot of tobacco under gnotobiotic conditions. *EMBO J.*, 8: 351-358.
- Wang, C., E. Knill, B.R. Glick and G. Defago, 2000. Effect of transferring 1-aminocyclopropane-1-carboxylic acid (ACC) deaminase genes into *Pseudomonas fluorescens* strain CHAO and its *gacA* derivative CHA96 on their growth-promoting and disease-suppressive capacities. *Can. J. Microbiol.*, 46: 898-907.
- Wang, Y., H.N. Brown, D.E. Crowley and P.J. Szanislo, 1993. Evidence for direct utilization of a siderophore, ferrioxamine B in axenically grown cucumber. *Plant Cell Environ.*, 16: 579-585.
- Wei, G., J.W. Koepper and S. Tuzun, 1991. Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology*, 81: 1508-1512.

- Wei, G., J.W. Kloepper and S. Tuzun, 1996. Induction of systemic resistance to cucumber disease and increase plant growth by plant growth-promoting rhizobacteria under field conditions. *Phytopathology*, 86: 221-224.
- Weller, D.M. and R. Cook, 1986. Increased growth of wheat by seed treatment with fluorescent pseudomonads and implications of Pythium control. *Can. J. Plant Pathol.*, 8: 328-344.
- Whitelaw, M.A., 2000. Growth promotion of plants inoculated with phosphate-solubilizing fungi. *Adv. Agron.*, 69: 99-152.
- Whitelaw, M.A., T.A. Hardenand and G.L. Bender, 1997. Plant growth promotion of wheat inoculated with *Penicillium radicum* sp. nov. *Aust. J. Soil Res.*, 35: 291-300.
- Xie, H., J.J. Pasternak, B.R. Glick, 1996. Isolation and characterization of mutants of the plant growth-promoting rhizobacterium *Pseudomonas putida* GR-122 that overproduce indoleacetic acid. *Curr. Microbiol.*, 32: 67-71.
- Yadav, J. and J.P. Verma, 2009. Response of wheat to PGPR and organic manures in cereal and legume based cropping sequences under nascent stage of organic farming. *Ann. Plant Soil Res.*, 11: 122-125.
- Yadav, J., J.P. Verma, S.K. Yadav and K.N. Tiwari, 2010. Effect of salt concentration and pH on soil inhabiting fungus *Penicillium citrinum* Thom for solubilization of *Tricalcium phosphate*. *Microbiol. J.*, 1: 1-7.
- Yami, Y.G. and F.K.A. El-Fattah, 1999. Towards integrated biofertilization management with free living and associative dinitrogen fixers for enhancing rice performance in the Nile delta. *Symbiosis*, 27: 319-331.
- Yami, Y.G., 1992. The effect of Cyanobacteria and Azolla on the performance of rice under different levels of fertilizer nitrogen. *Biotechnology*, 8: 132-136.
- Youssef, H, M. Monib, M. Fayez and N.A. Hegazi, 1998. Induction of paranodulation by 2,4-D, IAA and tryptophan in wheat inoculated with various diazotrophs. *Dev. Plant Soil Sci.*, 79: 43-150.
- Zahir, Z.A., M. Akram, M. Arshad and A. Khalid, 1998a. Improving Maize yield by inoculation with plant growth promoting rhizobacteria. *Pak. J. Soil Sci.*, 15: 7-11.
- Zahir, Z.A., M. Arshad and A. Khalid, 1998b. Promoting crop yield by *Azotobacter* inoculation. *J. Soil Sci.*, 15: 7-11.
- Zahir, Z.A., AS. Abbas. M. Khalid and M. Arshad, 2000. Substrate dependent microbially derived plant hormones for improving growth of maize seedling yield by inoculation with plant growth promoting rhizobacteria. *Pak. J. Biol. Sci.*, 3: 289-291.
- Zaidi, A., M.S. Khan and M.D. Amil, 2003. Interactive effect of rhizotrophic microorganisms on yield and nutrient uptake of chickpea (*Cicer arietinum* L.). *Eur. J. Agron.*, 19: 15-21.
- Zaidi, A. and M.S. Khan, 2007. Stimulatory effects of dual inoculation with phosphate solubilizing microorganisms and arbuscular mycorrhizal fungus on chickpea. *Aust. J. Exp. Agric.*, 47: 1016-1022.
- Zehnder, G., J.W. Kloepper, C. Yao, G. Wei, O. Chambliss and R. Shelby, 1997. Induction of systemic resistance in cucumber against cucumber beetles (Coleoptera: Chrysomelidae) by plant resistance. *Entomol. Exp. Appl.*, 83: 81-85.
- Zhang, S., A.L. Moyne, M.S. Reddy and J.W. Kloepper, 2002. The role of salicylic acid in induced systemic resistance elicited by plant growth-promoting rhizobacteria against blue mold of tobacco. *Biological Control*, 25: 288-296.