



# International Journal of Botany

ISSN: 1811-9700

**science**  
alert

**ANSI***net*  
an open access publisher  
<http://ansinet.com>

## ***Ipomoea aquatica*, An Underutilized Green Leafy Vegetable: A Review**

<sup>1</sup>K. Nagendra Prasad, <sup>1</sup>G.R. Shivamurthy and <sup>2</sup>S.M. Aradhya

<sup>1</sup>Department of Studies in Botany, University of Mysore,  
Manasagangotri, Mysore, 570 006, India

<sup>2</sup>Department of Fruit and Vegetable Technology,  
Central Food Technological Research Institute, Mysore, 570 020, India

**Abstract:** Fruits and vegetables are emerging fast as most economical and nutritious global foods. They were declared as healthy foods of the millennium and also described as nutraceutical foods of the century, owing to their health benefits. *Ipomoea aquatica* Forsk., a commonly grown green leafy vegetable is a rich source of vitamins, minerals, proteins, fibers, carotenes and flavonoids with many health benefits. It grows as a common weed in lakes and fresh water ponds but it is usually neglected because of lack of complete knowledge about this green leafy vegetable. This study describes useful information about *I. aquatica*, current research carried out so far and some of the medicinal properties identified and explored. This plant can be used as a natural source and as a nutraceutical substitute or as a functional food.

**Key words:** *Ipomoea aquatica*, bioactive compounds, antioxidant activity, hypoglycemic, anticancer

### **INTRODUCTION**

Food science has a growing literature about fruits and vegetables imparting greater benefits to humans. Considerable epidemiological evidence suggests an association between consumption of a diet high in fruits and vegetables and decreased risk of cardiovascular diseases, hypertension, diabetes, stroke and various forms of cancer (Brandt *et al.*, 2004; Craig, 1999). Although fruits and vegetables account for only 10% of total calories consumed, they make a significant contribution to overall health. Fruits and vegetables have a regular place in traditional Indian cuisine.

Among various types of vegetables, leafy vegetables are most commonly consumed in ancient Indian daily diet. The importance of leafy vegetables in the developing countries has been recognized only now due to their nutritional and medicinal value. Green leafy vegetables occupy an important place among food crops as these provide adequate amounts of crude fiber, carotene, a precursor of vitamin A, vitamin C, riboflavin, folic acid and mineral salts like calcium, iron, phosphorus etc. They are also fair sources of proteins, containing about 2-7% and they are equal to legumes, soybeans or whole egg (Aletor *et al.*, 2002; Rao *et al.*, 1990). They form cheap and best source of food. Green leafy vegetables are highly seasonal and are available in plenty at a particular season and can be easily cooked. However, edible green leafy

vegetables appear to be underutilized through out the world and, in some area, may even be diminished in use. Green leafy vegetables are rich source of a number of micronutrients and phytochemicals and it was the phytochemicals that appeared to provide much of the disease fighting power. In this context, this study provides some useful information of *I. aquatica*, about its nutritional importance, current work carried out and some important medicinal properties.

**Classification:** *Ipomoea aquatica* Forsk. belongs to the Class: Magnoliopsida; Order: Solanales; Family: Convolvulaceae; Sub class: Asteridae (Pullaiah, 1998).

**Botanical description:** *I. aquatica* is an aquatica or semi aquatica plant, trailing or floating, herbaceous, sometimes annual or perennial, with long, hollow stem possessing large number of air passages and rooting at the nodes, found throughout India (Fig. 1). Leaves are elliptic or ovate-oblong, cordate; flowers are infundibuliform, 2-2.5 cm long, white or pale purple, solitary; fruit is a capsule (Anonymous, 1959; Edie and Ho, 1969; Gamble, 1921; Payne, 1956; Synder *et al.*, 1981).

*I. aquatica* is found trailing on moist soil or mud along the margins of stagnant streams, fresh water ponds, ditches, marshes and wet rice fields; sometimes found floating on water bodies. It occurs both in wild and cultivated state and is easily propagated by cuttings. It



Fig. 1: *Ipomoea aquatica* in its natural habitat

grows rapidly producing dense masses of foliage within a few weeks of planting. It is one of the most popular green leafy vegetable. Its commonly called as aquatica morning glory, Chinese water spinach, kangkong, morning glory, swamp cabbage, swamp morning glory, water convolvulus, water spinach (Anonymous, 1959; Candlish *et al.*, 1987; Chen *et al.*, 1991; Edie and Ho, 1969; Payne, 1956; Synder *et al.*, 1981; Wills *et al.*, 1984).

**Origin and distribution:** *I. aquatica* is supposed to be originated in China (Edie and Ho, 1969). It is distributed through out India, Sri Lanka, Tropical Asia, Africa and Australia (Kritkar and Basu, 1952). The plant is grown wildly as weed in India and USA (Anonymous, 1959; Reed, 1977) while in South East Asia like Malaysia, China, Hong Kong, Singapore and Indonesia, the plant is grown commercially (Candlish *et al.*, 1987; Chen *et al.*, 1991).

**Composition:** The leaves of *I. aquatica* contain the following: moisture 90%, protein 3%, fiber 0.9%, fat 0.4%,

carbohydrate 4.3%, mineral matter 2%, nicotinic acid: 0.6 mg/100 g; riboflavin: 120 mg/100 g; vitamin C: 137 mg/100 g; vitamin E: 11 mg/100 g and ash: 1.4% (Anonymous, 1959). The young terminal shoots and leaves of *I. aquatica* are eaten as green leafy vegetable and in salads (Ismail *et al.*, 2004) and as fodder (Phimmasan *et al.*, 2004). *I. aquatica* leaves are also very rich in proteins (Ngamsaeng *et al.*, 2004), carotenes (Chen and Chen, 1992), amino acids like aspartic acid, threonine, serine, glutamic acid, proline, glycine, alanine, leucine, tyrosine, lysine, histidine and arginine (Rao and Vijay, 2002), minerals like sodium, potassium, calcium, iron, magnesium and zinc (Duc *et al.*, 1999), sugars like glucose, fructose, sucrose (Wills *et al.*, 1984), fiber, lipids and fats (Imbs and Pham, 1995), organic acids like malic acid, citric acid, oxalic acid (Wills *et al.*, 1984), vitamins (Duc *et al.*, 1999), starch (Candlish *et al.*, 1987), polyphenols like myricetin, quercetin, luteolin, apigenin, kaempferol (Chu *et al.*, 2000; Daniel, 1989; Miean and Mohamed, 2001), dihydroquercetin glycoside (Prasad *et al.*, 2005a) and ash (Ogle *et al.*, 2001) contents.

**Nutrition:** Moisture, nitrogen, protein and amino acids like aspartic acid, threonine, serine, glutamic acid, proline, glycine, alanine, cysteine, valine, methionine, isoleucine, leucine, tyrosine, lysine, histidine and arginine content of *I. aquatica* by Sutaria and Diego (1982) were estimated using AOAC methods. Alpha-tocopherol content of *I. aquatica* with other leafy vegetables were analysed by Candlish (1983). Wills *et al.* (1984) studied vitamins like vitamin C, thiamin, riboflavin, niacin, protein, fat, sugars like glucose, fructose, sucrose, starch, dietary fiber, organic acids like malic acid, citric acid, oxalic acid, ash and minerals like sodium, potassium, calcium, iron, magnesium and zinc contents for 15 Chinese vegetables including *I. aquatica*. Candlish *et al.* (1987) studied the dietary fiber (non-cellulose polysaccharide, cellulose, lignin) and starch content in vegetables that are extensively grown and consumed in Southeast Asia including *I. aquatica*. Rao *et al.* (1990) estimated the lipid contents like non polar lipids, glycolipids, phospholipids, fatty acids, amino acids like aspartine, threonine, serine, glycine, proline, alanine, leucine, tyrosine, histidine, aspergine, argenine, minerals like calcium, magnesium, iron, zinc and copper of *I. aquatica*.

Imbs and Pham (1995) analyzed total lipids, fatty acid, triglycerides content and phospholipid content of *I. aquatica*. Iron and vitamin C content of *I. aquatica* grown in three sites of Vietnam were investigated by Duc *et al.* (1999). The nutritional value of *I. aquatica* with yielding crops such as rice, sugarcane and maize were compared by Munger (1999). It is urged that investment in crops such as *I. aquatica* may be a suitable and cost-effective way to supplement the caloric and nutritional

value of current crop production. Micronutrient composition like ash, calcium, iron, zinc, carotenes like  $\alpha$  and  $\beta$  carotenes by Ogle *et al.* (2001) were determined. Iron, calcium,  $\beta$  carotene, ascorbic acid and oxalic acid content of *I. aquatica* consumed by the tribals of Purnia district of Bihar, India were analyzed by Rao and Vijay (2002). The dieting pattern of rabbits fed with water spinach were investigated by Phimmasan *et al.* (2004). Ngamsaeng *et al.* (2004) analyzed water spinach as protein supplements for ducks feed as basal diet.

**Carotenes:** Ortaliza *et al.* (1969) estimated the carotene contents and its availability of *I. aquatica*. The effects of *I. aquatica* on cholesterol metabolism in rats were studied by Chen *et al.* (1984). Chen *et al.* (1991) characterized the major carotenoids in water convolvulus (*Ipomoea aquatica*) by open-column, thin layer and high performance liquid chromatography. Comparing the absorption spectra and retention time with reference standards the compounds were identified as  $\beta$ -carotene, cryptoxanthin, lutein, lutein epoxide, violoxanthin and neoxanthin. Carotene composition and contents of *I. aquatica* were analyzed by Tee and Lim (1991) by AOAC and HPLC methods. They found that AOAC method gave a falsely elevated result, where as the HPLC method successfully separated the major carotenoids. The carotenoids identified were lutein and  $\beta$ -carotene and were compared to that of reference standards.

Carotenoids and chlorophylls in *I. aquatica* were determined by Chen and Chen (1992) using liquid chromatography. These carotenes and chlorophylls were analyzed by High Performance Liquid Chromatography (HPLC). Fourteen peaks were detected in HPLC, of which

12 pigments were identified. These pigments include neoxanthin, violaxanthin, cryptoxanthin, lutein epoxide, lutein, cis-lutein, chlorophyll b, chlorophyll b<sup>1</sup>, chlorophyll a,  $\beta$ -carotene and cis  $\beta$ -carotene. These samples were then compared with standards by thin layer chromatography.

Tee *et al.* (1996) investigated the biological utilization of carotenoids of *Ipomoea aquatica* using rats and found bioavailability of the major carotenoids was high, as evidenced by the accumulation of retinol of the experimental rats in relation to crystalline retinol concentrate. Wills and Ranga (1996) analyzed carotene contents of *I. aquatica* by reverse phase HPLC and gradient elution. The carotenes identified were zeaxanthin (Fig. 2a), lutein (Fig. 2b), anthraxanthin, flavoxanthin, auroxanthin, luteoxanthin, neoxanthin (Fig. 2c),  $\beta$ -carotene (Fig. 2d), violoxanthin (Fig. 2e), cryptoxanthin (Fig. 2f), neoxanthin a and neoxanthin b. These samples were compared to standards and identified. Provitamin A and carotenoids from *I. aquatica* of different maturity and origin were investigated by Hulshof *et al.* (1997). The carotene contents were different when collected from different places. Also, the carotene content of mature plant was high compared to that of young plants. Provitamin A carotenoid content of *I. aquatica* at different maturity and origin was reported by Paul *et al.* (1997).

**Flavonoids:** Daniel (1989) estimated the polyphenol contents of *I. aquatica* and identified quercetin 3'-methyl ether (Fig. 3a), quercetin 4'-methyl ether (Fig. 3b) and anthocyanins. Flavonoid contents of *I. aquatica* like mycetin, quercetin, luteolin and apigenin content were




Fig. 2: Carotenes from *Ipomoea aquatica* (a) Zeaxanthin, (b) Lutein, (c) Neoxanthin, (d)  $\beta$ -carotene, (e) Violoxanthin and (f) Cryptoxanthin




Fig. 3: Flavonoids from *Ipomoea aquatica*. (a) Quercetin 3'-methyl ether, (b) Quercetin 4'-methyl ether and (c) 7-O- $\beta$ -D-glucopyranosyl-dihydroquercetin-3-O- $\alpha$ -D-glucopyranoside

identified by Chu *et al.* (2000). Chlorophylls of *I. aquatica* were investigated by Miean and Mohamed (2001). Prasad *et al.* (2005a) isolated a 7-O- $\beta$ -D-glucopyranosyl-dihydroquercetin-3-O- $\alpha$ -D-glucopyranoside (Fig. 3c) and it was characterized using UV, MS and NMR data.

**Alkaloids:** Seven aliphatic pyrrolidine amides with branched and linear saturated  $C_{15}$ - $C_{19}$  acyl moieties were detected by Tofern *et al.* (1999) in the vegetative part (stem and leaves) of *I. aquatica*. One of the compounds was isolated and characterized as 1-(14-methylhexadecanoyl) pyrrolidine and compared with synthetic compound. Yajima and Yabuta (2001) investigated the synthesis and configuration of a novel pyrrolidine amide namely 1-(14'-methylhexadecanoyl) pyrrolidine from *I. aquatica*.

**Storage studies:** Kailasapathy and Koneshan (1986) studied the loss of ascorbate content in eight leafy vegetables including *I. aquatica* due to wilting, from harvest up to a period of 24 h at environmental (24.7-25.8°C) and under refrigeration (4.4°C) temperatures. Quality changes like browning and changes in leaf and stem tissues of *I. aquatica* during storage at various temperatures (1 and 20°C) was analyzed on 5th day by Hirata *et al.* (1987). The structural changes in leaf and

stem tissues of *I. aquatica* in relation to chilling injury and browning were investigated by Ose *et al.* (1990).

**Processing:** Candlish (1983) suggested that steaming of *I. aquatica* had highest carotene content than that of fresh and boiled one. The presence of cis-isomers of  $\beta$ -carotene in *I. aquatica* were examined before and after traditional Indonesian ways of cooking by Van der Pol *et al.* (1988). Yield of carotenoids of *I. aquatica* were analyzed by Chen and Han (1990) under different methods of cooking. Mazrizal *et al.* (1997) reported the retention of vitamin C, iron and  $\beta$ -carotene in *I. aquatica* under different cooking methods. Total carotenoid and  $\beta$ -carotene contents during processing like cooking, sun drying, frying and fermentation were investigated by Speek *et al.* (1988). Van der Pol *et al.* (1988) examined and compared the cis-carotene isomers ( $\alpha$ -carotene,  $\beta$ -carotene, neo- $\beta$ -carotene, neo-U- $\beta$ -carotene) contents of *I. aquatica* before and after cooking. Mazrizal *et al.* (1997) studied the retention of  $\beta$ -carotene of *I. aquatica* when prepared using different cooking methods like boiling, microwave steaming and frying.

**Tissue culture studies of *Ipomoea aquatica*:** Effect of light on growth and chlorophyll formation of *I. aquatica*

hairy roots were investigated by Masahiro *et al.* (1996). Methods for introducing foreign genes for *I. aquatica* and regenerating the plant by Masanori *et al.* (1997) were developed. The method comprises co-cultivating the nodes of the explant with *Agrobacterium tumefaciens* carrying the target foreign gene. Hirofumi *et al.* (2000) followed the development and characterization of a photoautotrophic cell line of *I. aquatica* hairy roots. The derived photoautotrophic hairy roots had high chlorophyll content and activities of 1,5-ribulose-biphosphate carboxylase content were more than those of the parent plant. Ninomiya *et al.* (2003) investigated the changes in chlorophyll content in phototautotrophic hairy roots of *I. aquatica*. Callus initiation and its antioxidant activity of *I. aquatica* was compared with that of the mother plant and it was found that the antioxidant activity was high in callus cultures (Prasad *et al.* 2006).

**Antioxidant studies:** Ismail *et al.* (2004) estimated the antioxidant and phenolic contents of *I. aquatica*. Antioxidant activity of leaf extracts and purified compound isolated from *I. aquatica* were determined by Prasad *et al.* (2004, 2005 a) by DPPH, metal chelation and lipid peroxidation methods.

**Medicinal importance:** In Ayurveda (Indian traditional medicine), the extracts of leaves of *I. aquatica* are orally administered to alleviate the disorders like jaundice and nervous debility. This plant is used medicinally in Southeastern Asia. It is effectively used against nosebleed and high blood pressure (Duke and Ayensu, 1985; Perry, 1980). The juice of *I. aquatica* is used as emetic in cases of opium and arsenic poisoning (Anonymous, 1959; Chopra *et al.*, 1956). Dried juice has purgative properties (Anonymous, 1959; Chopra *et al.*, 1956; Nadkarni, 1954). Leaves and stems are said to be cooling. In Assam, the plant is given for nervous and general debility (Anonymous, 1959; Chopra *et al.*, 1956). It is used also for piles (Anonymous, 1959), as anthelmintic (Anonymous, 1959; Nadkarni, 1954), used in leucoderma, leprosy, jaundice and liver complaints (Nadkarni, 1954).

This plant is supposed to possess an insulin-like principle according to indigenous medicine in Sri Lanka (Jayaweera, 1982). The medicinal uses of hitherto report includes: effect on liver diseases (Badruzzaman and Husain, 1992; Nadkarni, 1954), eye diseases (Jain and Verma, 1981), constipation (Samuelsson *et al.*, 1992) and the inhibition of prostaglandin synthesis (Tseng *et al.*, 1992). Mackeen *et al.* (1997) demonstrated moderate antinematodal activity of *I. aquatica* against pine wood nematode, *Bursaphelenchus xylophilus*. The aqueous

extract of this plant also possess hypoglycemic effect (Malalavidhane *et al.*, 2000, 2001, 2003). This plant also possesses antioxidant activity (Prasad *et al.*, 2004, 2005 a). This plant also shows moderate anticancer activity against Vero, Hep-2 and A-549 cancer cell lines (Prasad *et al.*, 2005 b).

## REFERENCES

- Aletor, O., A.A. Oshodi and K. Ipinmoroti, 2002. Chemical composition of common leafy vegetables and functional properties of their leaf protein concentrates. Food Chem., 78: 63-68.
- Anonymous, 1959. Wealth of India, Raw Materials. CSIR, New Delhi, 5: 237.
- Badruzzaman, S.M. and W. Husain, 1992. Some aquatica and marshy land medicinal plants from Hardoi district of Uttarpradesh. Fitoterapia, 63: 245-247.
- Brandt, K., L.P. Christensen, J. Hansen-moller, S.L. Hansen, J. Haraldsdottir, J. Jespersen, S. Purup, A. Kharazmi, V. Barkholt, H. Frokiaer and M. Kobaek-Larsen, 2004. Health promoting compounds in vegetables and fruits: A systematic approach for identifying plant components with impact on human health. Trends Food Sci. Technol., 15: 384-393.
- Candlish, J.K., 1983. Tocopherol content of some Southeast Asian Foods. J. Agric. Food Chem., 31: 166-168.
- Candlish, J.K., L. Gourley and H.P. Lee, 1987. Dietary fiber and starch contents of some Southeast Asian vegetables. J. Agric. Food Chem., 35: 319-321.
- Chen, M.L., D.F. Chia and J.Q. Run, 1984. Effect of dietary vegetable (Water convolvulus) on cholesterol metabolism in rats. J. Nutr., 114: 503-510.
- Chen, B.H. and L.H. Han, 1990. Effects of different cooking methods on the yield of carotenoids in water convolvulus (*Ipomoea aquatica*). J. Food Prot., 53: 1076-1078.
- Chen, B.H., S.H. Yang and I.H. Han, 1991. Characterization of major Carotenoids in water convolvulus (*Ipomoea aquatica*) by open-column, thin layer and high performance liquid chromatography. J. Chromatogr., 543: 147-155.
- Chen, B.H. and Y.Y. Chen, 1992. Determination of carotenoids and chlorophylls in water convolvulus (*Ipomoea aquatica*) by liquid chromatography. Food Chem., 45: 129-134.
- Chopra, R.N., S.L. Nayar and I.C. Chopra, 1956. Glossary of Indian Medicinal Plants. CSIR, New Delhi.
- Chu, Y.H., C.L. Chang and H.F. Hsu, 2000. Flavonoid content of several vegetables and their antioxidant activity. J. Sci. Food Agric., 80: 561-566.

- Craig, W.J., 1999. Health promoting properties of common herbs. Am. J. Clin. Nutr., 70 (suppl): S491-S499.
- Daniel, M., 1989. Polyphenols of some Indian vegetables. Curr. Sci., 58: 1332-1333.
- Duc, B.M., D. Humphries, I.T.B. Mai, A.H. Dao, T.M. Co, H.H. Nga and P.T. Kim, 1999. Iron and vitamin C content of commonly consumed foods in Vietnam. Asia-Pacific J. Clin. Nutr., 8: 36-38.
- Duke, J.A. and E.S. Ayensu, 1985. Medicinal Plants of China. Reference Publ., Algonac, MI.
- Edie, H.H. and B.W.C. Ho, 1969. *Ipomoea aquatica* as a vegetable crop in Hong Kong. Econ. Bot., 23: 32-36.
- Gamble, J.S., 1921. Flora of the Presidency of Madras, India.
- Hirata, K., K. Chachin and T. Iwata, 1987. The quality changes of some vegetables used in the tropical and subtropical area during storage at various temperatures. J. Jap. Soc. Hortic. Sci., 34: 566-573.
- Hirofumi, N., T. Masaki and T. Masahito, 2000. Development and characterization of a photoautotrophic cell line of Pak-Bung hairy roots. J. Biosci. Bioeng., 89: 151-156.
- Hulshof, P.J.M., C. Xu, P. Bovenkamp, P. Muhilal and C.E. West, 1997. Application of a validated method for the determination of provitamin A carotenoids in Indonesian foods of different maturity and origin. J. Agric. Food. Chem., 45: 1174-1179.
- Imbs, A.B. and L.Q. Pham, 1995. Lipid composition of ten edible seed species from North Vietnam. J. Am. Oil Chem. Soc., 72: 957-961.
- Ismail, A., Z.M. Marjan and C.W. Foong, 2004. Total antioxidant activity and phenolic contents in selected vegetables. Food Chem., 87: 581-586.
- Jain, S.P. and D.M. Verma, 1981. Medicinal plants in the folklore of Northeast Haryana. Natl. Acad. Sci. Letters (India), 4: 269-271.
- Jayaweera, D.M.A., 1982. Medicinal plants used in Ceylon. Part II, National Science Council, Colombo, Srilanka.
- Kailasapathy, K. and T. Koneshan, 1986. Effect of wilting on the ascorbate content of selected fresh green leafy vegetables consumed in Srilanka. J. Agric. Food Chem., 34: 259-261.
- Kritkar and Basu, 1952. Indian Medicinal Plants. Parbani Press, Kolkata.
- Mackeen, M.M., M.A. Ali, M.A. Abdullah, R.M. Nasir, N.B. Mat, R.R. Abdul and K. Kawazu, 1997. Antinematodal activity of some Malaysian plant extracts against the pine wood nematode, *Bursaphelenchus xylophilus*. Pestic. Sci., 51: 165-170.
- Malalavidhane, T.S., S.M. Wickramasinghe and E.R. Jansz, 2000. Oral hypoglycemic activity of *Ipomoea aquatica*. J. Ethnopharmacol., 72: 293-298.
- Malalavidhane, T.S., S.M. Wickramasinghe and E.R. Jansz, 2001. An aqueous extract of the green leafy vegetable *Ipomoea aquatica* is as effective as the oral hypoglycemic drug tolbutamide in reducing the blood sugar levels of wistar rats. Phytother. Res., 15: 635-637.
- Malalavidhane, T.S., S.M. Wickramasinghe, M.S. Perera and E.R. Jansz, 2003. Oral hypoglycemic activity of *Ipomoea aquatica* in streptozotocin-induced, diabetic wistar rats and type ii diabetes. Phytother. Res., 17: 1098-1100.
- Masahiro, K.O., N. Hirofumi, T. Masahito and T. Setsuji, 1996. Effect of light irradiation on growth and chlorophyll formation of Pakbung green hairy roots. J. Chem. Eng. Jap., 29: 1050-1054.
- Masanori, F., N. Atsuhiko and Y. Kazuya, 1997. Methods for introducing foreign genes into tropical aquatic plant *Ipomoea aquatica* and regenerating the plant. Patent CA Section 3.
- Mazrizal, A.M., D.W. Giraud and J.A. Driskell, 1997. Retention of vitamin C, iron and  $\beta$  carotene in vegetables prepared using different cooking methods. J. Food Qual., 20: 403-418.
- Miean, K.H. and S. Mohamed, 2001. Flavonoid (myricetin, quercetin, kaempferol, luteolin and apigenin) content of edible tropical plants. J. Agric. Food Chem., 49: 3106-3112.
- Munger, H.M., 1999. Enhancement of horticulture crops for improved health. Hortic. Sci., 34: 1158-1159.
- Nadkarni, 1954. Indian Materia Medica. 3rd Edn. Popular Books, Bombay.
- Ngamsaeng, A., S. Thy and T.R. Preston, 2004. Duckweed (*Lemna minor*) and water spinach (*Ipomoea aquatica*) as protein supplements for ducks fed broken rice as the basal diet. Livestock Res. Rural Dev., 16: 18-24.
- Ninomiya, K., Y. Oogami, M. Kino-Oka and M. Taya, 2003. Assessment of herbicidal toxicity based on non destructive measurement of local chlorophyll content in photoautotrophic hairy roots. J. Biosci. Bioeng., 95: 264-270.
- Ogle, B.M., A.D. Ha-Thi, G. Mulokozi and L. Hambræus, 2001. Micronutrient composition and nutritional importance of gathered vegetables in Vietnam. Int. J. Food Sci. Nutr., 52: 485-499.
- Ortaliza, I.C., I.F. Del Rosario, M.M. Caedo and A.P. Alcaraz, 1969. The availability of carotene in some Philippine vegetables. Phillipine J. Sci., 98: 123-131.
- Ose, H.K., K. Chachin and T. Iwata, 1990. Structural changes in cells and chilling injury of water convolvulus (*Ipomoea aquatica*) stored at low temperature. J. Jap. Soc. Hortic. Sci., 59: 171-177.

- Paul, J.M.H., C. Xu, P.V.D. Bovenkamp, A. Muhilal and C.E. West, 1997. Application of a validated method for the determination of pro-vitamin A Carotenoids in Indonesian foods of different maturity and origin. *J. Agric. Food Chem.*, 45: 1174-1179.
- Payne, W.J., 1956. *Ipomoea reptans*, Poir. A useful tropical fodder plant. *Trop. Agric. Trin.*, 33: 302-305.
- Perry, L.M., 1980. Medicinal Plants of East and Southeast Asia: Attributed Properties and Uses. MIT Press, Cambridge, MA.
- Phimmasan, H., S. Kongvongxay, P. Chhayty and T.R. Preston, 2004. Water spinach (*Ipomoea aquatica*) and Stylo 184 (*Stylosanthes guianensis* CIAT 184) as basal diets for growing rabbits. *Livestock Res. Rural Dev.*, 16: 46-59.
- Prasad, N.K., A.P. Kulkarni, M.S. Mahesh, S.M. Aradhya and G.R. Shivamurthy, 2004. Studies On antioxidant activity of selected vegetables using *in vitro* model. *Soc. Free Radical Res. India Bull.*, 3: 23-26.
- Prasad, N.K., S. Divakar, G.R. Shivamurthy and S.M. Aradhya, 2005a. Isolation of a free radical scavenging antioxidant from water spinach (*Ipomoea aquatica* Forsk.). *J. Sci. Food Agric.*, 85: 1461-1468.
- Prasad, N.K., R. Chandrashekar, G. Ashok, G.R. Shivamurthy, P. Vijayan and S.M. Aradhya, 2005b. Cytotoxic Properties of *Ipomoea aquatica* Forsk. Leaf. *Ind. J. Pharmacol.*, 37 (6): 397-398.
- Prasad, N.K., M. Shiva Prasad, S.M. Aradhya and G.R. Shivamurthy, 2006. Callus induction from *Ipomoea aquatica* Forsk. leaf and its antioxidant activity. *Ind. J. Biotechnol.*, 5: 107-111.
- Pullaiah, 1998. Taxonomy of Angiosperms. Regency Publications. New Delhi.
- Rao, K.S., R. Dominic, K. Singh, C. Kaluwin, E. Donals, R. Gwyn and P. Jones, 1990. Lipid, fatty acid, amino acid and mineral composition of five edible plant leaves. *J. Agric. Food Chem.*, 38: 2137-2139.
- Rao, T.V.R.K. and T. Vijay, 2002. Iron, calcium,  $\beta$ -carotene, ascorbic acid and oxalic acid contents of some less common leafy vegetables consumed by the tribals of purnia district of Bihar. *J. Food Sci. Technol.*, 39: 560-562.
- Reed, C.F., 1977. Economically important foreign weeds: Potential problems in the United States. W.S.D.A, Agric. Hand book.
- Samuelsson, G., M.H. Farah, P. Claeson, M. Hagos, M. Thulin, O. Hedberg, A.M. Warfa, A.O. Hassan, A.H. Elmi, A.D. Abdurahman, A.S. Elmis and Y.A. Abdi, 1992. Inventory of plants used in traditional medicine in Somalia: II. Plants of the families Combartaceae to Labiatae. *J. Ethnopharmacol.*, 37: 47-70.
- Speek, A.J.S. Speek and W.H.P. Schreurs, 1988. Total carotenoid and  $\beta$ -carotene contents of Thai vegetables and the effects of processing. *Food Chem.*, 27 (4): 245-257.
- Sutaria, P.B. and M.L. Diego, 1982. Essential amino acid analysis of selected Philippine vegetables and fruits. *The Philippine J. Sci.*, 3: 45-55.
- Snyder, G.H., J.F. Morton and W.G. Genung, 1981. Trials of *Ipomoea aquatica*, nutritious vegetable with high protein and nitrate extraction potential. *Proc. Fla. State Hortic. Soc.*, 94: 230-235.
- Tee, E.S. and C.H. Lim, 1991. Carotenoids composition and content of Malaysian vegetables and fruits by the AOAC and HPLC methods. *Food Chem.*, 41: 309-339.
- Tee, E.S., C.L. Lim, Y.H. Chong and S.C. Khor, 1996. A study of the biological utilization of carotenoids of carrots and swamp cabbage in rats. *Food Chem.*, 56: 21-32.
- Tofern, B., P. Mann, M. Kaloga, K.J. Siems, L. Witte and E. Eich, 1999. Aliphatic Pyrrolidine Amides from two tropical Convolvulaceous species. *Phytochemistry*, 52: 1437-1441.
- Tseng, C.F., S. Iwakami, A. Mikajiri, M. Shibuya, F. Hanaoka, Y. Ebizuka, K. Padmavinata and U. Sankawa, 1992. Inhibition of *in vitro* prostaglandin and leukotriene biosynthesis by cinnamoyl- $\beta$ -phenethylamine and n-acyl dopamine derivatives. *Chem. Pharm. Bull.*, 40: 396-400.
- Van Der Pol, F., S.U. Purnomo and H.A. Van Rosmalen, 1988. Trans-Cis isomerisation of Carotenoids and its effect on the vitamin A potency of some common Indonesian foods. *Nutr. Rep. Int.*, 37: 785-793.
- Wills, R.B.H., A.W.K. Wong, F.M. Scriven and H. Greenfield, 1984. Nutrient composition of Chinese vegetables. *J. Agric. Food Chem.*, 32: 413-416.
- Wills, R.B.H. and A. Ranga, 1996. Determination of carotenoids in Chinese vegetables. *J. Agric. Food Chem.*, 56: 451-455.
- Yajima, A. and G. Yabuta, 2001. Synthesis and absolute configuration of MQ-A<sub>3</sub> [1-(14'-methylhexadecanoyl) pyrrolidine], a novel aliphatic pyrrolidine amide from the tropical Convolvulaceae species. *Biosci. Biotechnol. Biochem.*, 65: 463-465.