

## Antioxidant, Analgesic and Toxic Potentiality of *Stephania Japonica* (Thunb.) Miers. Leaf

<sup>1</sup>M. Habibur Rahman, <sup>2</sup>M. Badrul Alam, <sup>3</sup>N.S. Chowdhury, <sup>1</sup>M.K. Jha, <sup>2</sup>M. Hasan,  
<sup>2</sup>M.M. Khan, <sup>2</sup>M.S. Rahman and <sup>4</sup>M. Ekramul Haque

<sup>1</sup>Department of Pharmacy, Bangladesh University, Dhaka, Bangladesh

<sup>2</sup>Department of Pharmacy, Atish Dipankar University of Science and Technology, Dhaka, Bangladesh

<sup>3</sup>Department of Pharmacy, Manarat International University, Dhaka, Bangladesh

<sup>4</sup>Department of Pharmacy, BRAC University, Dhaka, Bangladesh

**Abstract:** In the present study crude methanolic extract of *Stephania japonica* leaf was investigated for possible antioxidant, analgesic and toxic activity. The extract showed antioxidant activity in DPPH radical scavenging activity, nitric oxide scavenging activity and reducing power assays. In both DPPH radical and NO scavenging assay, the extract exhibited moderate antioxidant activity and the IC<sub>50</sub> values in DPPH radical scavenging and NO scavenging assays were found to be 105.55±1.06 and 129.12±0.15 µg mL<sup>-1</sup>, respectively while the IC<sub>50</sub> values of ascorbic acid were 12.30±0.11 and 18.64±0.22 µg mL<sup>-1</sup>, respectively. Reducing power activity of the extract increased in a dose dependent manner. Analgesic activity of the crude extract was evaluated using acetic acid-induced writhing model of pain in mice. The crude extract at 200 and 400 mg kg<sup>-1</sup> b.wt. doses displayed significant (p<0.001) reduction in acetic acid induced writhing in mice with a maximum effect of 75.89% reduction at 400 mg kg<sup>-1</sup> b.wt. which is comparable to the standard, diclofenac sodium (86.52%). The extract was also investigated for toxic potentiality using Brine Shrimp lethality bioassay. In this bioassay the extract showed significant toxicity to Brine Shrimp nauplii with the LC<sub>50</sub> value of 25.19±0.98 µg mL<sup>-1</sup>. The study clearly indicates that the extract possesses good analgesic and cytotoxic activity along with moderate antioxidant potential.

**Key words:** Antioxidant, analgesic, *in vitro*, *in vivo*, *Stephania japonica*, toxicity

### INTRODUCTION

Free radicals cause depletion of immune system antioxidants, change in gene expression and induce abnormal proteins and contribute to more than one hundred disorders in humans including atherosclerosis, arthritis, ischemia and reperfusion injury of many tissues, central nervous system injury, gastritis, cancer and AIDS (Hela and Abdullah, 2010; Kumpulainen and Salonen, 1999). Although organisms have endogenous antioxidant defenses produced during normal cell aerobic respiration against ROS, other antioxidants are taken both from natural and synthetic origin. Synthetic antioxidants are widely used but their use is being restricted now-a-days because of their toxic and carcinogenic effects. Thus, interest in finding natural antioxidants, without any undesirable effect, has increased greatly (Rechner *et al.*, 2002). Some natural antioxidant (e.g., rosemary and sage) are already exploited commercially either as antioxidant additives or a nutritional supplements but generally there is still a

demand to find more information concerning the antioxidant potential of plant species. It has been mentioned the antioxidant activity of plants might be due to their phenolic, flavonoid, tannin and proanthocyanidin compounds (Hela and Abdullah, 2010; Frankle and Meyer, 2000). Pain is a sensorial modality and primarily protective in nature but often causes discomfort. It is the most important symptom that brings the patient to physician. Analgesics relieve pain as a symptom without affecting its cause (Tripathi, 1999). Currently available analgesic drugs such as opiates and NSAIDs are not useful in all cases due to their adverse effects. In this respect new compounds with improved pain management capacity and fewer side effects are being sought with urgency.

*Stephania japonica* (Thunb.) Miers, (Family-Menispermaceae) is a slender twining shrub with greenish yellow flowers and large tubers. It is native to eastern and southern Asia and Australasia. In Bangladesh, it is grown in many areas. The leaves and roots are bitter and astringent and used in fevers, diarrhea, dyspepsia and urinary disease (Ghani, 2003).

As a part of present ongoing investigations on local medicinal plants of Bangladesh (Mazumder *et al.*, 2009) in this study, we reported antioxidant, analgesic and toxic potentiality of the leaves of *Stephania japonica*.

## MATERIALS AND METHODS

**Plant materials:** The plant, *Stephania japonica* (Thumb.) Miers was collected from the village Uthali under Jibannagar thana of Chuadanga district, Bangladesh during the month of August 2009. The plants were mounted on paper and the sample was identified by Mrs. Mahmuda Begum, senior scientific officer, Bangladesh National Herbarium, Dhaka, where the voucher specimen has been deposited. Its DACB Accession No. is 34,393.

**Chemicals:** Sodium nitroprusside were purchased from E. Merck (Germany). 1,1-diphenyl-2-picryl-hydrazyl (DPPH), sodium nitroprusside, ascorbic acid, quercetin and potassium ferric cyanide were purchased from Sigma Chemical Co. Ltd, (St. Louis, MO, USA). Diclofenac-Na was collected from Square Pharmaceuticals Ltd., Bangladesh. All other chemicals and reagents were of analytical grade.

**Preparation of plant extract:** The plant material was shade-dried with occasional shifting and then powdered with a mechanical grinder, passing through sieve #40 and stored in a tight container. The dried powder material (1.5 kg) was refluxed with MeOH for three hours. The total filtrate was concentrated to dryness, *in vacuo* at 40°C to render the MeOH extract for investigation.

**Animal:** For the experiment, Twenty swiss albino mice of either sex, 3-4 weeks of age, weighing between 20-25 g, were collected from the animal research branch of the International Center for Diarrheal Disease and Research, Bangladesh (ICDDR). Animals were maintained under standard environmental conditions (temperature: 24.0±1.0°C), relative humidity: 55-65% and 12 h light/ dark cycle) and had free access to feed and water *ad libitum*. The animals were acclimatized to laboratory condition for one week prior to experiments. All protocols for animal experiment were approved by the institutional animal ethical committee.

**Phytochemical screening:** The freshly prepared crude extract was qualitatively tested for the presence of chemical constituents. Phytochemical screening of the extract was performed using the following reagents and chemicals: Alkaloids with Dragendorff's reagent,

flavonoids with the use of Mg and HCl; tannins with ferric chloride and potassium dichromate solutions and saponins with ability to produce stable foam and steroids with Libermann- Burchard reagent. Reducing sugars with Benedict's reagent. These were identified by characteristic color changes using standard procedures (Ghani, 2003).

### *In vivo* analgesic screening

**Acetic acid-induced writhing test:** The analgesic activity of the samples was also studied using acetic acid-induced writhing model in mice. Test samples and vehicle were administered orally 30 min before intraperitoneal administration of 0.7% acetic acid but Diclofenac-Na was administered intraperitoneally 15 min before injection of acetic acid. After an interval of 5 min, the mice were observed for specific contraction of body referred to as 'writhing' for the next 10 min (Ahmed *et al.*, 2004).

### *In vitro* tests for antioxidant activity

**Free radical scavenging activity measured by 1,1-diphenyl-2-picryl-hydrazyl (DPPH):** The free radical scavenging activity of MeOH extract based on the scavenging activity of the stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical was determined by the method described by Braca *et al.* (2001). Plant extract (0.1 mL) was added to 3 mL of a 0.004% MeOH solution of DPPH. Absorbance at 517 nm was determined after 30 min and the percentage inhibition activity was calculated from  $[(A_0 - A_1)/A_0] \times 100$ , where  $A_0$  is the absorbance of the control and  $A_1$  is the absorbance of the extract/ standard.  $IC_{50}$  value was calculated from the equation of line obtained by plotting a graph of concentration ( $\mu\text{g mL}^{-1}$ ) versus % inhibition.

**Nitric oxide radical scavenging assay:** The procedure is based on the method, where sodium nitroprusside in aqueous solution at physiological pH spontaneously generates nitric oxide, which interacts with oxygen to produce nitrite ions that can be estimated using Griess reagent. Scavengers of nitric oxide compete with oxygen leading to reduced production of nitrite ions. For the experiment, sodium nitroprusside (10 mM) in phosphate buffered solution (pH 7.4) was mixed with different concentrations of plant extract of *S. japonica* dissolved in 10% DMSO and incubated at room temperature for 150 min. The same reaction mixture without the extract but the equivalent amount of the solvent used served as the control. After incubation, 0.5 mL of Griess reagent (1% sulfanilamide, 2%  $\text{H}_3\text{PO}_4$  and 0.1% N-(1-naphthyl) ethylenediamine dihydrochloride was added. The absorbance was measured at 546 nm and the percentage

inhibition activity was calculated from  $[(A_0 - A_1)/A_0] \times 100$ , where,  $A_0$  is the absorbance of the control and  $A_1$  is the absorbance of the extract/ standard (Sreejayan, 1997).  $IC_{50}$  value was calculated from the equation of line obtained by plotting a graph of concentration ( $\mu\text{g mL}^{-1}$ ) versus % inhibition.

**Reducing power activity:** The reducing power of *S. japonica* was determined by according to the method previously described by Oyaizu (1986). Extracts at different concentrations in 1 mL of 10% DMSO were mixed with 2.5 mL of phosphatebuffer (0.2M, pH 6.6) and 2.5 mL potassium ferricyanide  $[\text{K}_3\text{Fe}(\text{CN})_6]$  (1%) and then the mixture was incubated at  $50^\circ\text{C}$  for 30 min. Afterwards, 2.5 mL of trichloroacetic acid (10%) was added to the mixture which was then centrifuged at 3000 rpm for 10 min. Finally, 2.5 mL of upper layer solution was mixed with 2.5 mL distilled water and 0.5 mL  $\text{FeCl}_3$  (0.1%) and the absorbance was measured at 700 nm. Increased absorbance of the reaction mixture indicated increased reducing power.

**Brine shrimp lethality bioassay:** The cytotoxic activity of the plant was evaluated using Brine Shrimp lethality bioassay method (Meyer *et al.*, 1982) where 6 graded doses (viz., 5, 10, 20, 50, 100 and  $200 \mu\text{g mL}^{-1}$ ) were used. Brine shrimps (*Artemia salina* Leach) nauplii Ocean 90, USA were used as test organisms. For hatching, eggs were kept in brine with a constant oxygen supply for 48 h. The nature nauplii were then used in the experiment. DMSO was used as a solvent and also as a negative control. The median lethal concentration  $LC_{50}$  of the test sample after 24 h was obtained by a plot of percentage of the dead shrimps against the logarithm of the sample concentration. Vincristine sulfate was used as a reference standard in this case.

**Statistical analysis:** All the *in vitro* experimental results were Mean $\pm$ SEM of three parallel measurements. Results of *in vivo* study were given as Mean $\pm$ SEM. and data were evaluated by using student's ttest. pvalues $<$ 0.001 were regarded as significant.

## RESULTS

**Phytochemical screening:** Phytochemical analyses of the crude extract revealed the presence of steroid, alkaloid, tannin and glycoside (Table 1).

### *In vivo* Analgesic screening

**Acetic acid-induced writhing test:** Table 2 shows the effects of the extract of on acetic acid-induced writhing in

Table 1: Result of chemical group tests of the methanol extract of *Stephania japonica* leaf

Sample	Reducing						
	Alkaloid	Steroid	Flavonoid	sugar	Tannin	Saponin	Glycoside
ME of <i>Stephania japonica</i>	+++	++	-	-	+	-	+++

ME: Methanolic extract, +: Present, -: Absent; +++: Reaction intensity is high, ++: Reaction intensity is medium, +: Reaction intensity is normal

Table 2: Effects of the methanolic extract of *Stephania japonica* leaf on acetic acid-induced writhing in mice

Groups	Dose ( $\text{mg kg}^{-1}$ )	No. of writhing	Percentage of protection
I	Vehicle	28.2 $\pm$ 0.748	-
II	10	3.8 $\pm$ 1.008**	86.52
III	200	11.8 $\pm$ 0.765**	58.16
IV	400	6.8 $\pm$ 0.637**	75.89

Values are Mean $\pm$ SEM, (n =5); \*\*p $<$ 0.001, Student's t test as compared to vehicle control. Group-I animals received vehicle (1% Tween 80 in water), Group-II received Diclofenac-Na 10  $\text{mg kg}^{-1}$  body weight, Group-III and Group-IV were treated with 200 and 400  $\text{mg kg}^{-1}$  body weight (p.o.) of the crude extract of *S. japonica* leaf

Table 3: Antioxidant activities of the *S. japonica* leaf extract on DPPH and NO

Sample	<sup>a</sup> DPPH	<sup>b</sup> NO
	$IC_{50}$ ( $\mu\text{g mL}^{-1}$ )	$IC_{50}$ ( $\mu\text{g mL}^{-1}$ )
Methanolic extract of <i>S. japonica</i>	105.55 $\pm$ 1.06*	129.12 $\pm$ 0.15*
Ascorbic acid	12.30 $\pm$ 0.11	18.64 $\pm$ 0.22
Quercetin		27.69 $\pm$ 0.57

<sup>a</sup>DPPH is the free radical scavenging activity ( $IC_{50}$ :  $\mu\text{g mL}^{-1}$ ), <sup>b</sup>NO is the inhibition of nitric oxide production ( $IC_{50}$ :  $\mu\text{g mL}^{-1}$ ), \*p $<$ 0.001 by student's test for values between the sample and the control

mice. The oral administration of both doses of *S. japonica* extract significantly (p $<$ 0.001) inhibited writhing response induced by acetic acid in a dose dependent manner.

### *In vitro* antioxidant activity

**DPPH radical scavenging activity:** The percentage (%) scavenging of DPPH radical was found to be concentration dependent i.e., concentration of the extract between 10-200  $\mu\text{g mL}^{-1}$  greatly increasing the inhibition activity. The  $IC_{50}$  value of the extract was 105.55 $\pm$ 1.06  $\mu\text{g mL}^{-1}$ , as opposed to that of ascorbic acid ( $IC_{50}$  12.30 $\pm$ 0.11  $\mu\text{g mL}^{-1}$ ) which is a well known antioxidant (Table 3).

**Nitric Oxide (NO) scavenging activity:** Table 3 represents the result of nitric oxide scavenging activity ( $IC_{50}$  value) of the methanolic extract of *S. japonica* compared with various reference compounds such as ascorbic acid and quercetin.  $IC_{50}$  value of the extract of was 129.12 $\pm$ 0.15  $\mu\text{g mL}^{-1}$ , whereas, standard ascorbic acid and quercetin showed 18.64 $\pm$ 0.22 and 27.69 $\pm$ 0.57  $\mu\text{g mL}^{-1}$ , respectively.

**Reducing power ability:** For the measurement of the reductive ability, presat study investigated the  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$

Table 4: Brine shrimp lethality bioassay of crude methanol extract of *Stephania japonica* leaf

Test sample	Concentration ( $\mu\text{g mL}^{-1}$ )	Log Conc.	No. of dead shrimps (out of 10)	Mortality (%)	LC <sub>50</sub> of test sample ( $\mu\text{g mL}^{-1}$ )	LC <sub>50</sub> of Vincristine sulphate ( $\mu\text{g mL}^{-1}$ )
Methanolic extract of <i>S. japonica</i>	200	2.303	10	100	25.19±0.98	0.25±0.22
	200	2.303	10	100		
	160	0.204	10	100		
	80	1.903	9	90		
	40	0.602	7	70		
	20	0.301	4	40		
	10	1.000	2	20		




Fig. 1: Reducing power of Methanolic extract of *S. japonica*, ascorbic acid, galic acid and quercetin by spectrophotometric detection of  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  transformation. Results are Mean±SEM of three parallel measurements

transformation in the presence of methanolic extract. Like the antioxidant activity, the reducing power of *S. japonica* increased with increasing concentration of the sample. Figure 1 shows the reductive capabilities of the *S. japonica* compared with ascorbic acid, galic acid and quercetin.

**Brine Shrimp lethality bioassay:** The result of Brine Shrimp lethality bioassay is given in Table 4. *S. japonica* leaf extract displayed strong toxic potentiality. LC<sub>50</sub> value for the extract was found to be 25.19±0.98  $\mu\text{g mL}^{-1}$ .

## DISCUSSION

The oral administration of both doses of *S. japonica* extract significantly ( $p < 0.001$ ) inhibited writhing response induced by acetic acid in a dose dependent manner. Acetic acid-induced writhing model represents pain sensation by triggering localized inflammatory response. Such pain stimulus leads to the release of free arachidonic acid from tissue phospholipid (Ahmed *et al.*, 2006). The acetic acid induced writhing response is a sensitive procedure to evaluate peripherally acting analgesics. The response is thought to be mediated by peritoneal mast cells (Ribeiro *et al.*, 2000), acid sensing ion channels (Vogel and Vogel, 1997) and the prostaglandin pathways (Voilley, 2004).

Preliminary phytochemical screening showed the presence of tannin, alkaloid and glycoside in the plant extract. However, tetrandrine and fangchinoline alkaloids were isolated from *S. japonica* and showed anti-inflammatory effect through decrease leukotriene and prostaglandin generation. Furthermore tetrandrine has been shown to inhibit the production of TNF-alpha and IL-6 (Teh *et al.*, 1990; Xue *et al.*, 2008). Li *et al.* (2000) has been demonstrated that aconitum and *S. tetrandra* combinedly showed remarkable analgesic activity in rabbits and mice model. So, the observed analgesic activity may be attributed to these compounds. Moreover, recent studies suggest that the inflammatory tissue damage is due to the liberation of reactive oxygen species from phagocytes invading the inflammation sites (Parke and Sapota, 1996). Again the plant extracts demonstrated antioxidant action in the tested models. So it can be assumed that Cyclooxygenase (COX) inhibitory activity together with antioxidant activity may reduce the production of free arachidonic acid from phospholipid or may inhibit the enzyme system responsible for the synthesis of prostaglandins and ultimately relieve pain-sensation.

Polyphenolic compounds, like flavonoids, tannins and phenolic acids, commonly found in plants have been reported to have multiple biological effects, including antioxidant activity (Brown and Rice-Evans, 1998; Vinson *et al.*, 1995; Gil *et al.*, 1999; Kahkonen *et al.*, 1999). Fangchinoline and cepharanthine isolated from *Stephania rotunda* showed antioxidant activity in different *in-vitro* model (Gulcin *et al.*, 2010). Tannic acid present in the plant extract, as evident from phytochemical screening, may be responsible for the antioxidant action. NO scavenging capacity of the extract may help to arrest the chain of reactions initiated by excess generation of NO that are detrimental to the human health. Nitric oxide is also implicated for inflammation, cancer and other pathological conditions (Moncada *et al.*, 1991). A direct correlation between antioxidant capacity and reducing power of certain plant extracts has been reported (Tanaka *et al.*, 1988). The reducing properties are generally associated with the presence of reductones, which have been shown to exert antioxidant action by breaking the free radical chain by donating a hydrogen atom (Duh *et al.*, 1999).

The extract also showed significant cytotoxicity on Brine Shrimp nauplii. This may be due to the fact that *S. japonica* contains isotriline and triline, bisbenzylisoquinoline alkaloids which was previously reported to possess multidrug-resistance-reversing activity in human breast cancer cell line (Andrea and Chang, 1997).

### CONCLUSION

The study clearly indicates that the extract possesses antioxidant and cytotoxic substances. At the same time its ability to suppress abdominal writhes confirms the analgesic property of the extract. These findings justify the traditional uses of this plant in the treatment of diabetes, wounds, inflammatory conditions, worms, infestations and malarial fever. Further research is necessary for elucidating the active principles.

### REFERENCES

- Ahmed, F., M.S.T. Selim, A.K. Das and M.S.K. Choudhuri, 2004. Anti-inflammatory and antinociceptive activities of *Lippia nodiflora* Linn. *Pharmazie*, 59: 329-330.
- Ahmed, F., M.H. Hossain, A.A. Rahman and I.Z. Shahid, 2006. Antinociceptive and sedative effects of the bark of *Cerbera odollam* Gaertn. *J. Oriental Pharmacy Exp. Med.*, 6: 344-348.
- Andrea, M.H. and C.J. Chang, 1997. Multidrug-resistance modulators from *Stephania japonica*. *J. Nat. Prod.*, 60: 1193-1195.
- Braca, A., N.D. Tommasi, L.D. Bari, C. Pizza, M. Politi and I. Morelli, 2001. Antioxidant Principles from *Bauhinia terapotensis*. *J. Nat. Prod.*, 64: 892-895.
- Brown, J.E. and C.A. Rice-Evans, 1998. Luteolin rich artichoke extract protects low-density lipoprotein from oxidation *in vitro*. *Free Radical Res.*, 29: 247-255.
- Duh, P.D., Y.Y. Tu and G.C. Yen, 1999. Antioxidant activity of the aqueous extract of hamg Jyur (*Chrysanthemum morifolium* Ramat). *Lebensmittel-Wissenschaft Technol.*, 32: 269-277.
- Frankle, E.N. and A.S. Meyer, 2000. The problem in using one dimensional methods to evaluate multifunctional food and biological antioxidants. *J. Sci. Food Agric.*, 80: 1925-1941.
- Ghani, A., 2003. Medicinal Plants of Bangladesh. 2nd Revised Edn., Asiatic Society of Bangladesh, Dhaka, Bangladesh.
- Gil, M.I., F. Ferreres and F.A. Tomas-Barberan, 1999. Effect of postharvest storage and processing on the antioxidant constituents (flavonoids and vitamin C) of fresh-cut spinach. *J. Agric. Food Chem.*, 47: 2213-2217.
- Gulcin, I., R. Elias, A. Gepdiremen, A. Chea and F. Topal, 2010. Antioxidant activity of bisbenzylisoquinoline alkaloids from *Stephania rotunda*: Cepharanthine and fangchinoline. *J. Enzyme Inhibit. Med. Chem.*, 25: 44-53.
- Hela, A.E. and A. Abdullah, 2010. Antioxidant and antimicrobial activities of methanol extracts of some *Verbena* species: *In vitro* evaluation of antioxidant and antimicrobial activity in relation to polyphenolic content. *J. Applied Sci. Res.*, 6: 683-689.
- Kahkonen, M.P., A.I. Hopia, H.J. Vuorela, J.P. Rauha, K. Pihlaja, T.S. Kulaja and M. Heinonen, 1999. Antioxidant activity of plant extracts containing phenolic compounds. *J. Agric. Food Chem.*, 47: 3954-3962.
- Kumpulainen, J.T. and J.T. Salonen, 1999. Natural Antioxidants and Anticarcinogens in Nutrition, Health and Disease. Royal Society of Chemistry, UK., pp: 178-187.
- Li, X., S. Zhang, L. Qin, Z. Song, C. Mao and Q. Zhang, 2000. Experimental study of analgesic effect of *Combined aconitum* and *Stephania tetrandra*. *Chinese J. Integrative Med.*, 6: 111-111.
- Mazumder, M.E.H., M.B. Alam, M.A. Alam, M. Jamila and M.D.A. Muhsin, 2009. Investigation of secondary metabolites, antioxidant and cytotoxic activities of the leaf extract of *Portulaca oleracea* Linn. *Bangladesh J. Life Sci.*, 21: 17-22.
- Meyer, B.N., N.R. Ferrigni, J.E. Putnam, L.B. Jacobsen, D.E. Nichols and J.L. McLaughlin, 1982. Brine shrimp: A convenient bioassay for active plant constituents. *Planta Med.*, 45: 31-34.
- Moncada, S., R.M.J. Palmer and E.A. Higgs, 1991. Nitric oxide: Physiology, pathophysiology and pharmacology. *Pharmacol. Rev.*, 43: 109-142.
- Oyaizu, M., 1986. Studies on product of browning reaction prepared from glucose amine. *Jap. J. Nutr.*, 44: 307-315.
- Parke, D.V. and A. Sapota, 1996. Chemical toxicity and reactive Oxygen species. *Int. J. Occup. Med. Environ. Health*, 9: 331-340.
- Rechner, A.R., G. Kuhnle, P. Bremner, G.P. Hubbard, K.P. Moore and C.A. Rice-Evans, 2002. The metabolic fate of dietary polyphenols in humans. *Free Radic. Biol. Med.*, 33: 220-235.

- Ribeiro, R.A., M.L. Vale, S.M. Thomazzi, A.B. Paschoalato, S. Poole, S.H. Ferreira and F.Q. Cunha, 2000. Involvement of resident macrophages and mast cells in the writhing nociceptive response induced by zymosan and acetic acid in mice. *Eur. J. Pharmacol.*, 387: 111-118.
- Sreejayan, R.M.N., 1997. Nitric oxide scavenging by curcuminoids. *J. Pharm. Pharmacol.*, 49: 105-107.
- Tanaka, M., C.W. Kuie, Y. Nagashima and T. Taguchi, 1988. Applications of antioxidative Maillard reaction products from histidine and glucose to sardine products. *Nippon Suisan Gakkaishi*, 54: 1409-1414.
- Teh, B.S., W.K. Seow, S.Y. Li and Y.H. Thong, 1990. Inhibition of prostaglandin and leukotriene generation by the plant alkaloids tetrandrine and berbamin. *Int. J. Immunopharmacol.*, 12: 321-326.
- Tripathi, K.D., 1999. *Essentials of Medical Pharmacology*. 4th Edn., Jaypee Brothers Medical Publishers (P) Ltd., New Delhi, India, ISBN: 81-7179-633-8, pp: 432.
- Vinson, J.A., Y.A. Dabbagh, M.M. Serry and J. Jang 1995. Plant flavonoids, especially tea flavonols, are powerful antioxidants using an *in vitro* oxidation model for heart disease. *J. Agric. Food. Chem.*, 43: 2800-2802.
- Vogel, H.G. and W.H. Vogel, 1997. *Pharmacological Assays, Drug Discovery and Evaluation*. Springer Verlag, Germany, pp: 368-370.
- Voilley, N., 2004. Acid-sensing ion channels (ASICs): New targets for the analgesic effects of Non-Steroid Anti-Inflammatory Drugs (NSAIDs). *Curr. Drug Targets Inflamm. Allergy*, 3: 71-79.
- Xue, Y., Y. Wang, D.C. Feng, B.G. Xiao and L.Y. Xu, 2008. Tetrandrine suppresses lipopolysaccharide induced microglial activation by inhibiting NF- $\kappa$ B pathway. *Acta Pharmacol. Sin.*, 29: 245-251.