



International Journal of Pharmacology

ISSN 1811-7775

science
alert

ansinet
Asian Network for Scientific Information

Effect of Iodine Supplementation on Antioxidant Status of Normal and Alloxan Monohydrate in Toxicated Rats

¹Gabriel A. Agbor, ^{2,3}Ibrahim Taga, ²Danielle R. Nguindé, ³Mohsin A. Zaidi, ²L.G. Lehman, ³Illimar Altosaar and ⁴Jeanne Y. Ngogang

¹Centre for Research on Medicinal Plants and Traditional Medicine, Institute for Medical Research and Medicinal Plants Studies, P.O. Box 6163, Yaoundé, Cameroon

²Clinical Biochemistry and Nutrition Research Unit, Department of Biochemistry, Faculty of Science, University of Douala, Cameroon

³Biotechnology Laboratory, BMI, Faculty of Medicine, University of Ottawa, Canada

⁴Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Cameroon

Abstract: Oxidative stress is implied in numerous pathologies, including obesity, type 2 diabetes, atherosclerosis, inflammatory disorders and ageing. Antioxidants are known to neutralise the effects of oxidative stress. Hence, the present study was designed to evaluate the effect of iodine supplementation on the antioxidant status of alloxan monohydrate intoxicated rats. Two groups (A and C) of Wistar rats were kept normal (were not induced oxidative stress), while other two groups (B and D) were induced oxidative stress by alloxan monohydrate (150 mg kg⁻¹) intoxication. Group A and B animals were fed normal rodent diet while group C and D animals' diet was supplemented with iodine (KI, 25 mg kg⁻¹ diet). The feeding lasted 6 weeks and the animals were sacrificed and the plasma and red blood cells hemolysate prepared from whole blood samples were used for biochemical analysis. Alloxan monohydrate intoxication induced a collapse in the antioxidant system characterized by a significant drop (p<0.05) in antioxidant enzyme activity and ferric reducing antioxidant power as seen in group B animals. However, iodine supplementation tends to prevent this effect with significant increases (p<0.05) in bilirubin concentration, Ferric reducing antioxidant power and superoxide dismutase and catalase activities. A significant (p<0.05) reduction in the Thiobarbituric Acid Reactive Substance (TBARs), carbonyl and the thiol functionalities that were originally increased by alloxan monohydrate intoxication, further substantiate the antioxidant status of iodine. Thus, iodine has an antioxidant potential. The antioxidant potential of iodine obtained in this study is informative of the important role iodine plays in the human diet.

Key words: Oxidative stress, catalase, superoxide dismutase, carbonyl, thiol, TBARs

INTRODUCTION

Iodine is an antiseptic soluble in alcohol. The first living cells to produce oxygen earlier considered to be toxic were the algae containing high amount of iodine (Venturi and Venturi, 1999). Thus, the high iodine content played a protective antioxidant role in the algae. Iodine deficiency has been reported to activate antioxidant genes and cause DNA damage in the thyroid gland of rats and mice. Antioxidative enzymes in iodine deficient rats and mice showed increased mRNA expression. Thus, indicating increased radical burden that could lead to accumulation of oxidized adducts as seen in the thyroidal genomic DNA (Maier *et al.*, 2007). Dietary iodides have

earlier been reported to defend brain cells from lipid peroxidation (Wang *et al.*, 2004). The antioxidant action of iodide has also been described in isolated rabbit eyes. Potassium iodide supplementation at normal dose had a significant antioxidative ability in the retina of the rat eye (Wu *et al.*, 2003). As an essential element, iodine is derived from the diet and humans require a daily intake of iodine (as iodide) of about one tenth of a milligram. It has earlier been reported that iodine deficiency is still a problem in Cameroon (Taga *et al.*, 2004, 2008, 2009). Similar results have also been observed in Bangladesh (Ara *et al.*, 2010). Lack of proper nutrition limits the body's ability to properly develop, protect and work at its capacity. However, earlier observations suggested that

Corresponding Author: Gabriel A. Agbor, Centre for Research on Medicinal Plants and Traditional Medicine, Institute for Medical Research and Medicinal Plants Studies, P.O. Box 6163, Yaoundé, Cameroon
Tel: 237 7722674

long and un-necessary use of iodized salt may produce sub clinical hyperthyroidism and thyrotoxicosis (Khan *et al.*, 2003).

Diabetes has been related to oxidative stress either as a causative factor or as its consequence (Qia *et al.*, 2008). Oxidative stress is known to cause tissue damage occurring in several diseases, including diabetes (El-Missiry *et al.*, 2004). Several research results have established Alloxan monohydrate at a single dose regiment of 150 mg kg⁻¹ body weight as a standard method for induction of experimental diabetes (Ananthan *et al.*, 2003; Ekor *et al.*, 2010; Manoharan *et al.*, 2007; Mohamed and Faddah, 2007). Alloxan oxidizes glutathione (GSH) to form hydrogen peroxide (Nakazaki *et al.*, 2000; Bromme *et al.*, 2001) which though a stable molecule but can be dangerous in the presence of a metal ion. Induction of diabetes mellitus by alloxan in animals has been associated with lipid peroxide levels (Nishikawa *et al.*, 2000) and a corresponding depression in the antioxidant defense mechanism characterized by decreased super oxide dismutase, catalase and glutathione (El-Missiry *et al.*, 2004).

The role of iodine supplementation on the antioxidant status of normal and alloxan monohydrate intoxicated rats was evaluated in this study.

MATERIALS AND METHODS

Reagents: All reagents were of analytical grade purchased from Sigma Chemical (St Louis, MO).

Experimental animals: This piece of research was conducted from July, 2009 to December 2009 in the Laboratory of Pharmacology in the Centre of Research on Medicinal Plants and Traditional Medicine, Institute of Medical Research and Medicinal Plants Studies. Twenty female Wistar albino rats raised in the Institute of Medical Research and Medicinal Plants Studies, Yaoundé Cameroon were used for this study. The experiment was carried out in accordance with the 1996 Guide for the Care and Use of Laboratory Animals and the Animal Welfare Act that pertains to research of this nature. The rats were divided into 4 groups of 5 rats each and housed in wire meshed cages. The groups A and C were not induced oxidative stress (kept normal), while oxidative stress was provoked in groups B and D by intraperitoneal injection of a solution of alloxan monohydrate (150 mg kg⁻¹ animal weight single dose) as earlier described (Ekor *et al.*, 2010). Animals with hyperglycemia (blood glucose = 200 mg dL⁻¹) were considered as stressed. The group A and B animals were fed the normal rodent diet while the group C and D animals were fed the normal rodent diet

supplemented with potassium iodide (KI 25 mg kg⁻¹ of food). The feeding lasted 6 weeks and the animals were sacrificed after an overnight fast by cervical dislocation and blood (4 mL) collected by cutting the jugular vein immediately. The plasma prepared from the blood was stored at -20°C until analyzed. The red blood cells were then washed using physiological saline (0.9% NaCl). The resulting red blood cells were then haemolysed by the addition of cold distilled water. The hemolysate was then stored at -20°C until analysed.

Biochemical analysis: Biochemical parameters analysed from the plasma were total protein (Bradford, 1976), protein thiols groups (SH) (Ellman, 1959), protein Carbonyl Groups (CO) (Reznick *et al.*, 1992) and bilirubin (Friedman and Young, 2004). Blood glucose was analysed from whole blood using Glucometer (one touch ultra2). Antioxidant markers such as Superoxide dismutase (SOD) (McCord and Fridovich, 1969) and catalase (Beers and Sizer, 1952) were analyzed from the hemolysate. FRAP (Ferric Reducing Antioxidant Power) was analysed by employing the method of Benzie and Strain (1996).

Statistical analysis: All data obtained were subjected to statistical analysis and are presented as Means±Standard deviation. Analysis of Variance on Ranks was employed in groups' comparison. All pair wise multiple comparisons to determine significant difference between groups were carried out using Kruskal-Wallis test. The SPSS software version 10.1 was employed in these analyses. The level of significance was set at p<0.05 for all tests.

RESULTS

Table 1 presents the effect of iodine supplementation on antioxidant status of normal and alloxan monohydrate intoxicated rats. Group A and C animals (normal rats) were not intoxicated with alloxan monohydrate. However, Group C animals received iodine supplementation. The iodine supplementation improved the antioxidant status of the Group C rats significantly (p<0.05) as seen in the bilirubin concentration (89.73±23.13 µmol L⁻¹), FRAP (420.71±17.57 µM) and catalase activity (24.0±11.56 UI mg⁻¹ protein) but did not alter the SOD activity significantly (p>0.05). Alloxan monohydrate intoxication induced a collapse in antioxidant status of group B animals by significantly (p<0.05) decreasing the FRAP (292.15±35.52 µM) and the activities of SOD (118.22±0.16 UI mg⁻¹ protein) and CAT (10.69±6.69 UI mg⁻¹ protein). However, supplementation with iodine maintained the antioxidant status as seen in group D animals with FRAP SOD (129.74±0.11 UI mg⁻¹

Table 1: Effect of Iodine supplementation on antioxidant status of normal and alloxan monohydrate intoxicated rats

Parameters	Group A (Normal rats)	Group B (ALX rats)	Group C (Normal+iodine)	Group D (ALX+iodine)
Bilirubin ($\mu\text{mol L}^{-1}$)	60.43±23.43	42.55±31.45 ^a	89.73±23.13 ^a	129.50±15.40 ^b
FRAP (μM)	397.78±38.41	292.15±35.52 ^a	420.71±17.57 ^a	335.65±17.75 ^b
Total protein ($\mu\text{g L}^{-1}$)	229.32±42.95	218.67±75.53	238.97±21.65	234.99±52.35
Catalase (UI mg^{-1} protein)	13.95±7.57	10.69±6.69 ^a	24.00±11.56 ^a	22.00±9.01 ^b
SOD (UI mg^{-1} protein)	139.80±0.08	118.22±0.16 ^a	134.65±0.93	129.74±0.11 ^b

Value represented as Mean±SD of 5 rats in each group. ^aindicated significant difference ($p<0.05$) compared to Group A (normal rats fed normal rodent diet), ^bindicated significant difference ($p<0.05$) compared to Group B (alloxan intoxicated rats fed normal rodent diet). ALX = Alloxan monohydrate

Table 2: Effect of Iodine supplementation on blood glucose and oxidative stress parameters of alloxan monohydrate intoxicated rats

Parameters	Group A (Normal rats)	Group B (ALX rats)	Group C (Normal+iodine)	Group D (ALX+iodine)
Blood glucose (mg dL^{-1})	96.00±4.69	227.04±69.0 ^a	100.50±5.27	206.10±34.58
TBARS ($\mu\text{mol L}^{-1}$)	1.95±1.06	3.06±0.44 ^a	1.68±0.25	1.82±0.88 ^b
SH ($\mu\text{mol mg}^{-1}$ protein)	45.44±8.79	206.25±20.22 ^a	38.79±19.6	180.88±33.92 ^b
CO ($\mu\text{mol mg}^{-1}$ protein)	180.64±73.52	252.50±82.5 ^a	186.02±32.59	203.03±17.41 ^b

Value represented as Mean±SD of 5 rats in each group. ^a indicated significant difference ($p<0.05$) compared to Group A (normal rats fed normal rodent diet), ^bIndicated significant difference ($p<0.05$) compared to Group B (alloxan intoxicated rats fed normal rodent diet). ALX: Alloxan monohydrate

protein) values turning toward normal. Table 2 presents the effect of iodine supplementation on blood glucose and oxidative stress parameters of normal and alloxan monohydrate intoxicated rats. Intoxication of experimental animals with alloxan monohydrate induced hyperglycemia (227.04 ± 69.0 mg dL^{-1}) which was not significantly reduced ($p>0.05$) following supplementation with iodine. Supplementation of normal rats with iodine did not alter the blood glucose concentration either. Carbonyl (CO), thiol (SH) and lipid peroxidation represented by Thiobarbituric Acid Reactive Substances (TBARS) were assessed as markers of oxidative injury. TBARS, SH and CO were significantly ($p<0.05$) increased in alloxan intoxication 3.06 ± 0.44 $\mu\text{mol L}^{-1}$, 206.25 ± 20.22 $\mu\text{mol mg}^{-1}$ protein and 252.5 ± 82.5 $\mu\text{mol mg}^{-1}$ protein, respectively. The increase in TBARS concentration in the plasma of rats in Group B is an indication of lipid peroxidation. These increases in TBARS, SH and CO were prevented in the iodine supplemented group 1.82 ± 0.88 $\mu\text{mol L}$, 180.88 ± 33.92 $\mu\text{mol mg}^{-1}$ protein and 203.03 ± 17.41 $\mu\text{mol mg}^{-1}$ protein, respectively.

DISCUSSION

Alloxan monohydrate treatment is an established method for induction of diabetes. The mechanism of action has been attributed to DNA fragmentation in pancreatic islets and cell damage free radicals (Takasu *et al.*, 1991). Diabetes has been related to oxidative stress either as it is a causative factor or because of its consequence (Qia *et al.*, 2008). Oxidative stress and tissue damage are common phenomena in several diseases especially those linked to toxic agents' exposure (El-Missiry *et al.*, 2004). Thus, controlling oxidative stress may either help prevent development of diabetes or control its consequence which may lead to further harmful effects. The present study evaluated the effect of iodine supplementation on the antioxidant status

of alloxan monohydrate intoxicated rats. A mark oxidative stress effect was induced by alloxan administration evidenced by the significant decline of endogenous antioxidant enzymes (SOD and CAT) activities, in the erythrocytes of experimental animals studied (Group B). Similar results have been obtained in isolated pancreatic islets (Fisher and Humburger, 1980), erythrocytes (Yadav *et al.*, 1996), liver and testis (Soto *et al.*, 1998; El-Missiry and El-Gendy, 2000); brain (El-Missiry *et al.*, 2004) and forebrain (Kosenko *et al.*, 1999) of experimental animals.

Catalase has the function of detoxification of H_2O_2 , an effective inhibitor of SOD (De Duve and Baudhuin, 1996). Catalase scavenges the active oxygen metabolites, and protects the cells from toxic oxidation (Marklund *et al.*, 1982). The decreased catalase activity in alloxan intoxicated animals (Group B) served as an indicator of increased reactive oxygen species generation in diabetic rats (Makni *et al.*, 2010; Farombi and Ige, 2007). Alloxan in a single injection destroys the insulin-producing islet β -cells in order to cause diabetes in experimental animals (Rerup, 1970). This is mediated by redox reaction which produces superoxide radicals in or near the β -cells (Deamer *et al.*, 1971; Heikkila *et al.*, 1976; Grankvist *et al.*, 1979). Thus, superoxide dismutase, enzyme removing superoxide anion radicals, may acts prophylactically against alloxan-induced diabetes (Grankvist *et al.*, 1981). In the present study alloxan induced a collapse in the SOD activity. Similar decline in SOD activity in alloxan treated animals has been reported by Makni *et al.* (2010) and Farombi and Ige (2007). Supplementation with iodine reinstated the antioxidant status of alloxan intoxicated animals (Group D) suggesting that iodine may have a superoxide anion radical scavenging activity.

Diabetes may induce increased systemic oxidative stress which is the underlying cause of dysregulation of adipocytokines and development of metabolic syndrome and pathogenesis of various diseases (Brownlee, 2001).

Oxidative stress impairs glucose uptake in muscle and adipose tissue in diabetic conditions (Maddux *et al.*, 2001; Rudich *et al.*, 1998) and decreases insulin secretion from pancreatic B cells (Matsuoka *et al.*, 1997). This results to a hyperglycemic environment which may impair radical scavenging activity and hence exposing proteins and lipids to peroxidation. Earlier studies have reported increases in lipid (TBARS) and protein oxidation (CO) products in diabetic patients (Telci *et al.*, 2000; Kesavulu *et al.*, 2001; Farombi and Ige, 2007). This is in conformity with the results obtained in the present study characterized by a significant rise in TBARS, CO and SH concentrations in alloxan intoxication. The increase in thiol and carbonyl concentrations may be due to the oxidation of the side chains of certain amino acids resulting to the liberation of the thiol and carbonyl groups in the plasma (Pincemail *et al.*, 1999). The evaluation of protein carbonyls as a useful marker of oxidative stress is based on its long half life in circulating blood (Trombetta *et al.*, 2006; Dalle-Donne *et al.*, 2003). This suggests that iodine supplementation to prevent the rise in oxidative stress.

CONCLUSION

Iodine functions as an antioxidant by increasing antioxidant enzyme activity in both normal and alloxan intoxicated rats there by reversing the effect of alloxan intoxication and fortifying the antioxidant defense mechanism. Reducing the accumulation of oxidative stress markers such as carbonyl and thiol functionality and lipid peroxidation (collapse of antioxidant function) is also an important property of iodine observed in this study. It will be important to evaluate the oxidative stress and antioxidant statutes in iodine deficient endemic areas so as to have an understanding of the risks involved and a better intervention program.

REFERENCES

- Ananthan, R., C. Baskar, V. Narmathabai, L. Pari and K.M. Ramkumar, 2003. Antidiabetic effect of *Gymnema montanum* leaves: Effect on lipid peroxidation induced oxidative stress in experimental diabetes. *J. Ethnopharmacol.*, 48: 551-556.
- Ara, G., A. Melse-Boonstra, S.K. Roy, N. Alam, S. Ahmed, U.H.F. Khatun and T. Ahmed, 2010. Sub-clinical iodine deficiency still prevalent in bangladeshi adolescent girls and pregnant women. *Asian J. Clin. Nutr.*, 2: 1-12.
- Beers, Jr. R.F. and I.W. Sizer, 1952. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.*, 195: 133-140.
- Benzie, I.F. and J.J. Strain, 1996. The Ferric Reducing Ability of plasma (FRAP) as a measure of antioxidant power: The FRAP assay. *Anal. Biochem.*, 239: 70-76.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- Bromme, H.J., R. Weinandy, D. Peschke and E. Peschke, 2001. Simultaneous quantitative determination of alloxan, GSH and GSSG by HPLc. Estimation of the frequency of redox cycling between alloxan and dialuric acid. *Horm. Metab. Res.*, 33: 106-109.
- Brownlee, M., 2001. Biochemistry and molecular cell biology of diabetic complications. *Nature*, 414: 813-820.
- Dalle-Donne, I., D. Giustarini, R. Colombo, R. Rossi and A. Milzani, 2003. Protein carbonylation in human diseases. *Trends Mol. Med.*, 9: 169-176.
- De Duve, C. and P. Baudhuin, 1996. Peroxisomes (microbodies and related particles). *Physiol. Rev.*, 46: 323-357.
- Deamer, D.W., R.E. Heikkila, R.V. Panganamala, G. Cohen, D.G. Cornwell, 1971. The alloxan-dialuric acid cycle and the generation of hydrogen peroxide. *Physiol. Chem. Phys.*, 3: 426-430.
- Ekor, M., A.O. Odewabi, A.G. Bakre, K.S. Oritogun, T.E. Ajayi and O.V. Sanwo, 2010. Comparative evaluation of the protective effect of the ethanolic and methanolic leaf extracts of *Sida acuta* against hyperglycaemia and alterations of biochemical and haematological indices in alloxan diabetic rats. *J. Pharmacol. Toxicol.*, 5: 1-12.
- El-Missiry, M.A. and A.M. El-Gendy, 2000. Amelioration of alloxan induced diabetes mellitus and oxidative stress in rats by oil of *Eruca sativa* seeds. *Ann. Nutr. Metab.*, 44: 97-100.
- El-Missiry, M.A., A.I. Othman and M.A. Amer, 2004. L-Arginine ameliorates oxidative stress in alloxan-induced experimental diabetes mellitus. *J. Applied Toxicol.*, 24: 93-97.
- Ellman, G.L., 1959. Tissue sulfhydryl groups. *Arch. Biochem. Biophys.*, 82: 70-77.
- Farombi, E.O. and O.O. Ige, 2007. Hypolipidemic and antioxidant effects of ethanolic extract from dried calyx of *Hibiscus sabdariffa* in alloxan-induced diabetic rats. *Fundam. Clin. Pharmacol.*, 21: 601-609.
- Fisher, J. and S.A. Humburger, 1980. Inhibition of alloxan action in isolated pancreatic islets by superoxide dismutase, catalase and a metal chelator. *Diabetes*, 29: 213-216.
- Friedman, R.B. and D.S. Young, 2004. Effects of Disease on Clinical, Laboratory Tests. 3rd Edn., AACC Press, Washington, DC.

- Grankvist, K., S. Marklund and I.B. Taeljedal, 1981. Superoxide dismutase is a prophylactic against alloxan diabetes. *Nature*, 294: 158-160.
- Grankvist, K., S. Marklund, J. Sehlin and I.B. Taeljedal, 1979. Superoxide dismutase, catalase and scavengers of hydroxyl radical protect against the toxic action of alloxan on pancreatic islet cells *in vitro*. *Biochem. J.*, 182: 17-25.
- Heikkila, R., B. Winston and G. Cohen, 1976. Alloxan induced-diabetes evidence for hydroxyl radical as a cytotoxic intermediate. *Biochem. Pharmacol.*, 25: 1085-1092.
- Kesavulu, M.M., B.K. Rao, R. Giri, J. Vijaya, G. Subramanyam and C. Apparao, 2001. Lipid peroxidation and antioxidant enzyme status in Type 2 diabetics with coronary heart disease. *Diabetes Res. Clin. Pract.*, 53: 33-39.
- Khan, A., S. Akhter, M.M. Siddiqui, G. Nawab and K.N. Khattak, 2003. Continuous use of iodized salt may cause thyrotoxicoses in plain areas of North West Frontier province (NWFP) of Pakistan. *J. Medical Sci.*, 3: 423-428.
- Kosenko, E.A., A. Kaminskii and I. Kaminskii, 1999. Activity of antioxidant enzymes in liver and brain is decreased in the early stage of diabetes and this decrease depends on the function of NMDA-receptors. *Vopr. Med. Khim.*, 45: 304-308.
- Maddux, B.A., W. See, J.C. Lawrence, Jr., A.L. Goldfine, I.D. Goldfine and J.L. Evans, 2001. Protection against oxidative stress-induced insulin resistance in rat L6 muscle cells by micromolar concentrations of α -lipoic acid. *Diabetes*, 50: 404-410.
- Maier, J., H. van Steeg, C. van Oostrom, R. Paschke, R.E. Weiss and K. Krohn, 2007. Iodine deficiency activates antioxidant genes and causes DNA damage in the thyroid gland of rats and mice. *Biochim. Biophys. Acta, Mol. Cell. Res.*, 1773: 990-999.
- Makni, M., H. Fetoui, N.K. Gargouri, E.M. Garoui and N. Zeghal, 2010. Antidiabetic effect of flax and pumpkin seed mixture powder: Effect on hyperlipidemia and antioxidant status in alloxan diabetic rats. *J. Diabetes Complicat.*
- Manoharan, S., S. Silvan, K. Vasudevan and S. Balakrishnan, 2007. Antihyperglycemic and antilipidperoxidative effects of *Piper longum* (Linn.) dried fruits in alloxan induced diabetic rat. *J. Boil. Sci.*, 7: 161-168.
- Marklund, S.L., E. Holme and L. Hellner, 1982. Superoxide dismutase in extracellular fluids. *Clin. Chim. Acta*, 126: 41-51.
- Matsuoka, T., Y. Kajimoto, H. Watada, H. Kaneto and M. Kishimoto *et al.*, 1997. Glycation-dependent, reactive oxygen species-mediated suppression of the insulin gene promoter activity in HIT cells. *J. Clin. Invest.*, 99: 144-150.
- McCord, J.M. and I. Fridovich, 1969. Superoxide dismutase: An enzymic function for erythrocyte (Hemocytin). *J. Biol. Chem.*, 244: 6049-6055.
- Mohamed, A.M. and L.M. Faddah, 2007. Therapeutic effects of sildenafil citrate (Viagra) and/or vitamin E on some brain disorders of alloxan induced diabetes mellitus rats. *J. Medical Sci.*, 7: 932-941.
- Nakazaki, M., M. Kakei, K. Yaekura, N. Koriyama and S. Morimitsu *et al.*, 2000. Diverse effects of hydrogen peroxide on cytosolic Ca^{+2} homeostasis in rat pancreatic beta-cells. *Cell Struct. Funct.*, 25: 187-193.
- Nishikawa, T., D. Edelstein, X.L. Du, S. Yamagishi and T. Matsumura *et al.*, 2000. Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature*, 404: 787-790.
- Pincemail, J., M. Meurisse, R. Limet and J.O. Defraigne, 1999. Evaluating an individual's oxidative stress: A reality for doctors. *Vaisseaux, Coeur, Poumon*, 4: 148-154.
- Qia, X.Y., W.J. Chen, L.Q. Zhang and B.J. Xie, 2008. Mogrosides extract from *Siraitia grosvenori* scavenges free radicals *in vitro* and lowers oxidative stress, serum glucose and lipid levels in alloxan-induced diabetic mice. *Nutr. Res.*, 28: 278-284.
- Rerup, C.C., 1970. Drugs producing diabetes through damage of the insulin secreting cells. *Pharmacol. Rev.*, 22: 485-518.
- Reznick, A.Z., C.E. Cross, M.L. Hu, Y.J. Suzuki and S. Khwaja *et al.*, 1992. Modification of plasma proteins by cigarette smoke as measured by protein carbonyl formation. *Biochem. J.*, 286: 607-611.
- Rudich, A., A. Tirosh, R. Potashnik, R. Hemi, H. Kanety and N. Bashan, 1998. Prolonged oxidative stress impairs insulin-induced GLUT4 translocation in 3T3-L1 adipocytes. *Diabetes*, 47: 1562-1569.
- Soto, C.P., B.L. Perez, L.P. Favari and J.L. Reyes, 1998. Prevention of alloxan induced diabetes mellitus in the rat by silymarin. *Comp. Biochem. Physiol. Pharmacol. Toxicol. Endocrinol.*, 119: 125-129.
- Taga, I., M.L. Sameza, A.V. Kayo and J. Ngogang, 2004. Iodine levels in food and soil in different regions in Cameroon. *Sante* 14: 11-15.
- Taga, I., V.A.S. Oumbe, R. Johns, M.A. Zaidi, J.N. Yonkeu and I. Altosaar, 2008. Youth of West-Cameroon are at high risk of developing IDD due to low dietary iodine and high dietary thiocyanate. *Afr. Health Sci.*, 8: 180-185.

- Taga, I., G.A. Agbor, S. Djountsop, Z.A. Moshin, V.A. Oumbe, I. Altosaar and J.Y. Ngogang, 2009. Iodine deficiency in children 7-19 years old in Eastern Province of Cameroon. *Trop. Med. Int. Health.*, 14: 11-15.
- Takasu, N., I. Komiyada, T. Asawa, Y. Nagasawa and T. Yamada, 1991. Streptozocin- and alloxan-induced H₂O₂ generation and DNA fragmentation in pancreatic islets. H₂O₂ as mediator for DNA fragmentation. *Diabetes*, 40: 1141-1145.
- Telci, A., U. Cakatay, R. Kayali, C. Erdogan, Y. Orhan, A. Sivas and T. Akcay, 2000. Oxidative protein damage in plasma of type 2 diabetic patients. *Horm. Metab. Res.*, 32: 40-43.
- Trombetta, D., S. Gangemi, A. Saija, P.L. Minciullo and F. Cimino, 2006. Increased protein carbonyl groups in the serum of patients affected by thalassemia major. *Ann. Hematol.*, 85: 520-522.
- Venturi, S. and M. Venturi, 1999. Iodide, thyroid and stomach carcinogenesis: Evolutionary story of a primitive antioxidant?. *Eur. J. Endocrinol.*, 140: 371-372.
- Wang, J., Y. Ge, H. Ning and S. Wang, 2004. Effects of high Fluoride and low iodine on oxidative stress and antioxidant defense of the brain in offspring rats. *Fluoride*, 37: 264-270.
- Wu, Y.X., L.J. Li, G.Y. Chen, W.D. Zhao and M.C. Qiu, 2003. Effects of supplementation of different kinds of iodine on the antioxidative ability of retina in iodine deficient rats. *Zhonghua Yan Ke Za Zhi*, 39: 495-498.
- Yadav, P., S. Sarkar, N. Rathor, S. John, M. Kale and D. Bhatnagar, 1996. Ascorbic acid counteracts the prooxidant effect of alloxan in erythrocytes *in vitro*. *Curr. Sci.*, 71: 404-406.