



Review Article

Role of Phytoecdysteroids in Insect Pest Management: A Review

Mukesh Kumar Chaubey

Department of Zoology, Mahatma Gandhi Post Graduate College Gorakhpur, 273009 Uttar Pradesh, India

Abstract

Phytoecdysteroids are analogues of insect moulting steroid hormone, ecdysteroid. These are found in different plant groups and provide protection by deterring insects. Phytoecdysteroids have been synthesized in at least 27 families of *Pteridophyta*, 10 families of *Gymnospermae* and 74 families of *Angiospermae*. Chemically, phytoecdysteroids are triterpenoids, the group of compounds that includes triterpene saponins, phytosterols and phytoecdysteroids. These are polar steroids with sugar-like solubility properties. Over 300 phytoecdysteroids analogues have been identified so far and it has been speculated that there are over 1,000 possible structures in nature. These are synthesized from mevalonic acid and cholesterol. Phytoecdysteroids occur in relatively high concentration in many plants and comprise 0.001-3% of the dry weight. These have been isolated from all parts of plants in much higher amounts than those present in insects. Thus, plants are far better sources of ecdysteroids than insects. Different plant parts contain different amounts of ecdysteroids and that ecdysteroid concentration varies with season and geographical distribution of the plant. Ecdysteroids control insect development at all stages of the life cycle. Disruption of normal ecdysteroid level severely impairs insect development. Phytoecdysteroids mimic the insect ecdysteroid by binding to its receptors and eliciting a cascade of effects in insects. Phytoecdysteroids provide protection to plants by altering the normal levels of ecdysteroid hormone in adults and larvae in insects. Therefore, phytoecdysteroids can be an excellent replacement of synthetic insecticides in insect pest management programme.

Key words: Ecdysone, phytoecdysteroides, triterpenoids, moulting hormone, genetic manipulation

Received:

Accepted:

Published:

Citation: Mukesh Kumar Chaubey, 2018. Role of phytoecdysteroids in insect pest management. J. Agron., CC: CC-CC.

Corresponding Author: Mukesh Kumar Chaubey, Department of Zoology, Mahatma Gandhi Post Graduate College Gorakhpur, 273009 Uttar Pradesh, India
Tel: +91-9839427296

Copyright: © 2018 Mukesh Kumar Chaubey. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The author has declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

With the beginning of agricultural practices and storage of food grains, insects started damaging standing crops and grains in storage both qualitatively and quantitatively. Several synthetic pesticides have been developed and used to protect crops and grains from insect infestation. The continuous and extensive application of synthetic pesticides has developed the risk of ozone depletion, neurotoxicity, carcinogenicity, teratogenicity and mutagenicity in non-target animal species and cross- and multi-resistance in insects¹⁻⁴. These adverse effects have increased public awareness regarding human safety and possible environmental damage diverting attention towards other alternatives especially the use of plant products in insect pest management.

Plant botanicals include plant extracts. These are secondary plant metabolites synthesized by the plant for protective purposes. Some of these compounds are toxic to insects. These plant derived chemicals are called botanical pesticides. These target a broad range of insect pests. Plant insecticides act as toxicants⁵⁻⁸, as repellents⁹⁻¹¹, as antifeedants¹²⁻¹⁵, as oviposition deterrents^{9-11,13} and developmental inhibitory activities^{9-11,14-17}. Botanical insecticides do not affect the natural enemies of the pests adversely. Botanical pesticides are good alternatives to chemical pesticides as these are cheap, eco-friendly, cost-effective, target-specific and biodegradable. This study work synergistically and it is difficult for insect pests to develop resistance.

Ecdysteroid, insect moulting hormone, was first isolated from *Bombyx mori*¹⁸. This hormone governs all the aspects of transformation from larva to pupa to adult. A number of ecdysteroid analogues called phytoecdysteroids have been reported from plants. Phytoecdysteroids are a class of chemicals that plants synthesize for defense against phytophagous insects. Insects that ingest phytoecdysteroids and have not adapted to these defense molecules are subject to serious adverse effects, including weight loss, molting disruption and/or mortality¹⁹⁻²¹. Ecdysteroids control insect development at all stages of the life cycle and disruption of normal ecdysteroid level or its action can be expected to impair insect development severely by mimicking the function of insect ecdysteroid²². After the discovery of 20-Hydroxyecdysone, active form of ecdysone in the conifer, *Podocarpus nakai*²³, ecdysteroids have been detected in 27 families of the *Pteridophyta*, 10 families of *Gymnospermae* and 74 families of *Angiospermae*²⁴. Since phytoecdysteroids are found in primitive plants such as ferns, it was assumed that these secondary compounds appeared very early in plant

evolution. In the angiosperms, 74 families possess ecdysteroid containing species. A majority of these families are within eleven orders: *Liliales*, *Ranunculales*, *Urticales*, *Malvales*, *Violales*, *Capparales*, *Rosales*, *Sapindales*, *Polemoniales* and *Scrophulariales*.

Phytoecdysteroids occur in relatively high concentration in many plants and comprise 0.001-3% of the dry weight and have been isolated from all parts of plants in much higher amounts than those present in arthropods^{19,20}. They are far better source of ecdysteroids than arthropods²⁵. It is important to note that different plant parts contain different amounts of ecdysteroids and that ecdysteroid concentration varies with season and geographical distribution of the plant²⁶. Presence of the moulting hormone analogues in plants is suggestive of their role as allelochemicals, protecting them against insect predation by acting as antifeedants and/or disrupting the endogenous endocrine level of insects²⁷. These phytoecdysteroids provide little protection against specialized herbivores, which have evolved efficient detoxification mechanisms but represent a consistent barrier against non-adapted herbivores. Low concentrations (2-25 ppm) of phytoecdysteroids deter some insects, while other insects are resistant to very high concentrations (400-1000 ppm).

The role of phytoecdysteroids in plants is not clearly understood, as there are conflicting data supporting ecdysteroids as plant hormones (i.e., signaling molecules produced in plants) which travel to targeted locations, regulating specific cellular responses. Although initial reports show that ecdysteroids do not elicit any of the classical responses of known plant hormones, there are also data supporting ecdysteroids being physiologically active in plants²⁸. Phytoecdysteroids may play a role in plant defense. Production of 20-Hydroxyecdysteroids in spinach is elicited by both mechanical wounding and insect feeding. Phytoecdysteroids protect spinach from plant-parasitic nematodes and may confer a mechanism for nematode resistance²⁹. Although not conclusive, ecdysteroid's role in plants may be similar to the defensive role of lignins, whose production increases in response to pathogens. Application of these phytoecdysteroids on insects has been reported to result in marked growth and developmental disruption³⁰⁻³⁴. In this review, sources, structures and applications of phytoecdysteroids have been discussed. Phytoecdysteroids provide protection to plants by altering the normal levels of ecdysteroid hormone in adults and larvae of insects. Therefore, phytoecdysteroids can be applied against insects causing economic injury and can be proved an excellent replacement of synthetic insecticides in insect pest management programme.

CHEMISTRY OF PHYTOECDYSTEROIDS

Chemically, phytoecdysteroids are triterpenoids, a group of compounds that includes triterpene saponins, phytosterols and phytoecdysteroids. Ecdysteroids are polar steroids with sugar-like solubility properties³³. Animals and humans also synthesize some of the same products from the mevalonate pathway, such as ubiquinone and sterols. They can not, however, synthesize ecdysteroids. Phytoecdysteroids universally lack the polyhydroxylated side chain characteristic of ecdysteroids and are, therefore, more lipophilic³³. Unlike invertebrates, which are unable to synthesize ecdysteroids and must consume dietary phytosterols that are then converted into ecdysteroids, plants can synthesize ecdysteroids from mevalonic acid and cholesterol³⁵. Plants synthesize phytoecdysteroids from mevalonic acid in the mevalonate pathway in cell using acetyl-CoA as a precursor.

Most phytoecdysteroids possess a (cyclopentano-perhydrophenanthrene carbon skeleton) cholest-7-en-6-one carbon skeleton (C27), derived biosynthetically from cholesterol and phytosterols, often with a hydroxyl group in the 14-position³⁶. The carbon number can vary between C19-C29³⁵. The anellations of the rings are characteristic: The C/D ring junctions are generally trans, while the A/B ring junction is generally cis (5 β -H) and only rarely trans (5 α -H). Essential structural elements of phytoecdysteroids are a 7-en-6-one chromophore in ring B and a β -side chain at C-17. Phytoecdysteroids are highly hydroxylated with 2-8 hydroxyl groups. The common hydroxylated sites are 2 β , 3 β , 14 α , 20R, 22R and 25th position which together give rise to the highly biologically active ponasterone A (25-Deoxy-20-hydroxyecdysone)^{19,20}. The structural variations among different phytoecdysteroids include number, position and orientation of hydroxyl groups. Ecdysteroids hydroxylated at C-9, C-12, C-17, C-19, C-21, C-24, C-28 and C-29 occur only in plants. Further possibilities are additional/alternative double bond at one of locations of 4-5, 8-9, 9-11, 12-13, 14-15, 24-25, 25-26 or 24-28. Second oxogroup can occur in one of the positions of C-2, C-3, C-12, C-17, C-20 or C-22 (Fig. 1). Among the natural steroids the cardenolides, bufadienolides, sterines and bile acids have A/B cis anellation. Further common characteristic is the C-11-hydroxylation (in addition to C-1-, C-3-, C-5- and C-14-hydroxylations), which is usually frequent at ecdysteroids as well as for cardenolides and bufadienolides.

Although the most common ecdysteroid found in plants is 20-Hydroxyecdysone, over 300 ecdysteroid analogues have been identified so far in plant kingdom and it has been speculated that there are over 1,000 possible structures in




Fig. 1: General structure of ecdysteroid (heavy line makes the position of hydroxyl group in β -orientation, solid arrows show position of hydroxyl group in α or β -orientation and dotted arrows show position of hydroxyl group in α -orientation. Numbers show carbon number in the ecdysteroid skeleton


nature^{19,20} (Fig. 2). The various analogues differ in the number and site of hydroxylations, as well as the length and structure of the carbon side chain. Glycosylated and acetylated ecdysteroids have been described both in nature and in the laboratory³⁷. Many more plants have the ability to 'turn on' the production of phytoecdysteroids when under stress, animal attack or other conditions¹⁹.

The term phytoecdysteroid can also apply to ecdysteroids found in fungi, even though fungi are not plants. Fungi that produce phytoecdysteroids include *Achyranthes bidentata*³⁸, *Tinospora cordifolia*³⁹, *Pfaffia paniculata*⁴⁰, *Leuzea carthamoides*⁴¹, *Rhaponticum uniflorum*⁴², *Serratula coronata*⁴³, *Cordyceps* and *Asparagus*²⁰.


During discovery for anti-cancer compounds in plant extracts, Nakanishi *et al.*²³ isolated ponasterone A, a C27 polyhydroxy steroid closely related to the insect moulting hormone, from the plant *Podocarpus nakaii*. Soon after the isolation of ponasterone A, 20-Hydroxyecdysone, the major ecdysteroid of arthropods was reported in a conifer, *Podocarpus elatus*⁴⁴, *Polypodium vulgare* (fern)⁴⁵ and the angiosperm, *Achyranthes fauriei*⁴⁶. Later, both ecdysone and 20-Hydroxyecdysone were isolated from the ferns, *Pteridium aquilinum*⁴⁷ and *Polypodium vulgare*⁴⁸.

PLANT FAMILIES PRODUCING ECDYSTEROIDS


Approximately 6% of all plant species synthesize phytoecdysteroids¹⁹⁻²⁰ (Table 1). Among plant species, highest concentration of phytoecdysteroid is found in spinach, *Spinacia oleracea* (Chenopodiaceae) containing 50 $\mu\text{g g}^{-1}$




Ecdysone




20-Hydroxyecdysone




Ponasterone A




Poststerone




Poststerone C




Pterosterone




Ajugasterone C




Polypodine B




Muristerone A




2-Deoxy-20-hydroxyecdysterone




2-Deoxyecdysone




2-Deoxy polypodine



22-Deoxy-20-hydroxyecdysone



Cyasterone



24(28)-Dehydro makisterone A

Fig. 2: Continue




Fig. 2: Diversity of phytoecdysteroid structures

fresh weight⁴⁹. Phytoecdysteroids obtained from plants are ajugasterone in *Ajuga* (Lamiaceae) and *Vitex* (Verbenaceae); leuzeasterone in *Leuzea carthamoides* (Asteraceae), 2-Deoxy-20-hydroxyecdysone 3-Glucoside and 3-Epi-2-deoxy-20-hydroxyecdysone in *Tinospora cordifolia* (Menispermaceae)³⁹, ecdysterone and inokosterone in *Achyranthes bidentata* (Amaranthaceae), polypodine B in *Leuzea carthamoides* (Asteraceae)⁴¹ and ecdysterone, ajugasterone C, ajugasterone C-20, 22-monoacetonide in *Rhaponticum uniflorum* (Asteraceae)⁴². To date, 300 different phytoecdysteroids have been identified⁵⁰. Phytoecdysteroids differ in the number of

C-atoms present (24C-29C), the number, position and location of hydroxyl and keto groups on the steroid skeleton. The most commonly occurring phytoecdysteroid is 20-Hydroxyecdysone, followed by polypodine B. Typically, a phytoecdysteroid-containing species will contain 1-3 major ecdysteroids, which together make up 95% of the total ecdysteroid, together with plenty of minor ecdysteroids, forming a diverse cocktail of ecdysteroid structural analogues. Although most of the plant families have at least some species accumulate ecdysteroids, less than 2% of the world's flora has been investigated for the presence of ecdysteroids^{19,51}.

Table 1: Plant families and species containing phytoecdysteroids

Plant species	Family
<i>Achyranthes aspera</i>	Amaranthaceae ³⁸
<i>Achyranthes bidentata</i>	Amaranthaceae ³⁸
<i>Ajuga turkestanica</i>	Lamiaceae ⁷¹
<i>Cassia tora</i>	Caesalpinaceae ⁷²
<i>Cyanotis somaliensis</i>	Commelinaceae ⁵⁰
<i>Cyathula achyranthoides</i>	Amaranthaceae ⁵⁰
<i>Cyathula officinalis</i>	Amaranthaceae ⁵⁰
<i>Dianthus deltooides</i>	Caryophyllaceae ⁷³
<i>Leuzea carthamoides</i>	Asteraceae ⁷³
<i>Melandrium firmum</i>	Caryophyllaceae ⁷⁴
<i>Melandrium sachalinense</i>	Caryophyllaceae ⁷⁴
<i>Pfaffia iresinoides</i>	Amaranthaceae ⁷⁵
<i>Polypodium aurea</i>	Polypodiaceae ⁷⁵
<i>Polypodium glycyrrhiza</i>	Polypodiaceae ⁵⁰
<i>Polypodium vulgare</i>	Polypodiaceae ⁵⁰
<i>Rhaponticum uniflorum</i>	Asteraceae ⁴²
<i>Saponaria bellidifolia</i>	Caryophyllaceae ⁷³
<i>Serratula coronata</i>	Asteraceae ⁷⁶
<i>Spinacia oleracea</i>	Chenopodiaceae ⁴⁹
<i>Tinospora cordifolia</i>	Menispermaceae ³⁹

MECHANISM OF ACTION OF ECDYSTEROIDS

Hydroxyecdysone is a naturally occurring ecdysteroid which controls the ecdysis and metamorphosis in insects. Ecdysteroid is derived from enzymatic modification of cholesterol by p450 enzymes. The primary sources of 20-Hydroxyecdysone in larvae are the prothoracic gland, ring gland, gut and fat bodies in larvae. These tissues convert dietary cholesterol into 20-Hydroxyecdysone⁵². In the adult female, ovary is a source of 20-Hydroxyecdysone production⁵³. In adult males, fat body is the source of hydroxyecdysone. Its analogues called phytoecdysteroids are produced by various plants where these disrupt the development and reproduction of insect pests. This hormone is hydrophobic in nature, thus, it traverses lipid membranes and permeates the tissues of an organism. In insects, 20-Hydroxyecdysone acts through the ecdysone receptor. The ecdysone receptor is a nuclear receptor (a ligand-activated transcription factor), which controls development and contributes to other processes such as reproduction^{54,55}. The ecdysone receptor is a non-covalent heterodimer of two proteins viz., the EcR protein and ultraspiracle protein (USP). The ecdysteroid-binding pocket is located in the EcR subunit, but EcR must be dimerised with a USP for high-affinity ligand binding. The binding of ecdysone to receptor leads to the activation of ecdysone responsive genes and many other genes causing puffing of polytene chromosomes at over a hundred sites. Sites of puffing represent the position of genes to be expressed and thus, position of transcription. The DNA-binding domains of EcR and USP recognise specific short sequences in DNA and mediate the binding of the

heterodimer to ecdysone response elements (ECREs) in the promoters of ecdysone-responsive genes. The first gene to be activated are called early response genes. This is a code for proteins that are themselves gene regulatory factors either alone or in combination with hormones. They activate late-response genes that code for the proteins that actually cause structural changes, cell differentiation or apoptosis (programmed cell death). Ultimately, the activation cascade causes physiological changes that result in ecdysis (moulting)⁵⁶. The ecdysone receptor also binds to and is activated by phytoecdysteroids. Thus, phytoecdysteroids can mimic 20-Hydroxyecdysone of insects, bind insect ecdysone receptors and can elicit the same responses. These phytoecdysteroids induced responses at inappropriate time and stage causes feeding deterrence, abnormal development, altered pattern of ecdysis and finally death.

PHYTOECDYSTEROIDS AS INSECT GROWTH REGULATORS

Ecdysteroides inhibit the feeding in *Pieris brassicae* and *Mamestra brassicae* larvae when given at 200 mg kg⁻¹ fresh weight in sucrose solution⁵⁷ and inhibit drinking in *Dysdercus koenigii*, *D. fulvoviger* and *Spiloslethus pandrus* adult at 100 mg kg⁻¹ concentration⁵⁸. Jones and Firm⁵⁹ reported that ecdysone and 20-Hydroxyecdysone deter feeding in *P. brassicae* when incorporated above 5 mg kg⁻¹ diet. *Chilo partellus*, *Phyllobius pyri* and *P. argentatus* were deterred from feeding at concentration of 20-Hydroxyecdysone above 50-70 mg kg⁻¹, whereas, *Schistocerca gregaria*, *Hyponomeuta euonymella* and *Spodoptera littoralis* were not deterred at this concentration⁵⁹. Hydroxyecdysone and polypodine B regulate the length of feeding periods in larval instars and silk production when added to the diet of *B. mori*⁶⁰. Methanolic extract of *Ajuga remota* leaves and roots containing cyasterone and ecdysterone disrupt the moulting cycle in *B. mori*, *Peclinophora gossypiella* and *Spodoptera frugiperda*⁶¹.

Ingested 20-Hydroxyecdysone causes death without moulting, death following completion of promoted moulting or death during moulting in larvae of *B. mori*. Comparison of hydroxyecdysone with other phytoecdysteroids in *B. mori* and *P. gossypiella* shows ponasterone A as more potent effect⁶². Effective ecdysis inhibition has been observed in *P. gossypiella* with ponasterone A⁶³. Phytoecdysteroids like ecdysterone, polypodine B and ponasterone A when applied in 25-250 ppm in the diet induce ecdysial failure associated with the appearance of larvae having two head capsules and developmental anomalies during metamorphosis in larvae of *Acrolepiopsis assectella*³⁰. Slama *et al.*⁶⁴ have reported

hormonal activity of ecdysteroids in the larvae of *Galleria mellonella*, *Dermestes* sp. and *Sarcophaga bullata* and found cyasterone as the most effective lepidopteran ecdysteroid and turkesterone as the dipteran and coleopteran specific ecdysteroid⁶⁴. Kubo⁶⁵ conducted artificial diet feeding assay with the green bug, *Schizaphis graminum* and showed ajugasterone C to have 10 fold more potency in feeding deterrancy than 20-Hydroxyecdysone and 30 fold more potency than cyasterone⁶⁵.

Ecdysone in artificial diets of *B. mori* induces ultranumerary larval ecdyses³². Tanaka⁶⁶ has reported altered sensitivity of epidermis to 20-Hydroxyecdysone at 300 ppm of ecdysone. Only a very limited number of insect species have been examined for their susceptibility to dietary ecdysteroids. Most of the species tested belong to order *Lepidoptera* which includes phytophagous pests. Among the tested lepidopteran species some are highly susceptible to dietary 20-Hydroxyecdysone such as *A. assectella*, *Aglaisurticae*, *B. mori*, *C. partellus*, *Inachisio* and *P. gossypiella*, some are semi-tolerant like *Cynthia cardui* and *Tyria jacobaeae* while others such as *A. convolvuli*, *Helicoverpa* spp., *M. brassicae*, *Manduca sexta* and *S. littoralis* are highly resistant⁶⁷. The specific effects of ecdysteroids and their low toxicity formed the basis of application of phytoecdysteroids as safe insecticides⁶⁸. However, the phytoecdysteroids have limited application in the control of insect pest because of their environmental instability⁶⁹.

Ecdysteroids are compounds with low mammalian toxicity. The LD₅₀ values in mice are 6.4 and >9 g kg⁻¹, using intraperitoneal and oral administration of 20-Hydroxyecdysone, respectively. The 0.1 g kg⁻¹ dose of intravenous administration of 20-Hydroxyecdysone to rabbits has not been followed with any toxic reaction and sub-acute treatment of rats with 2 g kg⁻¹ day⁻¹ treatment neither gave toxic symptoms⁷⁰.

CONCLUSION

Phytoecdysteroids are toxic compounds that defend plants against phytophagous insects. These can be used as insecticidal tool in pest management programme, as these are eco-friendly, economic, target-specific and biodegradable.

SIGNIFICANCE STATEMENTS

Phytoecdysteroids are plant derived steroid hormones analogues to insect moulting hormone. These deter predation by non-adapted insect predators. This property of phytoecdysteroids can be exploited to develop new strategies for insect pest control. These plant derived

ecdysteroids can be used as lead molecules for developing insect controlling agents. New varieties of plants of economic interest can be developed with altered ecdysteroid levels/profiles through plant breeding or genetic manipulation.

REFERENCES

1. WMO., 1991. Scientific assessment of ozone depletion. WMO Report No. 25, World Meteorological Organization of the United Nations, Geneva, Switzerland.
2. Lu, F.C., 1995. A review of the acceptable daily intakes of pesticides assessed by WHO. Regul. Toxicol. Pharmacol., 21: 352-364.
3. UNEP., 2000. The montreal protocol on substances that deplete the ozone layer. United Nations Environment Programme, Ozone Secretariat, Nairobi, Kenya.
4. Beckel, H., I. Lorini and S.M.N. Lazzari, 2002. [Resistance of *Oryzaephilus surinamensis* (L.) (Coleoptera: Silvanidae) to insecticides pyrethroids and organophosphates used in stored wheat]. In: [Summaries and Minutes of III Technical Seminar of Trigo/XVII Reuniao of the Commission I South-Center Brazilian of Inquiry of Wheat], pp: 44, (In Portuguese).
5. Sadek, M.M., 2003. Antifeedant and toxic activity of *Adhatoda vasica* leaf extract against *Spodoptera littoralis* (Lep., Noctuidae). J. Applied Entomol., 127: 396-404.
6. Rodriguez-Saona, C.R. and J.T. Trumble, 1999. Effect of avocadofurans on larval survival, growth and food preference of the generalist herbivore, *Spodoptera exigua*. Entomol. Exp. Applic., 90: 131-140.
7. Ulrichs, C., I. Mewis, S. Adhikary, A. Bhattacharyya and A. Goswami, 2008. Antifeedant activity and toxicity of leaf extracts from *Porteresia coarctata* Takeoka and their effects on the physiology of *Spodoptera litura* (F.). J. Pest Sci., 81: 79-84.
8. Isman, M.B., 2002. Insect antifeedants. Pest. Outlook, 13: 152-157.
9. Chaubey, M.K., 2011. Insecticidal properties of *Zingiber officinale* and *Piper cubeba* essential oils against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). J. Biol. Active Plant Prod. Nat., 1: 306-313.
10. Chaubey, M.K., 2011. Fumigant toxicity of essential oils against rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionidae). J. Biol. Sci., 11: 411-416.
11. Chaubey, M.K., 2008. Fumigant toxicity of essential oils from some common spices against pulse beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). J. Oleo Sci., 57: 171-179.
12. Hummelbrunner, L.A. and M.B. Isman, 2001. Acute, sublethal, antifeedant and synergistic effects of monoterpenoid essential oil compounds on the tobacco cutworm, *Spodoptera litura* (Lep., Noctuidae). J. Agric. Food Chem., 49: 715-720.

13. Upadhyay, R.K., L. Rohtagi, M.K. Chaubey and S.C. Jain, 2006. Ovipositional responses of the pulse beetle, *Bruchus chinensis* (Coleoptera: Bruchidae) to extracts and compounds of *Capparis decidua*. J. Agric. Food Chem., 54: 9747-9751.
14. Malarvannan, S., R. Giridharan, S. Sekar, V.R. Prabhavathi and S. Nair, 2008. Bioefficacy of crude and fractions of *Argemone mexicana* against Tobacco caterpillar, *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae). J. Biopesticides, 1: 55-62.
15. Chaubey, M.K., 2007. Insecticidal activity of *Trachyspermum ammi*(Umbelliferae), *Anethum graveolens*(Umbelliferae) and *Nigella sativa*(Ranunculaceae) against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). Afr. J. Agric. Res., 2: 596-600.
16. Ray, D.P., D. Dutta, S. Srivastava, B. Kumar and S. Saha, 2012. Insect growth regulatory activity of *Thevetia nerifolia* Juss. against *Spodoptera litura* (Fab.). J. Applied Bot. Food Qual., 85: 212-215.
17. Chaubey, M.K., 2012. Fumigant toxicity of essential oils and pure compounds against *Sitophilus oryzae* L. (Coleoptera: Curculionidae). Biol. Agric. Hortic., 28: 111-119.
18. Butenandt, A. and P. Karlson, 1954. Über die Isolierung eines metamorphose-hormons der Insekten in kristallisierter Form. Zeitschrift für Naturforsch., 9b: 389-391.
19. Dinan, L., T. Savchenko and P. Whiting, 2001. On the distribution of phytoecdysteroids in plants. Cell. Mol. Life Sci., 58: 1121-1132.
20. Dinan, L., T. Savchenko and P. Whiting, 2001. Phytoecdysteroids in the genus *Asparagus* (Asparagaceae). Phytochemistry, 56: 569-576.
21. Schmelz, E.A., R.J. Grebenok, T.E. Ohnmeiss and W.S. Bowers, 2002. Interactions between *Spinacia oleracea* and *Bradysia impatiens*. A role for phytoecdysteroids. Arch. Insect Biochem. Physiol., 51: 204-221.
22. Dinan, L., P. Whiting, D. Alfonso and I. Kapetanidis, 1996. Certain withanolides from *Lochroma gesnerioides* antagonize ecdysteroid action in a *Drosophila melanogaster* cell line. Entomol. Exp. Applic., 80: 415-420.
23. Nakanishi, K., M. Koreeda, S. Sasaki, M.L. Chang and H.Y. Hsu, 1966. Insect hormones. The structure of ponasterone A, insect-moulting hormone from the leaves of *Podocarpus nakaii* Hay. Chem. Commun., 24: 915-917.
24. Camps, F., 1991. Plant Ecdysteroids and their Interaction with Insects. In: Ecological Chemistry and Biochemistry of Plant Terpenoids, Harborne, I.B. and E.A. Tomas-Barberan (Eds.). Chapter 14, Clarendon Press, Oxford, UK., ISBN-13: 9780198577393, pp: 331-376.
25. Hom, D.H.S. and R. Bergamasco, 1985. Chemistry of Ecdysteroids. In: Comprehensive Insect Physiology, Biochemistry and Pharmacology, Volume 7: Endocrinology I, Kerkut, G.A. and L.I. Gilbert (Eds.). Pergamon Press, London, UK., pp: 285-248.
26. Hikino, H., T. Okuyama, H. Jin and T. Takemoto, 1973. Screening of Japanese ferns for phytoecdysones. I. Chem. Pharmaceut. Bull., 21: 2292-2302.
27. Blackford, M., B. Clarke and L. Dinan, 1996. Tolerance of the Egyptian cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae) to ingested phytoecdysteroids. J. Insect Physiol., 42: 931-936.
28. Machackova, I., M. Vagner and K. Slama, 1995. Comparison between the effects of 20-hydroxyecdysone and phytohormones on growth and development in plants. Eur. J. Entomol., 92: 309-316.
29. Soriano, I.R., I.T. Riley, M.J. Potter and W.S. Bowers, 2004. Phytoecdysteroids: A novel defense against plant-parasitic nematodes. J. Chem. Ecol., 30: 1885-1899.
30. Arnault, C. and K. Slama, 1986. Dietary effects of phytoecdysones in the leek-moth, *Acrolepiopsis assectella* Zell. (Lepidoptera: Acrolepiidae). J. Chem. Ecol., 12: 1979-1986.
31. Elliger, C.A. and A.C. Waiss Jr., 1991. Insect Resistance Factors in *Petunia*. In: Naturally Occurring Pest Bioregulators, Hedin, P.A. (Ed.). American Chemical Society, Washington, DC., USA., ISBN-13: 9780841218970, pp: 210-223.
32. Tanaka, Y. and S. Takeda, 1993. Ecdysone and 20-hydroxyecdysone supplements to the diet affect larval development in the silkworm, *Bombyx mori*, differently. J. Insect Physiol., 39: 805-809.
33. Gilbert, L.I., R. Rybczynski and J.T. Warren, 2002. Control and biochemical nature of the ecdysteroidogenic pathway. Annu. Rev. Entomol., 47: 883-916.
34. Adler, J.H. and R.J. Grebenok, 1995. Biosynthesis and distribution of insect-molting hormones in plants—a review. Lipids, 30: 257-262.
35. Lafont R., 1997. Ecdysteroids and related molecules in animals and plants. Arch. Insect Biochem. Physiol., 35: 3-20.
36. Bathori, M., 2002. Phytoecdysteroids effects on mammals, isolation and analysis. Mini Rev. Med. Chem., 2: 285-293.
37. Maria, A., J.P. Girault, Z. Saatov, J. Harmatha, L. Dinan and R. Lafont, 2005. Ecdysteroid glycosides: Identification, chromatographic properties and biological significance. J. Chromatogr. Sci., 43: 149-157.
38. Gao, X., D. Wang and F. Li, 2000. Determination of ecdysterone in *Achyranthes bidentata* BL. and its activity promoting proliferation of osteoblast-like cells. Acta Pharmaceut. Sin., 35: 868-870.
39. Song, C.Q. and R.S. Xu, 1991. Phytoecdysones from the roots of *Tinospora capillipes*. Chin. Chem. Lett., 2: 13-14.
40. Courtheyn, D., B. Le Bizec, G. Brambilla, H.F. de Brabander and E. Cobbaert *et al.*, 2002. Recent developments in the use and abuse of growth promoters. Anal. Chim. Acta, 473: 71-82.
41. Pis, J., M. Budesinsky, K. Vokac, V. Laudova and J. Harmatha, 1994. Ecdysteroids from the roots of *Leuzea carthamoides*. Phytochemistry, 37: 707-711.

42. Zhang, Y.H., J.K. Cheng, L. Yang and D.L. Cheng, 2002. Triterpenoids from *Rhaponticum uniflorum*. *J. Chin. Chem. Soc.*, 49: 117-124.
43. Bathori, M., H. Kalasz, S.A. Csikkelne and I. Mathe, 1999. Components of *Serratula* species; screening for ecdysteroid and inorganic constituents of some *Serratula* plants. *Acta Pharmaceut. Hung.*, 69: 72-76.
44. Galbraith, M.N. and D.H.S. Horn, 1966. An insect-moulting hormone from a plant. *Chem. Commun.*, 24: 905-906.
45. Jizba, J., V. Herout and F. Sorm, 1967. Isolation of ecdysterone (crustecdysone) from *Polypodium vulgare* L. rhizomes. *Tetrahedron Lett.*, 8: 1689-1691.
46. Ogawa, S. and N. Nishimoto, 1967. [Isolation of the moulting hormones of insects from *Achyranthis radix*]. *J. Pharmaceut. Soc. Jpn.*, 87: 325-327, (In Japanese).
47. Kaplanis, J.N., M.J. Thompson, W.E. Robbins and B.M. Bryce, 1967. Insect hormones: Alpha ecdysone and 20-hydroxyecdysone in bracken fern. *Science*, 157: 1436-1438.
48. Heinrich, G. and H. Hoffmeister, 1967. Ecdyson als begleitsubstanz des ecdysterons in *Polypodium vulgare* L. *Cell. Mol. Life Sci.*, 23: 995-995.
49. Grebenok, R.J., S. Venkatachari and J.H. Adler, 1994. Biosynthesis of ecdysone and ecdysone phosphates in spinach. *Phytochemistry*, 36: 1399-1408.
50. Lafont, R., J. Harmatha, F. Marion-Poll, L. Dinan and I.D. Wilson, 2002. *The Ecdysone Handbook*. 3rd Edn., Cybersales Publ., Prague.
51. Dinan, L., T. Savchenko and P. Whiting, 2002. Chemotaxonomic significance of ecdysteroid agonists and antagonists in the Ranunculaceae: Phytoecdysteroids in the genera *Helleborus* and *Hepatica*. *Biochem. Syst. Ecol.*, 30: 171-182.
52. Thummel, C.S. and J. Chory, 2002. Steroid signaling in plants and insects-common themes, different pathways. *Genes Dev.*, 16: 3113-3129.
53. Handler, A.M., 1982. Ecdysteroid titers during pupal and adult development in *Drosophila melanogaster*. *Dev. Biol.*, 93: 73-82.
54. Laudet, V., 1997. Evolution of the nuclear receptor superfamily: Early diversification from an ancestral orphan receptor. *J. Mol. Endocrinol.*, 19: 207-226.
55. Riddiford, L.M., P. Cherbas and J.W. Truman, 2000. Ecdysone receptors and their biological actions. *Vitamins Hormones*, 60: 1-73.
56. Henrich, V.C., 2005. *The Ecdysteroid Receptor*. In: *Comprehensive Molecular Insect Science*, Gilbert, L.I., K. Iatrou and S.S. Gill (Eds.). Elsevier, Amsterdam, Netherland, ISBN-13: 9780444515162, pp: 243-285.
57. Ma, W.C., 1972. Dynamics of feeding responses in *Pieris brassicae* L. as a function of chemosensory input: A behavioural, ultrastructural and electrophysiological study. Mdedelingen Landbouwhogeschool Wageningen 72-11, pp: 1-162.
58. Schoonhoven, L.M. and I. Derksen-Koppers, 1973. Effects of secondary plant substances on drinking behaviour in some Heteroptera. *Entomol. Exp. Applic.*, 16: 141-145.
59. Jones, C.G. and R.D. Firn, 1978. The role of phytoecdysteroids in bracken fern, *Pteridium aquilinum* (L.) Kuhn as a defense against phytophagous insect attack. *J. Chem. Ecol.*, 4: 117-138.
60. Chou, W.S. and H.S. Lu, 1980. Growth Regulation and Silk Production in *Bombyx mori* L. From Phytoegenous Ecdysterone. In: *Progress in Ecdysone Research*, Hoffman, J.A. (Ed.). Elsevier, North Holland, pp: 281-297.
61. Kubo, I., J.A. Klocke and S. Asano, 1981. Insect ecdysis inhibitors from the East African medicinal plant *Ajuga remota* (Labiatae). *Agric. Biol. Chem.*, 45: 1925-1927.
62. Kubo, I., J.A. Klocke and S. Asano, 1983. Effects of ingested phytoecdysteroids on the growth and development of two lepidopterous larvae. *J. Insect Physiol.*, 29: 307313-311316.
63. Kubo, I. and J.A. Klocke, 1983. Isolation of Phytoecdysone as Insect Ecdysis Inhibitors and Feeding Deterrent. In: *Plant Resistance to Insects*, Hedin, P.A. (Ed.). Chapter 19, American Chemical Society, Washington, DC., USA., ISBN-13: 9780841207561, pp: 329-346.
64. Slama, K., N.K. Abubakirov, M.B. Gorovits, U.A. Baltaev and Z. Saatov, 1993. Hormonal activity of ecdysteroids from certain Asiatic plants. *Insect Biochem. Mol. Biol.*, 23: 181-185.
65. Kubo, B., 1993. Insect Control Agents from Tropical Plants. In: *Phytochemical Potential of Tropical Plants*, Downum, K.R., J.T. Romeo and H.A. Stafford (Eds.). Plenum Press, New York, USA., ISBN: 978-1-4899-1783-6, pp: 133-151.
66. Tanaka, Y., 1995. The different effects of ecdysone and 20-hydroxyecdysone on the induction of larval ecdysis in the silkworm, *Bombyx mori* (Lepidoptera: Bombycidae). *Eur. J. Entomol.*, 92: 155-160.
67. Blackford, M. and L. Dinan, 1997. The tomato moth *Lacanobia oleracea* (Lepidoptera: Noctuidae) detoxifies ingested 20-hydroxyecdysone, but is susceptible to the ecdysteroid agonists RH-5849 and RH-5992. *Insect Biochem. Mol. Biol.*, 27: 167-177.
68. Dhadialla, T.S., G.R. Carlson and D.P. Le, 1998. New insecticides with ecdysteroidal and juvenile hormone activity. *Ann. Rev. Entomol.*, 43: 545-569.
69. Dinan, L., 1989. Ecdysteroid Structure and Hormonal Activity. In: *Ecdysone: From Chemistry to Mode of Action*, Koolman, J. (Ed.). Chapter 36, George Thieme Verlag, Stuttgart, Germany, ISBN-13: 9780865773288, pp: 345-354.

70. Kholodova, Y.D., 2001. Phytoecdysteroids: biological effects, application in agriculture and complementary medicine. *Ukrainskii Biokhimicheskii Zhurnal*, 73: 21-29.
71. Syrov, V.N., Z. Saatov, S.S. Sagdullaev and A.U. Mamatkhanov, 2001. Study of the structure-anabolic activity relationship for phytoecdysteroids extracted from some plants of Central Asia. *Pharmaceut. Chem. J.*, 35: 667-671.
72. Lafont, R. and D.H.S. Horn, 1989. Phytoecdysteroids: Structures and Occurrence. In: *Ecdysone: From Chemistry to Mode of Action*, Koolman, J. (Ed.). Chapter 4, George Thieme Verlag, Stuttgart, Germany, ISBN-13:9780865773288, pp: 39-64.
73. Zibareva, L.N., L. Dinan and V.I. Eryomina, 2007. Screening of *Caryophyllaceae* species for phytoecdysteroids presence. *Rastitel'nye Resursy*, 43: 66-75.
74. Novozhilova, E., V. Rybin, P. Gorovoy, I. Gavrilenko and R. Doudkin, 2015. Phytoecdysteroids of the East Asian *Caryophyllaceae*. *Pharmacogn. Mag.*, 11: S225-S230.
75. Nishimoto, N., Y. Shiobara, M. Fujino, S.S. Inoue and T. Takemoto *et al.*, 1987. Ecdysteroids from *Pfaffia iresinoides* and reassignment of some ¹³CNMR chemical shifts. *Phytochemistry*, 26: 2505-2507.
76. Dinan, L., 2001. Phytoecdysteroids: Biological aspects. *Phytochemistry*, 57: 325-339.