



Journal of Applied Sciences

ISSN 1812-5654

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

A Review of Extraction Technology for Carotenoids and Vitamin E Recovery from Palm Oil

N. Othman, Z.A. Manan, S.R. Wan Alwi and M.R. Sarmidi
 Faculty of Chemical Engineering and Natural Resources Engineering,
 University Teknologi Malaysia, Skudai 81310 Johor, Malaysia

Abstract: Carotenoids and vitamin E (tocopherols, tocotrienols) are among the 1% minor valuable components in crude palm oil (*Elaeis guineensis*). These components have different nutritional functions and benefits to human health. Various technologies have been developed in order to recover these components from being destroyed in commercial refining of palm oil. These include saponification, selective solvent extraction, transesterification followed by molecular distillation and further purification by adsorption using synthetic resins, silica gel and reverse phase C18 silica, adsorption chromatography and membrane technology. Even though there are different technologies, but there is one same feature which is the use of solvent. Solvent plays an important role in most of the technologies. It can be used either as a pre-extraction solvent, main solvent or co-solvent. The problem of most solvents which are used nowadays is that they possess potential fire health and environmental hazards. Due to this, legislation is increasingly restricting the use of certain solvent chemicals. Hence, selection of the most safe, environmentally friendly and cost effective solvent is very important prior to design of alternative extraction methods. Chemical molecular product design is one of the methods that are becoming more popular nowadays for finding solvent with the desired properties prior to experimental testing.

Key words: Carotenoids, vitamin E, extraction, environment-friendly solvent

INTRODUCTION

Crude Palm Oil (CPO) contains approximately 1% of valuable minor components including carotenoids and tocopherols. Carotenes, which impart the distinctive orange-red color to palm oil, together with tocopherols contribute to the stability and nutritional value of the oil.

Both components are important in order to determine the quality characteristics of palm oil (Basiron and Sundram, 1998). Crude palm oil may undergo either physical or chemical refining, the former being the more common refining method in Malaysian palm oil refineries as shown in Fig. 1 (Nagendran *et al.*, 2000). The high temperature and vacuum used during deodorization and




Fig. 1: Commercial refining palm oil

Corresponding Author: Z.A. Manan, Faculty of Chemical Engineering and Natural Resources Engineering,
 University Teknologi Malaysia, Skudai 81310 Johor, Malaysia

deacidification are necessary for the removal of the undesirable components (free fatty acids, oxidation and decomposition products) as much as possible. Unfortunately, these conditions also result in the removal of some of the tocopherols and tocotrienols and the destruction of all the carotenoids present in crude palm oil (Gapor, 1990). Tocopherols, being natural antioxidants, need to be carefully preserved during milling, refining, fractionation and modification of palm oil (Gapor, 1990). Over several decades, various methods have been developed in order to recover carotenoids and vitamin E (tocopherols and tocotrienols) in Crude Palm Oil (CPO). These include saponification (Nesaretnam *et al.*, 2007; Eckey, 1945, 1949) selective solvent extraction (Nitsche *et al.*, 1999; Heidlas *et al.*, 1998; PORIM, 1989; Tanaka, 1986) transesterification followed by molecular distillation (Ooi and May, 2000) and further purification by adsorption using synthetic resins (Ooi *et al.*, 1994; Lion Corporation (1986), silica gel (Tan and Salleh, 1992; Ooi *et al.*, 1991) reverse phase C18 silica (Lion Corporation, 1993) adsorption chromatography (Bonnie, 2007; Goh and Toh, 1988; Baharin *et al.*, 2001) and membrane technology (Baharin *et al.*, 1998). This study reviews the three technologies based on solvent recovery that have been commercialized in industry. It also highlights the use of chemical product design for the selection of the best solvent which is safe, environmentally friendly and economical.

REVIEW OF SOLVENT EXTRACTION TECHNOLOGIES FOR RECOVERY OF CAROTENOIDS AND VITAMIN E FROM CRUDE PALM OIL (CPO)

The extraction technologies that have been developed to recover carotenoids and vitamin E from Crude Palm Oil (CPO) can be divided into three main categories.

Solvent extraction:

Applications: Solvent extraction is a mass transfer operation in which palm oil is contacted with an immiscible or nearly immiscible solvent that exhibits preferential affinity or selectivity towards carotenoids and tocopherols in palm oil. Solvent extraction is one of major commercial importance to the chemical and biochemical industries, as it is often the most efficient method of separation of valuable products from complex feedstock. Some extraction techniques involve partition between two immiscible liquids, others involve either continuous extractions or batch extractions. Because of environmental concerns, suitable solvent which gives less impact to the

environment are more preferred nowadays. The solvent can be a vapour, supercritical fluid, or liquid and the sample can be a gas, liquid or solid.

Solvent used: Solvent extraction which operates at normal condition commonly used hexane as a solvent to extract carotenoids and vitamin E from crude palm oil (Latip *et al.*, 2000; Chiu *et al.*, 2009). However, hexane poses potential fire, health and environmental hazards (Choo *et al.*, 1996). Short-chain alcohols, especially ethanol and isopropanol, have been proposed as alternative extraction solvents due to their greater safety and reduced probability of regulation (Ping and Gwendoline, 2006; IPCS, 1991). Alcohols tend to extract more non-glyceride materials than hexane, due to their greater polarity. Typically, alcohol-extracted oils contain more phosphatides and unsaponifiable compounds (Lusas *et al.*, 1991). Extraction is a process to separate a mixture into fractions or its constituents by suitable solvents. Organic solvents are partly toxic, inflammable and may cause explosive. These disadvantages cause to a lot of an attempt to do in solvent recovery and plant safety. Supercritical fluid is also been used for extraction since it can offer a highly selective process. High pressure technology is being used in Supercritical Fluid Extraction (SFE). Application of supercritical fluid extraction by using supercritical carbon dioxide (SC-CO₂) as solvent has advantages over other solvent extraction methods as carbon dioxide (CO₂) is non-hazardous and non-flammable (Watkins *et al.*, 1994; Johnson and Lusas, 1983) when compared to highly flammable petroleum-based solvent such as hexane or acetone. CO₂ is inert, non-toxic, non-flammable and recyclable besides leaving no residue in the products. Unfortunately, there are a few disadvantages of SFE (Supercritical Fluid Extraction Research Group, 2010) since it requires operating pressures that are typically much higher than ambient conditions, compression of solvent that requires elaborate recycling measures to reduce energy costs and also high capital investment for equipment.

Transesterification and molecular distillation: The solubility of a compound in solvent depends on its molecular weight, polarity and solvent strength (Puah *et al.*, 2007). Triglycerides in CPO have large molecular weights of 807-885. As a result, triglyceride has lower solubility in the solvent. The carotenoids are transported together with triglycerides. The Fatty Acids Methyl Ester (FAME) is more soluble than triglycerides. Transesterification process will convert the large molecules of triglycerides into smaller molecule of Fatty

Acids Methyl Ester (FAME), making it much easier to be separated (Puah *et al.*, 2008). A process involving neutralization and transesterification of palm oil, followed by molecular distillation of the esters are used to recover valuable minor components from crude palm oil. Molecular distillation process represents a special type of vaporization at low pressures and low temperatures. This method is possible to separate and purify molecules which have high molecular weight and thermally sensitive such as the vitamins. However, transesterification process converts the CPO irreversibly to methyl ester which is not edible and furthermore, also changes the quality of oil (Puah *et al.*, 2005). Further purification will be needed to produce crystalline carotenoids and tocopherols safe for consumption.

Adsorption

Applications: Carotenoids and tocopherols extraction by adsorption without a chemical conversion of palm oil has been reported by Ong and Boey (1980) and Mamuro *et al.* (1986). This method however can maintain an edible-oil quality and did not require converting CPO to methyl esters. A process of separating carotene from CPO by adsorption chromatography with a synthetic polymer adsorbent was developed by Goh and Toh (1988). However, this chromatographic process is still not commercially proven and may slow down the refining process if the process is to be introduced in the existing palm oil refining. Therefore, researcher has been trying to modify the process of carotene extraction from CPO by adsorption using synthetic adsorbent that could speed up carotene extraction process and at the same time maintain the edible oil quality of CPO.

Solvent used: Two types of solvent are commonly used in adsorption method. Alcohol such as ethanol and isopropanol (IPA) is used as initial solvent to build an initial layer on the surface area of the adsorbent in order to allow solute contact with liquid phase. The second solvent is n-hexane, which is preferable for eluting the carotene adsorbed on the hydrophobic surface of the adsorbent. The IPA has good solvency for CPO and the solution is almost homogeneous at 50°C. The adsorption ability for carotene in IPA is lower than that in ethanol. On the other hand, ethanol has poor solvency for CPO but more carotene tends to be adsorbed on the column. More ethanol volume will however be required for eluting the oil, because the elution speed of the oil is lower than in chromatography.

SOLVENT DESIGN-CHEMICAL PRODUCT DESIGN

As mentioned previously, solvent plays a major role in most of the technologies to recover carotenoids and

vitamin E from crude palm oil. Traditionally, to find a suitable solvent, researchers will conduct many experiments to test various possible solvents through trial and error approach. This is time consuming and also expensive. Nowadays, chemical product design is becoming more popular for finding a suitable solvent that meet a desirable target properties. Chemical product design method can be used to reduce the search time. Chemical product design is a reverse problem of property prediction, where, given the identity of the molecule and/or the molecular structure, a set of target properties is calculated. Moggridge and Cussler (2000) suggest four principal steps in their chemical product design which are defining the properties required on the product, generate idea that satisfy the target, choose the best set target and lastly produce the product. They proposed the use of a knowledge-based system to guide the user in identifying the target properties as well as selecting the corresponding property values. The selection of target properties should also be closely linked with what can be estimated and what must be measured. The knowledge-based system can help to reduce the number of experiments or to focus on a few specialized measurements from which a number of other target properties may be estimated. An example of existing knowledge based system is ICAS (Integrated Computer Aided System) software which is available at CAPEC (Computer Aided Process Engineering Center), Department of Chemical Engineering from Technical University of Denmark.

CONCLUSION AND FUTURE WORK

Even though there are many technologies to extract valuable minor components (carotenoids, tocopherols and tocotrienols), choosing the right solvent is an important part to determine the performance of the technology. However, there is still a need to search for a safe, environmentally friendly and economical solvent for those technologies. Computer-aided chemical product design can play an important role to screen possible solvents that meets these criteria.

Currently, study is under way at Universiti Teknologi Malaysia's Process Systems Engineering Centre (PROSPECT) to find possible potential solvents which are safe, environmental friendly and economical for recovery of carotenoids and vitamin E from crude palm oil by using chemical product design technique for all the different types of technology.

REFERENCES

- Baharin, B.S., K. Abdul Rahman, M.I. Abdul-Karim, T. Oyaizu, K. Tanaka, Y. Tanaka and S. Takagi, 1998. Separation of palm carotene from crude palm oil by adsorption chromatography with a synthetic polymer adsorbent. *J. Am. Oil Chem. Soc.*, 75: 399-404.
- Baharin, B.S., R.A. Latip, Y.B. Che-Man and R. Abdul Rahman, 2001. The effect of carotene extraction system on crude palm oil quality, carotene composition and carotene stability during storage. *J. Am. Oil Chem. Soc.*, 78: 851-855.
- Basiron, Y. and K. Sundram, 1998. Nutritional research on palm oil and its components. *PORIM Experience Palm Oil Dev.*, 30: 35-40.
- Bonnie, T.Y.P., 2007. Palm carotene concentrate from crude palm oil using vacuum liquid chromatography on silica gel. *J. Oil Palm Res.*, 19: 421-427.
- Chiu, M.C., C.M. Coutinho and L.A.G. Gonçalves, 2009. Carotenoids concentration of palm oil using membrane technology. *Desalination*, 245: 783-786.
- Choo, Y.M., S.C. Yap, C.K. Ooi, A.N. Ma, S.H. Goh and A.S.H. Ong, 1996. Recovered oil from palm-pressed fiber: a good source of natural carotenoids, Vitamin E and sterols. *J. Am. Oil Chem. Soc.*, 75: 599-602.
- Eckey, E.W., 1945. Carotene from palm oil. British Patent 567682.
- Eckey, E.W., 1949. Carotene from palm oil. US Patent 2460796.
- Gapor, M.T.A., 1990. Effect of refining and fractionation on vitamin E in palm oil. Proceedings of the 1989 PORIM International Palm Oil Development Conference, Sept. 5-9, 1989, Palm Oil Research Institute of Malaysia, Kuala Lumpur, Malaysia, pp: 261-265.
- Goh, S.W. and S.K. Toh, 1988. Recovery of carotenoids, tocopherols, tocotrienols and sterols from esterified palm oil. GB 2218989.
- Heidlas, J., G. Huber, J. Cully and U. Kohlrausch, 1998. Process for extraction of carotene from natural sources. US Patent 5714658. <http://www.freepatentsonline.com/5714658.html>.
- IPCS, 1991. n-Hexane Health and Safety Guide. WHO, Geneva.
- Johnson, L.A. and E.W. Lusas, 1983. Comparison of alternative solvents for oils extraction. *J. Am. Oil Chem. Soc.*, 60: 229-242.
- Latip, R.A., B.S. Baharin, Y.B. Che-Man and R. Abdul Rahman, 2000. Evaluation of different types of synthetic adsorbents for carotene extraction from crude palm oil. *Am. Oil Chem. Soc.*, 77: 1277-1282.
- Lion Corporation, 1986. A process for producing carotene from oils and fats. GB 2160874 A.
- Lion Corporation, 1993. Method for purification of carotene containing concentrate. EP 0242148.
- Lusas, E.W., L.R. Watkins and S.S. Koseoglu, 1991. Isopropyl alcohol to be tested as solvent. *Inform.*, 2: 970-976.
- Mamuro, H., Y. Kubota and H. Shiina, 1986. Carotene concentrates. Japanese Patent No. 61282357.
- Moggridge, G.D. and E.L. Cussler, 2000. An introduction to chemical product design. *Chem. Eng. Res. Des.*, 78: 5-11.
- Nagendran, B., U.R. Unithan, Y.M. Choo and K. Sundram, 2000. Characteristics of red palm oil, a carotene and vitamin E-rich refined oil for food uses. *Food Nutr. Bull.*, 21: 189-194.
- Nesaretnam, K., W.Y. Wong and M.B. Wahid, 2007. Tocotrienols and cancer: Beyond antioxidant activity. *Eur. J. Lipid Sci. Technol.*, 109: 445-452.
- Nitsche, M., W. Johannsbauer and V. Jordan, 1999. Process for obtaining carotene from palm oil. US Patent 5902890. <http://www.freepatentsonline.com/5902890.html>.
- Ong, A.S.H. and P.L. Boey, 1980. An improved method for extraction of carotene. British Patent No. 1562794.
- Ooi, C.K., Y.M. Choo and A.S.H. Ong, 1991. Recovery of carotenoids. US Patent 5,019,668.
- Ooi, C.K., Y.M. Choo, S.C. Yap, Y. Basiron and A.S.H. Ong, 1994. Recovery of carotenoids from palm oil. *J. Am. Oil Chem. Soc.*, 71: 423-426.
- Ooi, C.K. and Y.C. May, 2000. Recovery of carotenes. US Patent 6,072,092.
- PORIM, 1989. Recovery of carotenes. UK. Patent Application. GB 2212806 A.
- Ping, B.T.Y. and E.C.L. Gwendoline, 2006. Identification of lutein in crude palm oil and evaluation of carotenoids at various ripening stages of the oil palm fruit. *J. Oil Palm Res.*, 18: 189-197.
- Puah, C.W., Y.M. Choo, A.N. Ma and C.H. Chuah, 2005. Supercritical fluid extraction of palm carotenoids. *Am. J. Environ. Sci.*, 1: 264-269.
- Puah, C.W., Y.M. Choo, A.N. Ma and C.H. Chuah, 2007. Solubility of tocopherols and tocotrienols from palm oil in supercritical carbon dioxide. *J. Food Lipids*, 14: 377-385.
- Puah, C.W., Y.M. Choo, A.N. Ma and C.H. Chuah, 2008. Production of carotenoids and tocopherols concentrate from palm oil using supercritical carbon dioxide. *J. Oil Palm Res., Special Issue-October 2008*: 12-15.

- Supercritical Fluid Extraction Research Group, 2010. Supercritical fluid extraction: Introduction of the physico-chemical properties of the supercritical fluids. <http://sfe.vemt.bme.hu/angol/supercritical.html>.
- Tan, B. and M.H. Salleh, 1992. Integrated process for recovery of carotenoids and tocotrienols from oil. US Patent 05157132. <http://www.freepatentsonline.com/5157132.html>.
- Tanaka, Y., 1986. A process for producing carotene from oils and fats. British Patent 2160874.
- Watkins, L.R., S.S. Koseoglu, K.C. Rhee, E. Hernandez, M.N. Riaz, Jr. W.H. Johnson and S.C. Doty, 1994. New isopropanol system shows promise. *Ibid*, 5: 1245-1253.