



Plant Pathology Journal

ISSN 1812-5387

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Study of the Effects of a Chayotte (*Sechium edule*) Extract on the Plasmid Puc. 9.1 DNA

^{1,2,4}G.F. Diré, ⁴M.F.M.C. Coura, ⁴M.C.L. Almeida, ⁴S.D.D. Vasconcelos,
⁴P.R.A. Siqueira, ⁴R.M. Duarte, ⁴J.S. Rodrigues, ⁴J.C.S. Oliveira,
²M.L. Fernandes and ^{1,3}M. Bernardo-Filho

¹Departamentos de Biofísica e Biometria, Instituto de Biologia Roberto Alcântara Gomes,
Universidade do Estado do Rio de Janeiro, Av. 28 de setembro, 87, Rio de Janeiro

²Departamento de Patologia Geral, Faculdade de Ciências Médicas,
Universidade do Estado do Rio de Janeiro, Av. 28 de setembro, 87, Rio de Janeiro,

³Instituto Nacional do Câncer, Coordenadoria de Pesquisa, Praça Cruz Vermelha,
23, Rio de Janeiro, RJ, Brasil, 20230-130

⁴Laboratórios de Semiologia e Semiotécnica e Fisiologia do Exercício.
Campus R9/Taquara, Universidade Estácio de Sá. Rua André Rocha, 838, Rio de Janeiro

Abstract: Stannous chloride (SnCl₂) is employed as a reducing agent to obtain Technetium-99m-labelled radiopharmaceuticals in nuclear medicine kits, being injected endogenously in humans. Toxic effects of these kits were not studied, thus making it important to evaluate their impact in humans. The use of natural extracts as medicines is growing around the world. The chayotte (*Sechium edule*) is a subtropical vegetable with potent diuretic action. It is used in folk medicine due its hypotensor effect. In this study, plasmid deoxyribonucleic acid (DNA) was exposed to chayotte extracts (macerated) (0.1g mL⁻¹) in presence of stannous chloride (SnCl₂). Samples of the plasmid DNA were analyzed through agarose gel electrophoresis. The results show that the chayotte extract was capable of damaging the DNA in the presence and in the absence of SnCl₂.

Key words: Plasmidial DNA, chayotte, nuclear medicine, stannous chloride

INTRODUCTION

Stannous chloride can cause skin and mucosal irritation in humans and when this salt is injected into laboratory animals, it can produce stimulation and subsequent depression of the central nervous system (Gleason *et al.*, 1969). It has been suggested that SnCl₂ is a powerful genotoxic (McLean *et al.*, 1983; Oliver and Marzin, 1987), mutagenic (Singh, 1983; Tripathy *et al.*, 1990) and carcinogenic (Ashby and Tennant, 1991) compound. In nuclear medicine, SnCl₂ has been employed in scintigraphic test as Technetium-99m (99mTc) reducing agent. Besides the use of SnCl₂ in nuclear medicine, this salt is also used in dentistry (dentifrices) (Hallas and Cooney, 1981; McLean *et al.*, 1983; Rader, 1991; Budavery, 1996).

There are other sources of SnCl₂ to which human beings are exposed to such as from environmental contamination by biocide preparations containing organic compound dimethyl stannous chloride [SnCl₂ (CH₃)₂]

(Hallas and Cooney, 1981). It is hypothesized that the toxicity of SnCl₂ might be mediated by generation of reactive oxygen species (ROS) through the reaction: Sn²⁺ + O₂ + 2H⁺ → Sn⁴⁺ + H₂O₂. The generation hydrogen peroxide undergoes by Fenton reaction to generate [•]OH as follows: Fe²⁺ + H₂O₂ → OH[•] + [•]OH (4). It was also described that SnCl₂ mediates single strand breaks in plasmid DNA through ROS formation in a dose-dependent manner (Dantas *et al.*, 1996). In addition, the mutagenic potentiality of SnCl₂ was identified by *supF* gene mapping (Cabral *et al.*, 1998). It was also determined that *Escherichia coli* (*E. coli*) strains proficient in DNA repair mechanisms were more resistant to SnCl₂ treatment than deficient ones, suggesting that inactivation was due to DNA damage (Aherne and O'Brien, 1999). Biological effects of metals have been reported: (i) transition metals catalyze free radical production that can be related to aging processes and neurodegenerative diseases such as Alzheimer's diseases, Parkinson's disease and others (Stohs and

Bagchi, 1995); (ii) the association between human diseases and metal ions metabolism can be also demonstrated by Huntington's disease and amyotrophic lateral sclerosis; (iii) neurotoxic properties of aluminum and its etiopathogenetic role in Alzheimer's disease are still controversially argued and need further elucidation (Gutteridge *et al.*, 1985); (iv) abnormal copper metabolism detected in the brain tissue of Wilson's and Menke's patients remains to be fully explained (Gutteridge *et al.*, 1985); (v) substantial levels of zinc are found in the hippocampus and its brain deficiency can be the cause of several pathological events (Bettger and O'Dell, 1981), however; (vi) magnesium is successfully used for the treatment of migraine (Stohs and Bagchi, 1995). In addition, the molecular mechanisms underlying neurotoxicity associated with mercury, tin and manganese, also need further investigation. Tin is a heavy metal which has long been regarded as a contaminant of the environment (Wood, 1974). One of its inorganic salts, stannous chloride (SnCl₂), has been widely used in daily human life, to conserve soft drinks, in food manufacturing, as a result of processing and packaging. Studies on the biological effects of SnCl₂ revealed that it can generate reactive oxygen species (ROS) and breaks in deoxyribonucleic acid (DNA) (Caldeira-de-Araújo *et al.*, 1996) and induces lethality in *E. coli*, whose damage recovery depends on RecA-mediated repair (Bernardo-Filho *et al.*, 1994b). Medicinal plants, are mainly complex products with several components with different chemical and pharmacological characteristics (Moro and Basile, 2000). In addition, many of these products are also sold as dietary supplement, but, scientific information about their safe and effective use is hard to find because limited toxicological data are available on herbal remedies and support of rigorous clinical studies is lacking (Capasso *et al.*, 2000). The use of natural products as medicines has been growing in the entire world. Because of this fact, many studies with natural products are being developed and new drugs for treatments of diseases are being discovered. In the literature, the medicinal action mechanism of several plants has been described and different compounds, with various properties, have been isolated from the crude extracts (Leite *et al.*, 1986; Sallé, 1996). *Sechium edule* (Chayotte) a subtropical vegetable with potent diuretic action, is a cucurbitaceous species which is used as food or as medication in popular medicine. It was reported a case of severe hypokalemia pregnancy and that a chayotte preparation was implicated, as the potassium level returned to normal, without recurrence of hypokalemia, once the ingestion of this vegetable stopped (Jensen and Lai, 1986). Gordon (2000) described the hypotensor effect of the

chayotte. Diré *et al.* (2002), have shown that the extracts of chayotte (macerated and infusion) were capable of altering the labeling of blood elements with technetium-99m (99mTc) in a *in vivo* study. In other research Diré *et al.* (2001), have demonstrated that a chayotte extract (macerated) was able to alter the biodistribution of 99mTc as sodium pertechnetate (NaTcO₄) as well as the shape of red blood cells through a qualitative analysis. The effect of stannous ion has been abolished by extracts of some medicinal plants (Reiniger *et al.*, 1999; Melo *et al.*, 2001; Lima *et al.*, 2002, Silva *et al.*, 2002). Bernardo *et al.* (2002), described that the rutin, a compound isolated from *Ruta graveolens*, was not capable of damaging DNA, protecting DNA from the SnCl₂ redox action and inactivating the *Escherichia coli* (*E. coli* AB1157) culture.

Orsolio *et al.* (2005) described that many dietary constituents are chemopreventive in animal models and experiments with cultured cells are revealing various potential mechanisms of action. Compounds classified as blocking agents can prevent, or greatly reduce, initiation of carcinogenesis, or suppressing agents can act on cell proliferation. Caffeic acid (CA) and caffeic acid phenethyl ester (CAPE), members of the polyphenolic compounds, are present in high concentrations in medicinal plants and propolis, a natural beehive product. Based on the findings of the referred author it was postulated that the antitumor activity of polyphenolic compounds includes direct cytotoxic effects on tumor cells.

ROS are generated during a variety of cellular events with beneficial as well as deleterious effects to the organism (Halliwell, 1994). Some plant extracts may increase the effects of the deleterious actions of ROS (Lima *et al.*, 2001). In the present study, we have evaluated the influence of a chayotte extracts on the topology on gel electrophoretic of plasmid DNA submitted to SnCl₂.

MATERIALS AND METHODS

Characterization of the chayotte sample: Chayotte was purchased from a local market in Rio de Janeiro city, RJ, Brazil. To prepare the extract, 50 g of the skin of chayotte were mixture with 500 mL of water in an electric extractor. This preparation was filtered and this extract was considered 100%.

The presence of toxic compounds was evaluated and we did not find them in the extracts of chayotte used in our experiments. The method to verify the presence of these toxic products is based on inhibition of acetylcholinesterase in the presence of the pesticides (Cunha Bastos *et al.*, 1991). In this method, brain acetylcholinesterase is utilized as an *in vitro* detector of

organophosphorus and carbamate insecticides. Briefly, a preparation of acetylcholinesterase was obtained after extraction of a rat brain microsomal fraction with Triton X-100 and was incubated with the extract of cauliflower. Enzyme assay was performed by a potentiometric method based on the formation of acetic acid in the incubation mixture (preparation of acetylcholinesterase and extract of chayotte).

Nucleic acid manipulations: Plasmids were diluted, dispensed into eppendorf tubes (200 ng per tube) and incubated with 200 $\mu\text{g mL}^{-1}$ of SnCl_2 . To evaluate the influence of the extract of the chayotte in DNA breakage, a concentration on a par with 0.1 g mL^{-1} was used. In all cases, reaction mixtures were incubated at 37°C for 40 min. The analysis of the single breaks (SSB) formation was performed using 0.8% agarose gel electrophoresis in order to separate the conformations of plasmid DNA: form I supercoiled native conformation and form II open circle resulting from SSB. Aliquots from each sample (10 μL) were mixed to 2 μL of 6x concentrated loading buffer (0.25% xylene cyanol FF; 0.25% bromofenol blue; 30% glycerol) and applied in a horizontal gel electrophoresis chamber in Tris acetate-EDTA buffer at pH 8.0. After electrophoresis, the gel was stained with ethidium bromide (0.5 $\mu\text{g mL}^{-1}$) and the DNA bands were visualized by fluorescence in an ultraviolet (UV) transilluminator system. Permanent records were performed using a polaroid MP-4⁺system.

RESULTS

Figure 1 shows the electrophoresis in agarose gel of pUC 9.1 plasmid with SnCl_2 and/or the extract of macerated extract.

In Fig. 1 the electrophoretic mobility of plasmidial DNA in various experimental conditions (macerated extract) is shown.




Fig.1: Column 1: (Control: DNA + water), Column 2: (Chayotte 100%), Column 3: (SnCl_2 200 $\mu\text{g mL}^{-1}$), Column 4: (SnCl_2 - 200 $\mu\text{g mL}^{-1}$ + Chayotte 100%), Column 5: (oxidized Chayotte-10min), Column 6: (oxidized SnCl_2 - 10 min), Column 7 (oxidized Chayotte+ SnCl_2 - 10 min) and Column 8 (marker λ hind III). Photos of the gels were scanned

DISCUSSION

Much effort has focused on the identification of phytochemicals in plants, which exert biological effects. The knowledge of these effects are worthwhile and can help to prevent possible undesirable actions of crude extracts and/or purified substances isolated from various plants. Moreover, many times the results reported in the literature are controversies. This fact could be explained by (i) various experimental conditions and models, (ii) the characteristic and the concentration of the used material (crude extract, isolated fraction, purified substance or heated extract) and (iii) the specific condition of the growth of the studied plant. Many biological effects have been associated with the flavonoids and other antioxidant molecules (Aherne and Brien, 1999). Reactive oxygen species (ROS) have been implicated as the primary destructive intermediates in a wide range of environmental conditions as well as in an increasing number of humans disorders (mutagenesis, apoptosis, aging) (Hladik *et al.*, 1987). SnCl_2 has been used as a reducing agent (Bernardo-Filho *et al.*, 1994; Caldeira-de-Araujo *et al.*, 1996) in medical procedures.

Cytotoxic and genotoxic SnCl_2 -induced damage were demonstrated in *E. coli* and the effects appeared to be mediated by ROS (Caldeira-de-Araujo *et al.*, 1999; Dantas *et al.*, 1996,1999; Felzenszwalb *et al.*, 1998; Reiniger *et al.*, 1999).

The treatment of the *E. coli* strains AB1157 with SnCl_2 in presence of *Peumus boldus* (Reiniger *et al.*, 1999), *Cymbopogon citratus*, *Maytenus ilicifolia*, *Baccharis genistelloides* (Melo *et al.*, 2001), Rutin (Bernardo *et al.*, 2002) and with *Brassica oleracea* L. var *botrytis* (Lima *et al.*, 2002) induced protection of cells against the citotoxic- SnCl_2 effects. It could probably be due to oxidant properties of these extracts. However, the intensity of the protective effect against the SnCl_2 effects was dependent on the considered extract. Like observed in the study of the cauliflower extract (Lima *et al.*, 2002) the extract of chayotte was capable of inducing lesion of break type in the plasmid pUC 9.1 DNA (Lima *et al.*, 2001). In comparison with the Bernardo *et al.* (2002), study, it was verified that the rutin different of the chayotte extracts have not induced lesion in the DNA molecule. In this study the extracts of chayotte (macerated) in all concentrations tested have induced lesions in the DNA molecule. In the analysis of *Peumus boldus* extract was noticed that it has reduced or abolished the effect of SnCl_2 , although the lesive effect of boldine was observed when the highest concentration of this substance was used in the presence of the reducing agent despite boldine alone has not been capable of inducing alterations in the DNA. Silva *et al.* (2002), working with different cultures of

E. coli strains also have shown a reduction of the lethal effect induced by SnCl₂ on the survival of the cultures in the presence of *C. citratus*, *B. genistelloides*, *M. ilicifolia* and *P. boldus*. In the present work it was verified that the extract had induced breaks in the DNA molecule.

CONCLUSIONS

In general we can speculate that the extract of chayotte was capable of inducing damages in pUC. 9.1 DNA molecules and considering maybe due to the generation the free radicals which might induced lesions in the DNA molecule.

ACKNOWLEDGMENTS

The present study was supported by CNPq, CAPES, FAPERJ and UERJ.

REFERENCES

- Aherne, S.A. and M.N. O'Brien, 1999. Protection by the flavonoids myricetin, quercetin and rutin against hydrogen peroxidase-induced DNA damage in Caco-2 Nd Hep G2 cells. *Nutr. Cancer*, 34: 160-166.
- Ashby, J. and R.W. Tennant, 1991. Definitive relationships among chemical and mutagenicity for 301 chemicals tested by the U.S. NTP. *Mutat. Res.*, 257: 229-306.
- Bernardo-Filho, M., M.C. Cunha, J.O. Valsa, A. C. Araujo, F.C.P. Silva and A.S. Fonseca, 1994. Evaluation of potential genotoxicity of stannous chloride: inactivation, filamentation and lysogenic induction of *Escherichia coli*. *Food. Chem. Toxicol.*, 32: 477-479.
- Bernardo, L.C., M.B.N. Oliveira, C. R. Silva, F.J. S. Dantas, J.C.P. Mattos, A. Caldeira-de-Araújo, R.S. Moura and M. Bernardo-Filho, 2002. Biological effects of rutin on the survival of *Escherichia coli* AB1157 and on the electrophoretic mobility of plasmid pUC 9.1 DNA. *Cell. Mol. Biol.*, 48: 517-520.
- Bettger, W.J. and B.L.A. O'Dell, 1981. Critical physiological role of zinc in structure and function of biomembranes. *Life Sci.*, 28: 1425-1438.
- Budavery, S., 1996. The Merck Index. Merck. Whitehouse Station, NJ., pp: 1500-1501.
- Cabral, R.E.C., A.C. leitão, C. Lage, A. Caldeira-de-Araújo, M. Bernardo-Filho, F.J.S. Dantas and J.B. Cabral-Neto, 1998. Mutational potentiality of stannous chloride: An important reducing agent in the Tc-99m radiopharmaceuticals. *Mutat. Res.*, 408: 129-135.
- Caldeira-de-Araújo, A., F.J.S. Dantas, M.O. Moraes I. Felzenszwalb and M. Bernardo-Filho, 1996. Stannous chloride participate in the generation of reactive oxygen species. *J. Braz. Assoc. Adv. Sci.*, 48: 109-113.
- Capasso, R., A.A. Izzo, L. Pinto, T. Bifulco, C. Vitobello and M. Mascolo, 2000. Phytotherapy and quality of herbal medicines, *Fitoterapia*, 71: S58-S65.
- Cunha Bastos, V.L.F., J.F. Cunha Bastos and J.S. Lima, 1991. Brain acetylcholinesterase as an *in vitro* detector of organophosphorus and carbamate insecticides in the water. *Water Res.*, 7: 835-840.
- Dantas, F.J.S., M.O. Moraes, J.C.P. De Mattos, R.J.A.C. Bezerra, E.F. Carvalho, M. Bernardo-Filho and A. Caldeira-de-Araújo, 1999. Stannous chloride mediates single strand breaks in plasmid DNA through reactive oxygen species formation. *Toxicol. Lett.*, 110: 129-136.
- Diré, G., E. Lima, D. Mattos, M.B. Oliveira M.J. Pereira, S. Moreno, R. Freitas, M.L. Gomes and M. Bernardo-Filho, 2001. Effect of chayotte (*Sechium edule*) extract on the biodistribution of technetium-99m and on the morphometry of red blood cells. *J. Labelled Cpd. Radiopharm.*, 44: 648-650.
- Dantas, F.J.S., M.O. Moraes, E.F. Carvalho, J.O. Valsa, M. Bernardo-Filho and A. Caldeira-de-Araújo, 1996. Lethality induced by stannous chloride on *Escherichia coli* AB1157: participation of reactive oxygen species. *Food Chem. Toxicol.*, 34: 959-962.
- Diré, G.F., E.A.C. Lima, M.J.S. Pereira, M.B.N. Oliveira, S.R.F. Moreno, D.M.M. Mattos, R.L. Jales and M. Bernardo-Filho, 2002. Effect of a chayotte (*Sechium edule*) extract on the labeling of red blood cells and plasma proteins with technetium-99m: *In vitro* and *in vivo* studies. *Cell. Mol. Biol.*, 48: 751-755.
- Felzenszwalb, I., J.C.P. De Mattos, M. Bernardo-Filho and A. Caldeira-de-Araújo, 1998. Shark cartilage-containing preparation: protection against reactive oxygen species. *Food Chem. Toxicol.*, 36: 1079-1084.
- Gleason, M.N., R.E. Gosselin, C.H. Hodge and R.P. Smith, 1969. *Clinical Toxicology of Commercial Products (Acute Poisoning)*, Williams and Wilkins, Baltimore.
- Gordon, E.A., 2000. The antihypertensive effects of the Jamaican Cho-cho. *West Ind. Med. J.*, 1: 27-31.
- Gutteridge, J.M.C., G.J. Quilan, I. Clarke and B. Hallinwell, 1985. Aluminium salts accelerate peroxidation of membrane lipids stimulated by iron salts. *Biochem. Biophys. Acta*, 835: 441-447.
- Hallas, L.E. and J.J. Cooney, 1981. Tin and tin-resistance microorganisms in Chesapeake Bay. *Applied. Environ. Microbiol.*, 41: 466-471.
- Hallinwell, B., 1994. Free radicals and antioxidants: a personal view. *Nutr. Rev.*, 52: 256-258.
- Hladik III, W.B., G.B. Saha and K.T. Study, 1987. *Essentials of Nuclear Medicine Sci.* Williams and Wilkins, Baltimore, London.

- Jensen, L.P. and A.R. Lai, 1986. Chayote (*Sechium edule*) causing hypokalemia in pregnancy. *Am. J. Obstet. Gynecol.*, 5: 1048-1049.
- Lima, E.A.C., G. Diré, D.M.M. Mattos, M.N. Oliveira, J.C.P. Mattos, F.J.S. Dantas, A. Caldeira-de-Araújo and M. Bernardo-Filho, 2001. Effect of the leaf extract from cauliflower (*Brassica oleracea* L. Var. *Botrytis*) on the biodistribution of the radiopharmaceutical sodium pertechnetate in mice and on the electrophoretic mobility of plasmid pUC 9.1 DNA. *J. Labelled Cpd. Radiopharm.*, 44: 642-644.
- Leite, J.R., M. Seabra, L. de, E. Maluf, K. Assolant D. Suchecki, S. Tufik, S. Klepacz, H.M. Calil and E.A. Carlini, 1986. Pharmacology of lemongrass (*Cymbopogon citratus* Stapf). Assessment of eventual toxic, hypnotic and anxiolytic effects on humans. *J. Ethnopharmacol.*, 17: 75-83.
- Lima, E.A.C., G. Diré, D.M.M. Mattos, R.S. Freitas, M.L. Gomes, M.B.N. Oliveira, M.V.C. Faria, R.L. Jales and M. Bernardo-Filho, 2002. Effect of an extract of cauliflower (leaf) on the labeling of blood elements with technetium-99m and on the survival of *Escherichia coli* AB1157 submitted to the treatment with stannous chloride. *Food. Chem. Toxicol.*, 40: 919-923.
- McLean, J.R.N., D.H. Blakey, G.R. Douglas and J.R. Kaplan, 1983. The effect of stannous and stannic (tin) chloride on DNA in Chinese hamster ovary cells. *Mutat. Res.*, 119: 195-201.
- Melo, S.F., S.F. Soares, R.F. Costa, C.R. Silva, M.B.N. Oliveira, R.J.A.C. Bezerra, A. Caldeira-de-Araújo and M. Bernardo-Filho, 2001. Effect of the *Cymbopogon citratus*, *Maytenus ilicifolia* and *Baccharis genistelloides* extracts against the stannous chloride oxidative damage in *Escherichia coli*. *Mutat. Res.*, 496: 33-38.
- Moro, C.O. and G. Basile, 2000. Obesity and medicinal plants, *Fitoterapia*, 71: S73-S82.
- Oliver, P. and D. Marzin, 1987. Study of genotoxic potential of 48 inorganic derivatives with the SOS chromotest. *Mut. Res.*, 189: 263-269.
- Orsolich, N., S. Terzic, Z. Mihaljevic, L. Sver and I. Basic, 2005. Effects of local administration of propolis and its polyphenolic compounds on tumor formation and growth. *Biol. Pharm. Bull.*, 28: 28-29.
- Rader, J.I., 1991. Anti-nutritive effects of dietary tin. *Adv. Exp. Med. Biol.*, 289: 509-524.
- Reiniger, I.W., C.R. Silva, I. Felzenszwalb J.C.P. Mattos, J.F. Oliveira, F.J.S. Dantas, A.A.C. Bezerra and M. Bernardo-Filho, 1999. Boldine action against the stannous chloride effect. *J. Ethnopharmacol.*, 68: 345-348.
- Sallé, J.L.O., 1996. *Totum Em Fitoterapia*. Robe Editorial, São Paulo, pp: 237.
- Silva, C.R., M.B.N. Oliveira, S.F. Melo F.J.S. Dantas, J.C.P. Mattos, R.J.A.C., Bezerra, A. Caldeira-de-Araújo, A. Duatti and M. Bernardo-Filho, 2002. Biological effects of stannous chloride, a substance that can produce stimulation or depression of the central nervous system. *Brain Res. Bull.*, 59: 213-216.
- Singh, I., 1983. Introduction of reverse mutation and mitotic gene conversion by some metal compounds in *Saccharomyces cerevisiae*. *Mutat. Res.*, 117: 149-152.
- Stohs, S.J. and D. Bagchi, 1995. Oxidative mechanisms in the toxicity of metal ions. *Free Radic. Biol. Med.*, 18: 321-336.
- Tripathy, N.K., F.E. Wurgler and H. Frei, 1990. Genetic toxicity of six carcinogens and six non carcinogens in the *Drosophila* wing spot test. *Mut. Res.*, 242: 169-180.
- Wood, J.M., 1974. Biological cycles for toxic elements in the environment. *Sci.*, 183: 1049-1052.