

## Comparative Nematotoxicity and Fungitoxicity of Crude and Partitioned Three Plant Extracts

Anjorin Samuel Toba

Department of Crop Production, Federal University of Technology, Minna, Nigeria

**Abstract:** The nematotoxicity and fungitoxicity of crude and partitioned ethanolic leaf extracts of *Azadirachta indica*, *Blumea perotitiana* and *Lippia multiflora* were compared *in vitro* involving *Meloidogyne* root-knot nematode and *Rhizoctonia* root-rot fungus. The experiment which involved a Complete Randomized Design (CRD) with three replicates indicated that the nematode mortality and radial growth of the mycelia of *R. solani* applied with ethanolic crude extracts of each of the tested plant leaves was relatively less than their respective partitioned extract. Since, the purification process of natural plant products especially in developing countries is slow and cumbersome and might render pesticidal products to be ineffective, a well-prepared effective crude plant extract will be easier for the resource-poor farmers to afford and renew.

**Key words:** Comparative, nematotoxicity, fungitoxicity crude, partitioned, plant extracts

### INTRODUCTION

*Meloidogyne incognita* and *javanica* are the most pathogenic species of nematode to most crops in the guinea savanna zone of Nigeria (Adesiyun *et al.*, 1990) and could cause up to 64% yield reduction (Khan *et al.*, 1996). They are numerous and adaptable to many soil ecology where they feed on roots, live and reproduce entirely within the soil or root tissue. *R. solani* causes root and stem rots of several young crop plants (Sinclair, 1982). Yield losses of up to 50% have been attributed to *R. solani* attack in cowpea, soyabean and rice (Akem, 1991). The association of nematode and fungus often showed additive interaction leading to more severe disease symptoms and lower yield of several crops.

The use of synthetic pesticides to control pathogens are associated with myriads of problems which include the high cost of procuring and applying them (Salako, 2002). The side or residual effects from their extensive use and their persistence lead to substantial phytotoxicity, soil and water pollution, extermination of beneficial organisms and development of pesticide resistance by many organisms (Kohli *et al.*, 1999). The search for the new alternative fungus-nematode management strategies that are cheap, eco-friendly and environmentally safe is imperative (Rotimi and Moens, 2001). The use of several crude botanicals against pathogenic nematodes and fungi has been reported, but only few farmers or researchers have investigated the partitioned extracts. This study

hereby compared the nematocidal and fungicidal efficacies of the crude and partitioned ethanolic leaf extracts of *Azadirachta indica*, *Blumea perotitiana* and *Lippia multiflora in vitro*. Potent plant-derived portion could be further purified or formulated into botanical pesticide.

### MATERIALS AND METHODS

The leaves of *A. indica* (neem), *B. perotitiana* (iron weed) and *L. multiflora* (Lippia) were collected locally from Kuje and Abuja Municipal blocks of the Federal Capital Territory, Abuja and washed with clean water. The shade-dried leaves were ground into powder with an electric blender and 62.5 g of each leaf powder was cold extracted by soaking in 200 mL ethanol for 3 days. The extractable material was separated from the extraction solvent with Bochi rotary evaporator at 50°C. The jelly-like crude ethanolic extracts was concentrated with water bath and subsequently partitioned by first subjecting 20 g each of the extracts was soxhlet extracted with n-Hexane at 50-60°C for 24 h. The resulting solution was concentrated in rotary vacuum evaporator to yield an oily brown mass of 1.2 g coded H-p. The defatted residual ethanolic extract mass was added with 100 mL of distilled water and agitated properly. The mixture was decanted and filtered with cheese cloth. The filtrate which was a mixture of the extract and water was poured in a separating funnel and successfully partitioned with ethyl acetate and n-butanol to yield 1.1 g golden brown ethyl acetate soluble portion coded Ee-p and a 1.8 g dark brown

n-butanol soluble portion coded Be-p respectively. The residual aqueous dark brown portion mass of 3.2 g was dried and coded Ae-p.

The duly identified preserved sample of *R. solani* inoculum was obtained in the microbiology laboratory of Federal University of Technology, Minna was sub-cultured and grown on a prepared aseptic Oxoid medium in sterile Petri dishes. They were incubated at 28°C and after 72 h, fan-like radial mycelia grew from the rhizomorphs. *Meloidogyne* sp. used was multiplied on *Celosia argentea* in the garden and the inoculum was collected from the roots and rhizosphere of infected plants. In the experimental set up, the potted seeds in each replicate were treated with carbofuran i.e. Furadan® (100 Kg ha<sup>-1</sup>) and Maneb + Zinc i.e. Mycotrin® (0.6 Kg/100 L of water/ha). Also a pot per replicate had 50 mL of distilled water and these served as the control. The experiment involved a Complete Randomized Design (CRD) with 3 replicates. Data collected was subjected to Analysis of variance (ANOVA) and treatment means were separated by Duncan's Multiple Range Test (DMRT) using SAS (1997) package.

## RESULTS AND DISCUSSION

At 24 h exposure, the effect of the crude extract of *B. perotitiana* on percentage mortality of *Meloidogyne* spp. was significantly higher ( $p \leq 0.05$ ) than any other treatments, except that of carbofuran (Table 1). Among the extract treatment, n-hexane soluble portion had the least nematocidal effect but significantly higher than that of the control. At 48 h exposure the carbofuran application recorded the highest percentage mortality (99.33%). This was followed by that of *B. perotitiana* crude extract (86.22%) but was significantly less ( $p \leq 0.05$ ) than that of carbofuran. The n-hexane soluble portion of *B. perotitiana* recorded the least nematocidal effect (35.95%) but this was significantly higher ( $p \leq 0.05$ ) than that of the control (1.33%). At 4 DAI, the effect of mycotrin® treatment had the least mycelia radial growth (1.67 mm). This was followed by the radial growth of neem crude extract applied plots (11.58 g) but was significantly higher ( $p \leq 0.05$ ) than that of the mycotrin® (Table 2).

Among the leaf extracts applied, ethyl acetate soluble portion recorded the least fungistatic efficacy with a radial growth of (32.55 mm). However, it was significantly higher ( $p \leq 0.05$ ) than the radial growth in the control (37.83 mm). At 8 DAI, the trend of mycelia radial growth was similar to that of 4 DAI as it almost increased proportionately in their radial growth. At 8 DAI mycotrin®-applied fungus recorded the highest percentage reduction in the radial

Table 1: Percentage Mortality of *Meloidogyne* spp. exposed to Partitioned Ethanolic leaf extracts for 24 and 48 h

Treatment	Mortality at 24 h exposure(%)	Mortality at 48 h exposure (%)
Neem crude extract	73.38 <sup>b</sup>	82.88 <sup>b</sup>
N-hexane-soluble portion	40.9 <sup>f</sup>	43.73 <sup>e</sup>
Ethyl acetate-soluble portion	49.6 <sup>e</sup>	51.15 <sup>d</sup>
N-butanol-soluble portion	51.6 <sup>de</sup>	55.75 <sup>d</sup>
Blumea crude extract	82.90 <sup>ab</sup>	86.22 <sup>b</sup>
N-hexane-soluble portion	34.13 <sup>g</sup>	35.95 <sup>f</sup>
Ethyl acetate- soluble portion	43.93 <sup>f</sup>	44.18 <sup>e</sup>
N-butanol-soluble portion	53.65 <sup>d</sup>	55.03 <sup>d</sup>
Lippia crude extract	57.98 <sup>e</sup>	61.01 <sup>c</sup>
N-hexane-soluble portion	39.33 <sup>f</sup>	41.48 <sup>e</sup>
Ethyl acetate-soluble portion	42.23 <sup>f</sup>	43.85 <sup>e</sup>
N-butanol-soluble portion	48.50 <sup>e</sup>	54.80 <sup>d</sup>
Carbofuran	91.76 <sup>a</sup>	99.33 <sup>a</sup>
Control (distilled water)	1.16 <sup>h</sup>	1.33 <sup>e</sup>

\*Column means followed by the same letter(s) are not significantly different ( $p \geq 0.05$ ) by DMRT

Table 2: Radial growth of *R. solani* (mm) on Partitioned Ethanolic Extract and Mycotrin®- mended PDA at 4 and 8 DAI

Treatment	Radial 4Days After Inoculation (5000 umg mL <sup>-1</sup> )	Growth (m) 8Days After Inoculation	% Reduction 8Days After Inoculation
Neem Crude Extract	11.58 <sup>h</sup>	13.08 <sup>e</sup>	70.94 <sup>e</sup>
n-Hexane soluble portion	19.83 <sup>f</sup>	24.68 <sup>e</sup>	45.15 <sup>e</sup>
Ethyl acetate soluble portion	24.63 <sup>de</sup>	29.50 <sup>d</sup>	34.44 <sup>d</sup>
n- Butanol soluble portion	14.50 <sup>g</sup>	16.73 <sup>f</sup>	62.89 <sup>f</sup>
Blumea Crude Extract	11.83 <sup>h</sup>	13.43 <sup>g</sup>	70.15 <sup>e</sup>
n-Hexane soluble portion	25.68 <sup>d</sup>	33.23 <sup>c</sup>	26.15 <sup>e</sup>
Ethyl acetate soluble portion	32.55 <sup>b</sup>	31.60 <sup>cd</sup>	29.78 <sup>cd</sup>
n- Butanol soluble portion	14.30 <sup>g</sup>	15.03 <sup>g</sup>	66.60 <sup>f</sup>
Lippia Crude Extract	12.45 <sup>e</sup>	13.75 <sup>g</sup>	69.44 <sup>e</sup>
n-Hexane soluble portion	22.23 <sup>e</sup>	33.93 <sup>c</sup>	24.60 <sup>e</sup>
Ethyl acetate soluble portion	29.15 <sup>c</sup>	39.75 <sup>b</sup>	11.67 <sup>g</sup>
n- Butanol soluble portion	14.83 <sup>g</sup>	16.15 <sup>g</sup>	64.11 <sup>f</sup>
Mycotrin®	1.67 <sup>i</sup>	1.80 <sup>h</sup>	98.22 <sup>h</sup>
Control	37.83 <sup>a</sup>	45.00 <sup>a</sup>	0.00 <sup>a</sup>

\*Column means followed by the same letter(s) are not significantly different ( $p \geq 0.05$ ) by DMRT

growth of the mycelia of *R. solani*. Among the leaf extracts used, neem crude extract had the highest radial growth reduction (70.94%). The least fungistatic treatment was the *Lippia* ethyl acetate soluble portion-applied media with 11.67% radial growth reduction. This might be due to qualitative and quantitative differences in the bioactive principles present in them.

The result of *in vitro* assessment of the fungicidal and nematocidal potential of partitioned ethanolic leaf extracts of the selected plants indicated the bioactivity of group of compounds in the extracts thus could serve as basis for further purification or characterization of the active agents. The higher percentage mortality of *Blumea* crude extract than that of carbofuran at 24 h exposure might be due to the incomplete dissolution of the carbofuran granules and low concentration of the nematocidal active ingredients in the water used for the test. However, at 48 h exposure, highest percentage mortality was observed from carbofuran-applied medium

probably due to the complete dissolution of the granules and the release of the lethal substance which inhibited the enzyme acetyl cholinesterase at the cholinergic synapses in the nematode nervous system (Singh, 2005). Rotimi and Moens (2002) and Tang'an *et al.* (2002) reported that extraction methods employed and the level of concentration of an extract greatly affects the pesticidal efficacy of botanicals.

The radial growth of the mycelia of *R. solani* applied with ethanol crude extracts of each of the tested plant leaves was relatively less than their respective partitioned extract. This might be due to the synergistic relationship of bioactive ingredients in the crude extracts that was capable of exhibiting fungicidal efficacy. As soon as each crude extract is partitioned, their bioactive efficacy dwindles. Rodriguez *et al.* (1996) observed that the quantity of bioactive metabolites in an extract is directly proportional to the mass of the extract. Singh *et al.* (2001) reported that the use of plant crude extract pesticide instead of the formulations from the purified bioactive ingredients is justified since their active ingredients are normally too complex and therefore the process of simplification or isolation may lead to loss of activity.

### CONCLUSION

The purification process of natural plant products especially in developing countries is slow and cumbersome and might render pesticidal products to be ineffective. For a local use, it seems unnecessary to further partition the crude extract in order not to break the bond of synergism in them. A well-prepared crude plant extract could be effective and will be easier for the resource-poor farmers to afford and renew.

### ACKNOWLEDGEMENT

I wish to express my profound gratitude to Mallam Kudu of Department of Microbiology and Mrs P. Fadipe of Department of Chemistry of Federal University of Technology, Minna-Nigeria for their technical support.

### REFERENCES

- Adesiyan, S.O., F.E. Caveness, M.O. Adeniyi and B. Fawole, 1990. Nematode Pests of tropical crops. PTF low price. Heinemann. Educational books (Nigeria) Plc, pp: 9-17.
- Akem, C.N., 1991. Cowpea disease: Biology identification, control and management. IITA Research Guide, pp: 39.
- Khan, M.R., M.W. Khan and A.A. Khan, 1996. Effect of *Meloidogyne incognita* on dry weight, root gall and root nodulation of chickpea and cowpea cultivars. *Test of Agrochemicals and Cultivars*, 17: 70-71.
- Kohli, R.K., D.R. Batish, H.P. Singh and V. Arora, 1999. Allelopathy as a tool for pest and weed management. *J. Punjab Acad. Sci.*, 1: 127-131.
- Rodriguez, I.S.L., J.D. Wolfender J.D. Msonthi and K. Hostettmann, 1996. Quantitative and qualitative analysis of saponins from berries of cultivated *Phytolaca dodecandra* by Lc/Uv and Lc/Ms. *Phytochemistry*, 36: 283-293.
- Rotimi, M.O. and M. Moens, 2001. Potentials of Botanicals as alternative to synthetic nematicides. Paper presented at the 30th Annual Conference of the Nigerian Society for Plant Protection, University of Agriculture, Abeokuta.
- Rotimi, M.O. and M. Moens, 2002. The use of leaf extracts of some herbs in the control of *Meloidogyne incognita*. Paper presented at the 30th Annual Conference of the Nigerian Society for Plant Protection, University of Agriculture, Abeokuta.
- SAS, 1997. Statistical Analysis Software: User's guide for personal computers Release 6.12 SAS Institute, Inc., Cary, N.C. USA., pp: 956.
- Salako, E.A., 2002. Plant protection for the resource-poor farmers a keynote address at Nigerian Society for Plant Protection. 30th Annual Conference. UNAAB. Abeokuta.
- Sinclair, J.B., 1982. Reaction of *Rhizoctonia solani* isolates to certain chemicals. *Plant Dis. Repr.*, 44: 474-477.
- Singh, N.P.I., R.K. Kohli and D.R. Batish, 2001. Allelopathy in Agro-ecosystems. *J. Crop Prod. (Food Products Press)* an imprint of the Haworth Press, Inc., 4 (2): 1-44.
- Singh, R.S., 2005. The Non-Fumigant Nematicides. In: *Plant Diseases*. 8th Edn. Published by Vijay Pramlani for Oxford and IBH, New Delhi, India, pp: 645-683.
- Tang'an, B.N.S., S.A. Adebitan, S.D. Abdul and J.A. Ogidi, 2002. Fungicidal activity of some plant extracts on *Cercospora sojina* Hara: The causal agent of frog-eye leaf spot of soyabean. Paper presented at the 30th Annual Conference of the Nigerian Society for Plant Protection, University of Agriculture, Abeokuta.