

Nutritional Value of Cassava Wastes Ensiled with *Albizia saman* Pod as Feed for Ruminants in Off Season

O.J. Babayemi, U.A. Inyang, O.J. Ifut and L.J. Isaac
Department of Animal Science, University of Ibadan, Nigeria

Abstract: Forages/feed conservation offers strategic and sustainable solutions to the off season feeds for ruminants. Against this background the nutritive value of ensiled cassava wastes and *Albizia saman* was studied. Secondary metabolites and chemical composition of ensiled Cassava Wastes (CSW) with *Albizia saman* Pods (ASP) were determined. *In vitro* gas production of CSW and ASP at 24 h incubation was assessed. The ensiled mixtures were: 100% CSW, 75% CSW+25% ASP, 50% CSW+50% ASP, 25% CSW + 75% ASP and 100% ASP. Saponin was detected in 50% ASP inclusion and 100% ASP silages while tannin was recorded for 100% ASP silage. Crude Protein (CP) content ranged from 4.81% in 100% CSW to 24.50% in 100% ASP silages. The CP values increased significantly ($p < 0.05$) with increasing inclusion of ASP. Metabolizable Energy (ME), Organic Matter Digestibility (OMD) and Short Chain Fatty Acid (SCFA) obtained for all silage mixtures were significantly ($p < 0.05$) different from each other. Potential gas production and potentially degradable fractions (a+b) differed significantly ($p < 0.05$) amongst each other while rate of fermentation did not differ ($p > 0.05$). Total gas production did not differ ($p > 0.05$) at 3 h while for other hours (6-24 h) there were significant ($p < 0.05$) differences with 100% CSW silage being significant over others while 100% ASP had the lowest in all the hours observed. Methane (mL^{-1} 200 mg DM) production ranged from 7-27, the highest being from 100% CSW while the least was observed in 100% ASP. The findings of this study showed that an inclusion level of 50, 75 and even 100% ASP could support small ruminants during period of drought as against sole feeding of cassava wastes.

Key words: Cassava wastes, *Albizia saman* pod, chemical composition, gas production silage, ruminant

INTRODUCTION

The major limitation to ruminant production in many tropical regions is poor nutrition. Treatment of fibrous feedstuffs, supplementation of tropical roughages with leguminous fodder trees and shrubs and low cost nitrogenous sources (Makkar, 2002) and use of agricultural by-products that have little or no utilizable value for man are promising methods to alleviate nutrient deficiencies during the off season. Cassava wastes, an industrial/kitchen waste from cassava tuberous root processing has recently become a feedstuff of great interest in livestock nutrition in Nigeria (Otukoya and Babayemi, 2008). This abundant waste product can be used as a basal diet, source of energy but its low nitrogen content (2-3% CP), unless upgraded cannot sustain growth, production and reproductive functions in ruminants. Fermentation has been used to enhance the nutritive potentials of cassava and its wastes for both human and livestock consumption (Aro, 2008). There is an abundance of leguminous trees/browses for animals to feed on during dry season. Several researches have reported a positive growth response in animals fed leguminous browse plants (Babayemi and Bamikole, 2006;

Babayemi *et al.*, 2006) and pods (Sawe *et al.*, 1998) as supplements. *Albizia saman* is a fast growing, multi-purpose tree found in the tropics useful for its fruits (pods) and leaves as fodder (NFTA, 1995). The leaf is not acceptable to ruminants as the reason might be largely due to the presence of antinutritional factors. The pods are high in crude protein but the seed is hard to digest by ruminants (Jolaosho *et al.*, 2006; Durr, 2001). The *in vitro* Gas Production (GP) method is widely used for evaluation of nutritive value of different classes of feeds. The popularity of *in vitro* GP stems mainly from the ability to exercise experimental control, the capacity to non-destructively screen a large number of substrates, the kinetic information obtained and low costs. Hence, this study was designed to determine the chemical composition, secondary metabolites and *in vitro* gas production of cassava wastes ensiled with *Albizia saman* pod.

MATERIALS AND METHODS

Feed materials/formulation: Cassava wastes, (consisting of peel and skin of tuber) were collected from Mokola in Ibadan and the *Albizia saman* pods were hand picked

from underneath the trees within the campus of the University of Ibadan, Ibadan. The feedstuffs were mixed for silage into five treatments: 100% CSW, 75% CSW+25 % ASP, 50% CSW+50% ASP, 25% CSW+75% ASP and 100% ASP. Salt was added at 0.25% per treatment. The ensiling lasted for 6 months.

Chemical analysis: Dried and ground samples of the silage were used for chemical analysis. Crude protein, crude fibre, ether extract and ash in silage were determined according to methods of AOAC (1990). Neutral Detergent Fibre (NDF), Acid Detergent Fibre (ADF) and Acid Detergent Lignin (ADL) were determined using the methods of Van Soest *et al.* (1991).

In vitro gas production: Rumen fluid was obtained from three West African dwarf female goats. The method for collection was as described by Babayemi and Bamikole (2006) using suction tube in goats. Incubation procedure was as reported by Menke and Steingass (1988). Incubation was carried out at 39±1°C and the volume of gas production was measured at 3, 6, 9, 12, 15, 18, 21 and 24 h. At post incubation period, 4 mL of NaOH (10 M) was introduced to estimate methane production as reported by Fievez *et al.* (2005). Data for gas production were fitted to an exponential equation as proposed by Orskov and McDonald (1979):

$$GP = a + b(1 - \exp^{-ct})$$

Where:

- GP = Gas production (mL) at time t
- a + b = The gas potential production
- c = The rate of gas production (mL⁻¹ h)

The Organic Matter Digestibility (OMD%) and metabolizable energy (ME, MJ kg⁻¹ DM) were calculated.

Secondary metabolites determination: Saponin and tannin was determined qualitatively as described by Babayemi *et al.* (2004). About 2 g of each ground sample were weighed into extraction bottles, using two extraction solvents, namely methanol and petroleum ether.

Statistical analysis: Data were analysed using analysis of variance (SAS, 1999). Significant means were separated using the Duncan (1955) multiple range F-test. Experimental model of the design is:

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij}$$

Where:

- Y_{ij} = Individual observation
- μ = General mean of population
- α_i = Treatment effect
- ε_{ij} = Composite error effect

RESULTS AND DISCUSSION

In Table 1, the Dry Matter (DM) and chemical composition of the ensiled feedstuffs are shown. The highest (24.50%) and the least (4.81%) crude protein content were obtained in 100% ASP and 100% CSW silages, respectively. The CP values showed an increasing trend (p<0.05) as the ASP amount increased in the silage mixtures. There was significant (p<0.05) difference in the NDF, ADF and ADL values. Table 2 shows secondary metabolites of the Cassava wastes ensiled with *Albizia saman* pods. Saponin was detected as low in 50% ASP inclusion and medium in 100% ASP silages while it was negative for others. Tannin was detected in 100% ASP silage alone while it showed a negative observation for other treatments.

Table 3 shows that is the *in vitro* gas production of ensiled CSW together with ASP for 24 h. From the 6-24 h significant (p<0.05) differences were observed in all the treatments with 100% CSW silages being the highest from 9-24 h but similar to 50% inclusion level of ASP silage at the 6th. The 100% ASP silage recorded the lowest gas production. Gas volume increased with increasing hours for all the treatments signifying microbe action on the substrates. Figure 1 shows *in vitro* gas production graphically. Metabolizable Energy (ME), Organic Matter Digestibility (OMD) and Short Chain Fatty Acids (SCFA) of the ensiled mixtures are shown in Table 4. The value

Table 1: Chemical composition (%) of cassava wastes ensiled with *A. saman* pod

Parameters	Silage mixtures					SEM
	100% CSW	75 CSW + 25 ASP	50 CSW + 50 ASP	25 CSW + 75 ASP	100 ASP	
DM	31.79 ^a	46.36 ^b	46.36 ^b	53.44 ^{ab}	60.53 ^a	1.65
ASH	6.00 ^{ab}	7.00 ^a	5.00 ^b	5.00 ^b	5.00 ^b	0.30
CP	4.81 ^d	10.06 ^c	16.19 ^b	16.63 ^b	24.50 ^a	0.10
CF	10.00 ^c	12.00 ^a	8.00 ^d	10.00 ^c	11.00 ^b	0.13
EE	14.00 ^b	14.00 ^b	12.00 ^c	13.00 ^c	15.00 ^a	0.16
NDF	40.00 ^d	38.00 ^d	46.00 ^c	49.00 ^b	53.00 ^a	0.50
ADF	27.00 ^c	30.00 ^b	24.00 ^d	27.00 ^c	42.00 ^a	0.34
ADL	11.00 ^c	11.00 ^c	8.00 ^d	14.00 ^b	20.00 ^a	0.26
CELL	16.00 ^c	19.00 ^b	16.00 ^c	13.00 ^d	22.00 ^a	0.40
HEMC	13.00 ^b	8.00 ^c	22.00 ^a	22.00 ^a	11.00 ^{bc}	0.62

^{a, b, c, d} means on the same row with different superscripts differ significantly (p<0.05)

Table 2: Secondary metabolites of cassava wastes ensiled with *Albizia saman* pod

Treatments (%)	Secondary metabolites		
	Saponin	Tannin 0.2 mL FeCl ₃	Tannin 1.0 mL FeCl ₃
100 CSW	Negative	Negative	Negative
75 CSW+25 ASP	Negative	Negative	Negative
50 CSW+50 ASP	Low	Negative	Negative
25 CSW+75 ASP	Negative	Negative	Negative
100 ASP	Medium	Positive	positive

Table 3: *In vitro* gas production (mL⁻¹ 200 mgDM) of CSW ensiled with ASP for 24 h

Hours	Silage mixtures					SEM
	T1	T2	T3	T4	T5	
3	2.000	4.333	3.333	4.667	2.667	0.46
6	3.667 ^c	11.333 ^a	6.000 ^{bc}	8.667 ^{ab}	6.000 ^{bc}	0.52
9	4.667 ^c	17.000 ^a	8.333 ^{bc}	11.000 ^b	9.000 ^b	0.74
12	5.667 ^c	21.667 ^a	11.667 ^b	13.000 ^b	13.000 ^b	0.82
15	8.667 ^c	24.333 ^a	14.333 ^{bc}	14.667 ^b	15.667 ^b	1.04
18	9.000 ^c	27.000 ^a	16.333 ^b	17.000 ^b	17.333 ^b	0.89
21	10.667 ^c	29.333 ^a	18.667 ^b	20.333 ^b	18.667 ^b	1.21
24	12.667 ^c	33.333 ^a	20.667 ^b	22.333 ^b	21.333 ^b	1.12

^{a,b,c,d} means on the same row with different superscripts differ significantly (p<0.05). T1: 100% ASP; T2: 100% CSW; T3: 25% CSW+75% ASP; T4: 50% CSW+50% ASP and T5: 75% CSW+25% ASP

Table 4: *In vitro* fermentation characteristics of CSW ensiled with ASP for 24 h

Treatment	Fermentation characteristics					Estimated parameters		
	a+b	b	c	Y	t	ME	OMD	SCFA
100 CSW	33.33 ^a	29.00 ^a	0.059	14.33	07.00	7.04 ^a	50.58	0.86 ^a
75C+25A	21.33 ^b	18.67 ^b	0.064	11.33	10.00	5.71 ^b	42.93 ^b	0.57 ^b
50C+50A	22.33 ^b	17.67 ^b	0.042	08.67	06.00	6.18 ^b	45.92 ^{ab}	0.59 ^b
25C+75A	20.67 ^b	17.33 ^b	0.067	12.33	12.00	5.99 ^b	43.99 ^b	0.55 ^b
100ASP	12.67 ^c	10.67 ^c	0.052	07.00	12.00	5.35 ^b	40.42 ^b	0.36 ^c
SEM	1.12	0.96	0.006	01.49	01.32	0.15	1.00	0.03

^{a,b,c,d} means on the same column with different superscripts differ significantly (p<0.05). b = insoluble but fermentable fractions; a+b = potential extent of gas production; c = fractional rate constant; SCFA = Short Chain Fatty Acid; SEM = Standard Error Mean

for the ME, OMD and SCFA ranged from 5.31 in 100% ASP to 7.04 in 100% CSW, 40.42 in 100% ASP to 50.58 in 100% CSW and 0.36 in 100% ASP to 0.86 in 100% CSW, respectively. There were significant differences (p<0.05) among the ensiled mixtures in ME, OMD and SCFA. In all these parameters, 100% CSW silage was observed to be the highest and 100% ASP the lowest. Methane (mL⁻¹ 200 mg DM) production for 24 h as showed in Fig. 2 ranged from 7-27 among the silages, the least being from 25% inclusion of ASP silage (7 mL) and the highest from 100% CSW silage (27 mL).

In most cases, feedstuffs that show high capacity for gas production are also observed to be synonymous for high methane production. The values for CP was higher than the minimum protein requirement of 10-12% recommended by ARC (1985) for ruminants except for 100% CSW which was low (4.81%) and 25% inclusion of ASP which was within range. The increase in CP in the mixtures is as a result of ASP inclusion and these values compares favourably with other leguminous pods as cited




Fig. 1: *In vitro* gas production of cassava wastes ensiled with *Albizia s. pod* at 24 h. T1: 100% ASP; T2: 100% CSW; T3: 25% CSW + 75% ASP; T4: 50% CSW + 50% ASP and T5: 75% CSW + 25% ASP




Fig. 2: Methane gas production of cassava wastes ensiled with *Albizia.s. pod* at 24 h

by other researchers. Ngwa *et al.* (2001) reported that *Leucaena leucocephala* whole pods contained 24.69% CP. Fibre fractions (NDF, ADF and ADL) were higher as the ASP inclusion increased in the mixtures except for some that deviated from this trend as noticed in ADF, fraction. The high fibre values were as a result of hardness seed of the ASP. It has been noted to be indigestible by ruminants (Durr, 2001). Ngwa *et al.* (2001) reported 40.80 and 28.50% for NDF and ADF, respectively in *Leucaena leucocephala* whole pod. A high fibre level occasions indigestibility, low feed intake due to slow degradability of feed and hence reduced performance. The detection of saponins and phenols in the shown study did not agree with that of Otukoya (2007) of absence of Saponin and tannin in pods of *Albizia saman*. However, it agreed with the submission of Hosamani *et al.* (2005) that reported the presence of tannin at a concentration of 2.95%.

The highest gas production was obtained from 100% CSW and the lowest from 100% ASP and this may be attributed to the presence of tannins as shown in Table 2 although, it contained high crude protein. High crude protein in feed enhances microbial multiplication in the rumen which in turn determines the extent of fermentation (Babayemi, 2007). The tannin in ASP might have been responsible for its low gas production and a trend could be observed from Table 3 in that the gas decreased with

increasing inclusion of ASP. However, the observed low value for estimated parameters Table 4 of 100% ASP silage may be due to their low net gas production since gas production reflects more on the digestible energy rather than on protein and fat (Krishna and Gunther, 1987). The low value of the estimated parameters can also be attributed to the high fibre content prevalent in the silage.

The 100% ASP recorded a high content in NDF, ADF and even Acid Detergent Lignin (ADL) values when compared to other silages and were significantly ($p < 0.05$) different. Moreover, the low gas production may also be adduced to the presence of antinutritive factors. Saponin detected in 100% ASP silage is a defaunating agent (Teferedegne, 2000) while tannin reduces ruminal protein degradation due to its great affinity for protein molecules (Frutos *et al.*, 2004). Feedstuffs that are inherent with certain antinutritive factor had been reported to be low in ME and OMD (Aregheore and Abdulrazak, 2005).

This might have accounted for the low values recorded by 100% ASP silage and its mixtures. Methane production indicates energy loss to ruminants and many tropical feedstuffs have been implicated to induce methanogenesis (Babayemi and Bamikole, 2006).

CONCLUSION

The study has revalidated the potential of cassava peel and *Albizia saman* pods as a livestock feed resource in the dry season. Chemical composition showed that the ensiled feedstuffs might not only be practical protein supplements but also as an energy resource. However, 75 and 50% inclusion of ASP may be good enough in supplementing the diet of small ruminants together with an energy diet.

REFERENCES

- AOAC, 1990. Official Methods of Analysis. 15th Edn., Association of Analytical Chemist, Washington DC., pp: 69-88.
- ARC, 1985. Agricultural Research Council. The nutrient requirements of farm animals. ARC, London.
- Aregheore, E.M. and S.A. Abdulrazak, 2005. Estimation of organic matter digestibility and metabolizable energy content of agro-industrial wastes using *in vitro* gas production. Nig. J. Anim. Prod., 32: 79-87.
- Aro, S.O., 2008. Improvement in the nutritive quality of cassava and its by-products through microbial fermentation. Afr. J. Biotech., 7: 4789-4797.
- Babayemi, O.J. and M.A. Bamikole, 2006. Effects of *Tephrosia candida* DC leaf and its mixtures with Guinea grass on *in vitro* fermentation changes as feed for ruminants in Nigeria. Pak. J. Nutr. 5: 14-18.
- Babayemi, O.J., 2007. *In vitro* fermentation characteristics and acceptability by West African dwarf goats of some dry season forages. Afr. J. Biotechnol., 6: 1260-1265.
- Babayemi, O.J., D. Demeyer and V. Fievez, 2004. *In vitro* fermentation of tropical browse seeds in relation to their content of secondary metabolites. J. Anim. Feed Sci., 13: 31-34.
- Babayemi, O.J., F.T. Ajayi, A.A. Taiwo, M.A. Bamikole and A.K. Fajimi, 2006. Performance of West African dwarf goats fed *Panicum maximum* and concentrate diets supplemented with Lablab (*Lablab purpureus*), Leucaena (*Leucaena leucocephala*) and Gliricidia (*Gliricidia sepium*) foliage. Nig. J. Anim. Prod., 33: 102-111.
- Duncan, D.B., 1955. Multiple range and multiple F-tests. Biometrics, 11: 1-42.
- Durr, P.A., 2001. The biology, ecology and agroforestry potential of the rain tree *Samanea saman* (Jacq.) Merr. Agrofor. Syst., 51: 223-237.
- Fievez, V., O.J. Babayemi and D. Demeyer, 2005. Estimation of direct and indirect gas production in syringes: A tool to estimate short chain fatty acid production requiring minimal laboratory facilities. Anim. Feed Sci. Technol., 123: 197-210.
- Frutos, P., G. Hervas, F.J. Giraldez and A.R. Mantecon, 2004. Review. Tannins and ruminant nutrition. Spanish J. Agric. Res., 2: 191-202.
- Hosamani, S.V., N.K.S. Gowda and S.D. Kololgi, 2005. Evaluation of chemical, nutritive and feeding value of rain tree pods. <http://www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20053101705>.
- Jolaosho, A.O., B.O. Oduguwa, O.S. Onifade and O.J. Babayemi, 2006. Effects of ingestion by cattle and immersion in hot water and acid on the germinability of rain tree (*Albizia saman*) seeds. Tropical Grasslands, 40: 244-253.
- Krishna, G. and K.D. Gunther, 1987. The usability of the Hohenheim gas test for evaluating *in vitro* organic matter and protein degradability at rumen level of some agro-industrial by products. Landwirtschliche Forschung, 40: 281-286.
- Makkar, H., 2002. Development and use of rumen molecular techniques for predicting and enhancing productivity. Proposal on New CRP on Rumen Molecular Tools, NAFA. <http://www.naweb.iaea.org/nafa/aph/meetings/gene-proposal.pdf>.
- Menke, K.H. and H. Steingass, 1988. Estimation of the energetic feed value from chemical analysis and *in vitro* gas production using rumen fluid. Anim. Res. Devt., 28: 7-55.

- NFTA, 1995. NFT highlights: A quick guide to useful nitrogen fixing trees from around the world. NFTA 95-02. <http://www.food-security.info/Winrock%20Archive/Albizia%20saman-Winrock.pdf>.
- Ngwa, A.T., I.V. Nsahlai and M.L.K. Bonsi, 2001. The rumen digestion of dry matter, nitrogen and cell wall constituents of the pods of *Leucaena leucocephala* and some *Acacia* species. *J. Sci. Food Agric.*, 82: 1-9.
- Orskov, E.R. and I. McDonald, 1979. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J. Agric. Sci.*, 92: 499-503.
- Otukoya, F.K., 2007. Evaluation of pods of *Albizia saman* as feed supplement for West African dwarf goats. Ph.D. Thesis, University of Ibadan, Ibadan.
- Otukoya, F.K. and O.J. Babayemi, 2008. Supplementation of *Leucaena leucocephala* hay as protein enrichment for Cassava peels in West African dwarf goats. *J. Food Agric. Environ.*, 6: 247-250.
- SAS, 1999. Statistical Analysis System, SAS/STAT Users Guide. SAS Institute Inc., Cary, North Carolina, USA.
- Sawe, J.J., J.K. Tuitoek and J.M. Ottaro, 1998. Evaluation of Common tree leaves or pods as supplements for goats on range area of Kenya. *S. Rum. Res.*, 28: 31-37.
- Teferedegne, B., 2000. New perspective on the use of tropical plants to improve ruminant nutrition. *Proc. Nutr. Soc.*, 59: 209-214.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis, 1991. Methods for dietary fibre, neutral detergent fibre as non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74: 3583-3597.