

Measuring Radon and Radium Concentrations in 120 Samples of Drinking Water Sources, Springs and Rivers of Shandiz, Zoshk and Abrdeh Regions

¹A. Binesh, ²S. Mohammadi, ³A.A. Mowlavi and ²P. Parvaresh

¹Department of Physics, Payam Nour University, Fariman, Iran

²Department of Physics, Payam Nour University, Mashhad, Iran

³Department of Physics, Sabzevar Tarbiat Moallem University, Sabzevar, Iran

Abstract: Radioactive element radium produces radon gas with the alpha decay. Radon is a colorless gas without smell, weak, ineffective and over 50% radiation of the annual dose human body is involved. The two heavy radioactive elements in the natural decay chain from uranium and thorium is produced. Radon through breathing, eating and drinking enters the body. Alpha emission from gas and other radiations emitted from daughter nuclei of its short life makes serious damage to the respiratory system and into the human digestive, therefore after smoking this radiation is the second risk factor of lung cancer. In this study, the concentration of radium and radon in water sources, springs and rivers of Shandiz, Zoshk and Abrdeh regions (Mashhad-Iran) and using light and portable PRASSI system is measured. Total 120 samples including 38 samples of drinking water, 56 river water samples and 26 samples of spring water has been tested. A total of 19 samples had concentrations >11 (Bq L⁻¹), the reference level set by the U.S. Environmental Protection Agency. Radium concentration of all samples was <1 , only sample No. 21 related to drinking water of Shandiz city is about 2.2 (Bq L⁻¹).

Key words: Measuring radon, radium, drinking water, Shandiz, Zoshk and Abrdeh regions, PRASSI system, agency

INTRODUCTION

²²²Rn radioisotope with half-life 3.8 days from the ²³⁸U decay series and ²²⁰Rn with half-life 55 sec from ²³²Th decay series are produced. Radon gas is alpha emitter and enters the body with breathing, eating and drinking. In addition to radon exposure, its daughter nuclei which are very short lived as sediment in the inner membrane respiratory or digestive body remain and increase the body absorption dose (United States Environmental Protection Agency, 1991; UNSCEAR, 1998; ICRP, 1993; IARC, 1988; Mowlavi *et al.*, 2009; Baykara and Dogru, 2006). Natural exposure of people is about 50% of radon gas that many people in the annual risk of cancers of the respiratory and gastrointestinal die. Thus, measurement of radon in water and air is very important and many studies have been done in this area (IARC, 1988; Mowlavi *et al.*, 2009; Baykara and Dogru, 2006; Vogianis *et al.*, 2004; Field *et al.*, 2001; Tayyeb *et al.*, 1998; Yu *et al.*, 1994; Mancini and Giannelli, 1995; Alabdulaaly, 1999; Schmitz and Nickels, 2001; Mortazavi, 2000). This study is the first report measuring radon and radium in drinking water and springs in the region.

MATERIALS AND METHODS

In this study, to measure radon in water samples PRASSI system has been used. This system is lightweight and portable device that has the ability to measure radon concentration in water, soil and air.

Figure 1 shows the system set up of measurement including bubbler and drier column. PRASSI pumping circuit operates with constant fallow rate at 3 L min⁻¹ in order to degassing the water sample properly.

Its detector is a scintillation cell coated with ZnS (Ag) 1830 cm³ volume. The sensitivity of this system in continuous mode is 4 Bq m⁻³ during the integration time 1 h.

Numbers shown by the device is based on Bq m⁻². Using relationship Eq. 1, radon gas density is calculated based on (Bq L⁻¹).

$$Q_{Rn} \left(\frac{Bq}{l} \right) = Q_{PRASSI} \times \frac{V_{tot}(m^3)}{V(l)} \times \left[\exp\left(-\frac{Ln2}{3.8 \times 24} t\right) \right] \quad (1)$$

Where:

- Q_{PRASSI} = The value recorded by the device
- V_{tot} = The total volume of air connections
- V = The volume sample and within the brackets is a correction factor in the delay measurement




Fig. 1: View from the device to measure radon in water

RESULTS AND DISCUSSION

Measuring radon in water samples: In this study, 120 samples of radon concentration of water including drinking water, rivers and springs of Shandiz, Zoshk and Abrdeh regions and adjacent villages (Mashhad-Iran) have been measured. The third column in Table 1, radon concentration samples that have been ordered from low to high. Also, the radon gas density results are shown in histogram of Fig. 2 and 3. Seen that only 83/15% of the samples, the last 19 samples in Table 1 have concentrations >11 ($Bq L^{-1}$) particularly the sample number 120 that related to the spring in the village of Zoshk has concentration about 32 ($Bq L^{-1}$).

Measuring radium in water samples: For measuring radium in water samples, the water samples have kept in the bottles for 35 days to let radon reach the equilibrium with radium. So, by measuring, we obtain radium concentration in the samples. Figure 4 shows the

Table 1: Radon and radium concentration data of different water samples

| Sample number | Water sample | Q_{Ra} ($Bq L^{-1}$) | Q_{Rn} ($Bq L^{-1}$) |
|---------------|--|--------------------------|--------------------------|
| 1 | Zoshk river | 0.000 | 0.000 |
| 2 | River 14 km before Abrdeh | 0.000 | 0.221 |
| 3 | River 1 km before Zoshk | 0.000 | 0.000 |
| 4 | River 10 km before Zoshk | 0.000 | 0.237 |
| 5 | River 2 km after Zoshk | 0.000 | 0.159 |
| 6 | River 6 km after Abrdeh | 0.302 | 0.000 |
| 7 | Abrdeh spring water | 0.327 | 0.000 |
| 8 | Zoshk drinking water (No. 1) | 0.331 | 0.047 |
| 9 | River 1.5 km after Zoshk | 0.379 | 0.097 |
| 10 | Abrdeh drinking water (No. 1) | 0.545 | 0.000 |
| 11 | River of shandiz waterfall (No.1) | 0.555 | 0.654 |
| 12 | River 2.3 km after Abrdeh | 0.590 | 0.095 |
| 13 | River 2.5 km after Zoshk | 0.636 | 0.059 |
| 14 | River 1.3 km after Zoshk | 0.695 | 0.097 |
| 15 | Zoshk drinking water (No. 2) | 0.922 | 0.000 |
| 16 | Shandiz waterfall | 1.041 | 0.177 |
| 17 | River 2.8 km after Abrdeh | 1.180 | 0.163 |
| 18 | River 0.8 km after Zoshk | 1.299 | 0.018 |
| 19 | River 1.8 km after Zoshk | 1.305 | 0.000 |
| 20 | River 2.7 km after Abrdeh | 1.345 | 0.000 |
| 21 | Shandiz drinking water (No.1) | 1.412 | 2.176 |
| 22 | River 2.5 km after Zoshk | 1.525 | 0.059 |
| 23 | Shandiz drinking water (No.2) | 1.994 | 0.000 |
| 24 | Abrdeh drinking water (No. 2) | 1.641 | 0.163 |
| 25 | River 2.3 km after Zoshk | 1.763 | 0.000 |
| 26 | Zoshk drinking water (No. 3) | 1.853 | 0.141 |
| 27 | Upper Abrdeh drinking water (N0.1) | 1.936 | 0.308 |
| 28 | River 0.7 km after Zoshk | 2.241 | 0.096 |
| 29 | Zoshk spring water (No. 1) | 2.307 | 0.034 |
| 30 | River 2.7 km after Zoshk | 2.352 | 0.000 |
| 31 | Shandiz drinking water (No.3) | 2.416 | 0.498 |
| 32 | River 0.8 km after Zoshk spring water | 2.433 | 0.170 |
| 33 | Lower Abrdeh drinking water (No.1) | 2.476 | 0.000 |
| 34 | Shandiz drinking water near the mosque | 2.476 | 0.000 |
| 35 | Shandiz drinking water (No.4) | 2.629 | 0.854 |
| 36 | Upper Abrdeh drinking water (N0.2) | 2.698 | 0.062 |
| 37 | River 5 km after Abrdeh | 2.833 | 0.000 |
| 38 | River 1.7 km after Zoshk | 2.873 | 0.208 |
| 39 | Lower Abrdeh drinking water (N0.2) | 2.876 | 0.000 |
| 40 | Lower Abrdeh spring water | 3.049 | 0.215 |
| 41 | Shandiz drinking water (No.5) | 3.153 | 0.652 |
| 42 | River of shandiz waterfall (No.1) | 3.215 | 0.137 |

Table 1: Continued

| Sample number | Water sample | Q_{Ra} (Bq L ⁻¹) | Q_{Rn} (Bq L ⁻¹) |
|---------------|--------------------------------------|--------------------------------|--------------------------------|
| 43 | Lower Abrdeh drinking water (N0.3) | 3.227 | 0.4910 |
| 44 | River 1.3 km after Zoshk | 3.269 | 0.0000 |
| 45 | River beginning Zoshk | 3.418 | 0.0740 |
| 46 | River 5.5 km after Abrdeh | 3.492 | 0.0000 |
| 47 | Shandiz drinking water (No.6) | 3.619 | 0.7870 |
| 48 | River at Zoshk | 3.796 | 0.0000 |
| 49 | River 5.9 km after Abrdeh | 4.012 | 0.0130 |
| 50 | River 2.4 km after Abrdeh | 4.172 | 0.2850 |
| 51 | River 0.5 km after Zoshk | 4.230 | 0.1330 |
| 52 | Shandiz drinking water (No.7) | 4.231 | 0.0000 |
| 53 | River 1.5 km after Zoshk | 4.237 | 0.0510 |
| 54 | Upper Abrdeh drinking water (N0. 3) | 4.254 | 0.0000 |
| 55 | Upper Abrdeh drinking water (N0. 4) | 4.375 | 0.0000 |
| 56 | River 2.6 km after Abrdeh | 4.729 | 0.0000 |
| 57 | River 1.2 km after Zoshk | 4.883 | 0.0000 |
| 58 | Lower Abrdeh drinking water (N0. 4) | 4.895 | 0.2610 |
| 59 | Shandiz drinking water (No.8) | 4.967 | 0.0000 |
| 60 | Lower Abrdeh drinking water (N0.5) | 5.051 | 0.1108 |
| 61 | River of shandiz waterfall (No. 2) | 5.058 | 0.3170 |
| 62 | River 3.5 km after Abrdeh | 5.081 | 0.0590 |
| 63 | Lower Abrdeh spring water | 5.130 | 0.2440 |
| 64 | River 0.1 km after lower Abrdeh | 5.255 | 0.0000 |
| 65 | River 1.6 km after Zoshk | 5.431 | 0.0570 |
| 66 | Upper Abrdeh spring water | 5.441 | 0.0440 |
| 67 | River 0.2 km after Zoshk | 5.453 | 0.1990 |
| 68 | Abrdeh drinking water (N0. 3) | 5.482 | 0.0000 |
| 69 | River 4 km before Abrdeh | 5.579 | 0.1330 |
| 70 | River 5 km before Abrdeh | 5.675 | 0.0000 |
| 71 | River 0.5 km after Abrdeh | 5.692 | 0.0940 |
| 72 | Zoshk spring water (No. 2) | 5.727 | 0.0000 |
| 73 | Upper Abrdeh drinking water (No. 5) | 6.141 | 0.0870 |
| 74 | Lower Abrdeh drinking water (N0. 6) | 6.574 | 0.0470 |
| 75 | Abrdeh drinking water (N0. 4) | 6.907 | 0.2880 |
| 76 | Spring water 1 km after Zoshk | 7.020 | 0.0000 |
| 77 | Lower Abrdeh drinking water (N0. 7) | 7.111 | 0.2570 |
| 78 | River 2.8 km after Zoshk | 7.150 | 0.0000 |
| 79 | Abrdeh drinking water (N0. 5) | 7.530 | 0.2880 |
| 80 | River 0.2 km after lower Abrdeh | 7.591 | 0.0960 |
| 81 | Lower Abrdeh spring water (No. 1) | 7.631 | 0.1320 |
| 82 | River 2.9 km after Zoshk | 7.867 | 0.2910 |
| 83 | Zoshk spring water (No. 3) | 7.895 | 0.0000 |
| 84 | River 4.5 km after Abrdeh | 7.969 | 0.0000 |
| 85 | Abrdeh drinking water (N0. 6) | 8.131 | 0.1780 |
| 86 | Zoshk drinking water (No. 4) | 8.155 | 0.0580 |
| 87 | Zoshk drinking water (No. 5) | 8.310 | 0.0000 |
| 88 | Zoshk spring water (No. 4) | 8.327 | 0.0000 |
| 89 | River 0.4 km after Zoshk | 8.356 | 0.0000 |
| 90 | Zoshk drinking water (No. 6) | 8.603 | 0.0540 |
| 91 | Lower Abrdeh drinking water (N0. 8) | 8.630 | 0.4370 |
| 92 | Zoshk spring water (No. 5) | 9.034 | 0.1830 |
| 93 | Zoshk spring water (No. 6) | 9.056 | 0.2800 |
| 94 | River 2.5 km after Abrdeh | 9.931 | 0.0189 |
| 95 | River of shandiz waterfall (No. 3) | 10.124 | 0.0000 |
| 96 | Qelqeli spring water | 10.402 | 0.0830 |
| 97 | Zoshk drinking water (No. 7) | 10.721 | 0.0014 |
| 98 | Lower Abrdeh drinking water (N0. 9) | 10.729 | 0.0000 |
| 99 | Zoshk drinking water (No. 8) | 10.915 | 0.0052 |
| 100 | Lower Abrdeh drinking water (N0. 10) | 10.992 | 0.0220 |
| 101 | Shandiz drinking water (No. 9) | 11.199 | 0.0000 |
| 102 | Spring water 0.5 km after Zoshk | 11.360 | 0.1270 |
| 103 | River 1 km before Zoshk | 11.434 | 0.2070 |
| 104 | Lower Abrdeh drinking water (N0. 11) | 11.595 | 0.0960 |
| 105 | River 2 km after Zoshk | 11.778 | 0.4330 |
| 106 | Zoshk spring water (No. 7) | 13.055 | 0.1330 |
| 107 | River 1 km after Zoshk | 13.058 | 0.0910 |
| 108 | Zoshk spring water (No. 8) | 13.761 | 0.0026 |
| 109 | Zoshk spring water (No. 9) | 14.43 | 0.1830 |
| 110 | Spring water 0.1 km after Zoshk | 14.577 | 0.0000 |
| 111 | Spring water 2 km after Zoshk | 14.863 | 0.2070 |
| 112 | Zoshk drinking water (No. 9) | 15.755 | 0.0000 |
| 113 | River 0.5km before Zoshk | 16.324 | 0.0000 |

Table 1: Continued

| Sample number | Water sample | Q_{Ra} (Bq L ⁻¹) | Q_{Rn} (Bq L ⁻¹) |
|---------------|-------------------------------------|--------------------------------|--------------------------------|
| 114 | Spring water at Zoshk | 16.344 | 0.0000 |
| 115 | River of shandiz waterfall (No. 4) | 17.363 | 0.3540 |
| 116 | Upper Abrdeh drinking water (No. 6) | 17.879 | 0.2070 |
| 117 | Lower Abrdeh spring water (No. 2) | 18.445 | 0.0470 |
| 118 | River 1.5 km after Abrdeh | 18.578 | 0.0000 |
| 119 | Spring water 0.7 km after Zoshk | 21.495 | 0.0100 |
| 120 | Spring water 1.5 km before Zoshk | 31.881 | 0.6600 |




Fig. 2: The histogram of radon gas concentration in 120 water samples of Shandiz, Zoshk and Abrdeh regions




Fig. 3: Locations of sampling in Shandiz, Zoshk and Abrdeh regions




Fig. 4: The histogram of radium concentration in different water samples

histogram of radium concentration in different water samples as well as the data are listed in fourth column of Table 1. It is notice that all the radium concentration of samples was <1 (Bq L^{-1}), except sample number 21, drinking water of Shandiz region is about 2.2 (Bq L^{-1}).

CONCLUSION

Measurement results of radon concentration in the water samples shows that only 15.83% sample concentrations are higher than the normal 11 (Bq L^{-1}). This limit by United States Environmental Protection Agency (EPA) as normal is defined and 148 (Bq L^{-1}) is limit the amount of action or reaction that radon should be reduced (Mowlavi *et al.*, 2009). Any sample has not this amount of concentration but most amount of radon concentration with 32 (Bq L^{-1}) is related to spring in Zoshk that is almost one-fifth of the reaction too. Radium concentration of all samples, except sample number 21, drinking water of Shandiz is small and <1 (Bq L^{-1}). Therefore, radon and radium concentration in the water of the regions is not high and this is appropriate.

REFERENCES

Alabdulaaly, A.I., 1999. Occurrence of radon in the central region groundwater of Saudi Arabia. *J. Environ. Radioactivity*, 44: 85-95.
 Baykara, O. and M. Dogru, 2006. Measurements of radon and uranium concentration in water and soil samples from east anatolian active fault systems (Turkey). *Radiat. Meas.*, 41: 362-367.

Field, R.W., D.J. Steck and B.J. Smith, 2001. The lowa radon long gas exposure and lung caner study-phase I: Residential radon gas exposure and long cancer. *Sci. Total Environ.*, 272: 67-72.
 IARC, 1988. Man-made mineral fibres glasswool (group 2B) glass filaments (group 3) rockwool (group 2B) slagwool (group 2B) ceramic fibres (group 2B). *Summar. Evaluat.*, 43: 39-39.
 ICRP, 1993. International commission on radiological protection: Protection against radon-222 at home and at work. *Ann. ICRP*, 23: 1-45.
 Mancini, C. and G. Giannelli, 1995. Determination of waterborne ^{222}Rn concentrations using AC canisters. *Health Physics*, 69: 403-405.
 Mortazavi, S.M.J., 2000. Biological effects of prolonged exposure to high levels of natural radiation in Ramsar, Iran. *Proceeding of International Conference on Radiation and its Role in Diagnosis and Treatment*, Tehran, Iran.
 Mowlavi, A.A., A. Shahbahrami and A. Binesh, 2009. Dose evaluation and measurement of radon concentration in some drinking water sources of the Ramsar region in Iran. *Isotopes Environ. Health Stud.*, 45: 269-272.
 Schmitz, J. and R.M. Nickels, 2001. Exposure to radon and decay products in water works. *Sci. Total Environ.*, 272: 347-348.
 Tayyeb, Z.A., A.R. Kinsara and S.M. Farid, 1998. A study on the radon concentrations in water in Jeddah (Saudi Arabia) and the associated health effects. *J. Environ. Radioactivity*, 38: 97-104.
 UNSCEAR, 1998. United Nations Scientific Committee on the effect of Atomic Radiation: Sources, Effects and Risks of Ionizing Radiation. United Nations, New York.
 United States Environmental Protection Agency, 1991. National primary drinking water regulations radionuclides: Proposed rules. *Fed. Regist.*, 56: 33050-33050.
 Vogiannis, E., D. Nikolopoulos, A. Louizi and C.P. Halvadakis, 2004. Radon variations during treatment in thermal spas of Lesvos Island (Greece). *J. Environ. Radioact.*, 75: 159-170.
 Yu, K.N., Z.J. Guan, M.J. Stokes and E.C. Young, 1994. A preliminary study on the radon concentrations in water in Hong Kong and the associated health effects. *Applied Radiat. Isot.*, 45: 809-810.