

Susceptibility Faecal Isolates of Commensal *E. coli* from Cattle to Fluoroquinolones and Cephalosporins

¹A.O. Ajayi, ²O. Oluduro, ³O.A. Olowe and ^{1,4}O. Famurewa

¹Department of Microbiology, University of Ado-Ekiti, P.M.B. 53536, Ado-Ekiti, Nigeria

²Department of Microbiology, Faculty of Science, Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria

³Department of Medical Microbiology and Parasitology, College of Health Sciences, Ladoke Akintola University of Technology, Osogbo, Nigeria

⁴College of Science, Engineering and Technology, Osun-State University, P.M.B. 4494, Oke-Baale, Osogbo, Nigeria

Abstract: About 500 isolates of *Escherichia coli* were recovered from apparently healthy cattle and tested for their susceptibility to third generation cephalosporins and extended spectrum beta-lactamase producers were detected using the double disk synergy test. Representative isolates were selected on the basis of their cephalosporin resistance patterns for plasmids analysis and mating experiment using *E. coli* 25922 as recipient. Resistance to cefoxitin was highest as 395 (79.0%) showed resistance to the antibiotic while resistance to ceftriaxone was the lowest with 87 (17.4%) isolates. Seventy two isolates were confirmed to be extended spectrum beta-lactamase producers. Two of 13 representative isolates selected among the extended spectrum beta lactamase producers carried single plasmids while the remaining 11 isolates contain multiple plasmids ranging from 1.2-22.2 kb. The cephalosporin-resistant isolates also carried plasmids with sizes ranging from 4.32-32 kb. Twelve isolates among the extended spectrum beta-lactamase producers that carried plasmids successfully transferred the plasmids to the recipient while all the cephalosporin-resistant isolates successful transferred plasmids to the recipient strain. The use of third-generation cephalosporins in animal medicine should be regulated.

Key words: Cephalosporins, animal medicine, regulated plasmids, spectrum, beta-lactamase, Nigeria

INTRODUCTION

The cephalosporins are powerful broad-spectrum antibiotics widely accepted as for use in clinical and animal medicine. They have been successfully utilised for effective treatment and control of human infections and more recently are increasingly being deployed for various uses in animal care and husbandry. They demonstrate vast bactericidal action against gram-positive and negative bacteria and their uses have been extended for use in ruminants, cattle and poultry following the success achieved for their use in human medicine. Furthermore, recent reports have indicated that larger quantities of cephalosporins are being deployed for use in animals and this have fuelled concerns regarding the potential detrimental effect of the use of cephalosporins in humans and livestock and in particular, the emergence of cephalosporin resistant bacteria which have been confirmed to be due to their increasing use in animals (Singer *et al.*, 2008). Another contributory factor for the

increasing incidence of resistance among enteric bacteria to cephalosporins is the use in livestock of third generation cephalosporins which were originally licensed for use for treatment and prevention of infections in humans due to their broad spectrum and bactericidal activity (Tenover, 2006).

For instance, some third-generation cephalosporin with many applications in human medicine including the treatment of severe salmonellosis, diarrhoea and other mild or intestinal diseases caused by enteric bacteria in humans (Whichard *et al.*, 2005).

In addition, their uses in treatment of animal infections, particularly pneumoniae, mastitis, postpartum metritis and necrotising pododermatitis have been documented (Daniels *et al.*, 2009) while they found greater roles in prevention of infections in livestock, enhancement of weight gain to increase beef production and their incorporation into animal food and drinks. The various mechanisms of resistance against cephalosporins have been reported. The most common

mechanism of resistance against third-generation cephalosporins is through the production of extended spectrum beta-lactamases called cephalosporinases that act against the core beta-lactam ring of the antibiotics (Whichard *et al.*, 2005). In gram negative pathogens, cephalosporinases confer resistance against cefotaxime, aztreonam, ceftazidime and ceftriaxone and the cephalosporin-resistant bacteria in livestock have become prevalent worldwide (Livermore *et al.*, 2007; Madec *et al.*, 2008). Extended spectrum beta-lactamases are always coded by plasmids when they occur in gram negative bacteria and various plasmid-mediated cephalosporinases that occur in enteric bacteria have been described (Zioga *et al.*, 2009).

MATERIALS AND METHODS

Bacterial isolation: About 500 isolates of *E. coli* were recovered from faeces of apparently health cattle using sterile faecal swab sticks. The faecal swabs were inoculated directly onto eosine methylene blue agar plates and isolates with morphological appearance of characteristic green metallic sheen were presumptively identified as *E. coli*. All isolates were further confirmed using standard biochemical tests.

Antibiotic susceptibility tests: Antibiotic susceptibility tests were carried out to determine their susceptibility to third-generation cephalosporins and also to determine whether the isolates are extended-spectrum β-lactamase producers using the double disk synergy test. The cephalosporins used were: aztreonam (25 µg), ceftazidime (25 µg), ceftriaxone (25 µg) and ceftazidime (25 µg) (Oxoid, UK). In the disk synergy test, disks containing combination of amoxicillin-clavulanic acid was placed at the center of Mueller-Hinton agar plates and the respective cephalosporin disks were placed between 20-30 mm equidistant from the amoxicillin-clavulanic acid. Plates were incubated at 35°C for 24 h and checked for enhanced zones of inhibition. Those isolates that showed resistance to individual cephalosporins but are non-extended spectrum beta-lactamase producers were also examined and those isolates which showed multiple resistance to the cephalosporins were noted.

Plasmid analysis: Plasmid analysis was carried out on two separate sets of representative isolates: fifteen extended spectrum beta-lactamase producers and non-extended spectrum beta-lactamase cephalosporin-resistant isolates of *E. coli*. The modified alkaline method for plasmid extraction described by Birnboim and Doly (1979) was used for plasmid extraction. Plasmids bands

were separated and using 1% agarose gel electrophoresis and visualised using the ultra violet transillumination.

Conjugation studies: Mating experiments were carried out on all isolates that were confirmed to carry plasmids. The plasmid-free recipient strain *E. coli* 25922 was obtained at the department of Microbiology, Obafemi Awolowo University, Ile-Ife. The donors and the recipient strain were mated in separate tubes in 1:9 vol/vol. transconjugants were selected on Mueller-Hinton agar plates containing ampicillin (50 µg mL⁻¹) and tetracycline (50 µg mL⁻¹). The frequency of conjugation was estimated on all transconjugant plates.

RESULTS AND DISCUSSION

About 72 isolates of *E. coli* showed enhanced zone of diffusion indicating that they are presumptively extended spectrum beta-lactamase producers. Resistance to ceftazidime was highest while resistance to ceftriaxone was the lowest (Table 1). About 158 isolates showed multiple cephalosporin resistance and eleven different cephalosporin resistance patterns were observed among those isolates that showed multiple cephalosporin-resistance (Table 2). Among the representative ESBL producing isolates, five different plasmid profiles were observed comprising between two and four plasmid bands (Fig. 1) while the representatives of the cephalosporin resistant isolates yielded four plasmid

Table 1: Antibiotic resistance of isolates against single cephalosporins

Antibiotics	Number (%) n = 500
Ceftazidime	395 (79.0)
Amoxicillin-clavulanic acid	325 (65.0)*
Ceftazidime	242 (48.4)
Aztreonam	202 (40.4)
Ceftriaxone	87 (17.4)

Table 2: Multiple cephalosporin resistance phenotypes

S/N	No. of cephalosporins	-----Resistance phenotypes-----					No. of isolates
1	2	Amc	Fox				60
2		Fox	Caz				06
3		Atm	Fox				11
Total							77
4	3	Amc	Fox	Caz			06
5		Amc	Atm	Fox			32
6		Atm	Fox	Caz			02
Total							40
7	4	Amc	Atm	Fox	Caz		13
8		Atm	Fox	Caz	Cro		06
9		Amc	Fox	Caz	Cro		02
10		Amc	Atm	Fox	Cro		05
Total							26
11	5	Amc	Atm	Caz	Cro	Fox	12
Total							12

Amc: Amoxicillin-clavulanic acid, Atm: Aztreonam, Caz: Ceftazidime, Cro: Ceftriaxone, Fox: Cefoxitin




Fig. 1: Plasmid profile of representative isolates of extended spectrum β -lactamase producers among *E. coli*; Lane 1: E662, 22 kb; Lane 2: E660, 8.9, 3.7, 1.6 and 12 kb; Lane 3: E647, 933 and 569 kb; Lane 4: E595, 933, 16 and 12 kb; Lane 5: E563, 933 kb; Lane 6: E513, 933, 1.6 and 12 kb; Lane 7: E506, 933, 1.6 and 12 kb; Lane 8: E496, 933, 1.6 and 12 kb; Lane 9: E494, 9.42, 16 and 12 kb; Lane 10: E480, 9.44 and 12 kb; Lane 11: E469, 933, 3.7, 16 and 12 kb; Lane 12: E434, 933, 3.7, 1.6 and 12 kb; Lane 13: E429, 933, 3.7, 1.6 and 12 kb; Lane M: Marker, Lambda Hind III dige 1




Fig. 2: Plasmid profile of representative isolates of cephalosporin-resistant *E. coli* (GEL₁); Lane 1: E649, 686 kb; Lane 3: E643, 4.3 kb; Lane 4: E630, 192 and 433 kb; Lane 5: E608, 559 kb; 432 kb; Lane 7: E563, 792 and 530 kb; Lane 8: E543, 792 kb, 5.30 kb; Lane 9: E410, 7.92 kb

profiles with on or two plasmids (Fig. 2 and 3). In the mating experiment, the frequency of conjugation among the ESBL producing isolates ranged between zero and 3.1×10^{-3} (Table 3) while it ranged between 1.4×10^{-5} and 1.8×10^{-3} among the cephalosporin-resistant isolates (Table 4 and 5).

Five hundred isolates of commensal *Escherichia coli* were recovered from apparently healthy cattle and they were subjected to antibiotic susceptibility tests to determine resistance to third-generation cephalosporins




Fig. 3: Plasmid profile of representative isolates of cephalosporin-resistant *E. coli* (GEL₂); Lane M: Marker, Lambda hind III dige 1; Lane 1: E397, 23.3 kb; Lane 2: E322, 34.0 and 26.7 kb; Lane 3: E155, 233 kb; Lane 4: E157, 34.0 kb; Lane 6: E567, 34.0 and 23.3 kb; Lane 7: E255, 23.3 kb

Table 3: Result for transconjugants selected among extended spectrum β -lactamase producers

Lanes	Isolate code	Frequency of conjugation (Transconjugants /donor cells)	MIC (mg L ⁻¹)
1	E662	1.2×10^{-4}	64
2	E660	2.7×10^{-4}	64
3	E647	1.7×10^{-5}	32
4	E595	2.6×10^{-5}	64
5	E536	0.0	64
6	E513	2.2×10^{-3}	64
7.	E506	2.4×10^{-5}	>64
8	E496	2.0×10^{-4}	32
9.	E494	2.7×10^{-5}	32
10	E480	3.5×10^{-5}	16
11.	E469	1.8×10^{-5}	>64
12	E434	2.9×10^{-5}	32
13	E429	3.1×10^{-3}	64

and also to determine whether they are extended spectrum beta-lactamase producers using the double disk synergy test. First, it was observed in this study that the prevalence of cephalosporin resistance among the five hundred representative isolates was very high with resistance against cefoxitin being highest as 395 (79.0%) isolates showed resistance to the antibiotic while resistance to ceftriaxone was the least.

Third-generation cephalosporins were used because of their heavy use in humans and animals, probably due to their improved activity, enhanced spectrum of activity and improved pharmacokinetic that ensure penetration of body tissues within the shortest time. These improved qualities make them suitable for use in cattle and other animals. It was also observed that some of the isolates are

Table 4: Result for transconjugants selected among cephalosporin-resistant *E. coli*

Lanes	Isolate code	Frequency of conjugation (Transconjugants /donor cells)	MIC (mg L ⁻¹)
1	E 649	2.5×10 ⁻⁵	64
2	E 643	1.4×10 ⁻⁵	64
3	E 630	3.0×10 ⁻⁴	32
4	E 608	1.8×10 ⁻⁵	64
5	E 563	1.7×10 ⁻⁵	64
6	E 543	2.2×10 ⁻⁵	64
7	E 410	1.9×10 ⁻³	16
8	E 397	2.7×10 ⁻⁵	64
9	E 322	3.2×10 ⁻⁴	64
10	E 155	3.6×10 ⁻⁵	64
11	E 157	1.5×10 ⁻⁴	32
12	E 567	2.0×10 ⁻⁵	32
13	E 255	1.8×10 ⁻³	16

Table 5: Plasmid profile of cephalosporin-resistant isolates (GEL 2)

Lanes	Strain code	Plasmid bands (Approx. kb)	Cephalosporin resistance phenotypes
M	-	-	-
1	E 397	23.3	Atm/Fox/Caz/Cro
2	E 322	34.0	Amc/Atm/Fox/Caz
		26.7	
3	E 155	23.3	Amc/Fox/Caz/Cro
4	E 157	34.0	Amc/Fox/Caz/Cro
5	E 71	-	Amc/Atm/Fox/Caz
6	E 567	34.0	Amc/Atm/Fox/Cro
		25.5	
7	E 255	23.3	Amc/Atm/Fox/Caz

Amp: Ampicillin, Col: Colistin, Cot: Cotrimoxazole, Gen: Gentamicin, Nal: Nalidixic-acid, Nit: Nitrofurantoin, Tet: Tetracycline, Strep: Streptomycin, Amc: Amoxicillin-clavulanic acid, Atm: Aztreonam, Fox: Cefoxitin, Cro: Ceftriaxone, Caz: Ceftazidime

extended spectrum β-lactamase producers and this could be responsible for the high incidence of cephalosporin-resistance among the isolates. Production of β-lactamases is the major mechanism through which bacteria develop resistance to cephalosporins. By definition, extended spectrum β-lactamases are enzymes elaborated by organisms and the inactivate penicillins and cephalosporins through hydrolysis of the β-lactam core ring in the antibiotics (Daniels *et al.*, 2009).

In this study, we detected and characterised plasmids that ranged between 22.2-1.2 kb among the *E. coli* isolates which produced extended spectrum beta-lactamases. On the contrary, those plasmids with narrower range of sizes between 23.3 and 4.35 kb were conspicuous among the cephalosporin-resistant isolates. Profiles of multiple plasmids of between two to four plasmids were more predominant among the extended spectrum beta-lactamase producing *E. coli* strains under study while the multiple profiles of plasmids were comparatively less pronounced among the cephalosporin-resistant strains with a maximum of two plasmid bands (Table 6 and 7).

Previous studies have reported and characterised plasmids carried by many enteric bacteria which were confirmed resistant not only to common antibiotics but

Table 6: Plasmid profile of ESBL producing isolates

Lanes	Isolate code	Plasmid bands (Approx.) (kb)	Cephalosporin resistance phenotypes
M	-	-	-
1	E 662	22.2	-
2	E 660	8.9, 3.7, 1.6, 1.2	Cro/Caz
3	E 647	9.33, 569 bp	Fox
4	E 595	9.33, 1.6, 1.2	Fox
5	E 536	9.33	Amc
6	E 513	9.33, 1.6, 1.2	Amc/Fox/Caz
7	E 506	9.33, 1.6, 1.2	Fox/Caz
8	E 496	9.33, 1.6, 1.2	Fox
9	E 494	9.42, 1.6, 1.2	Fox
10	E 480	9.44, 1.2	Fox
11	E 469	9.33, 3.7, 1.6, 1.2	Fox
12	E 434	9.33, 3.7, 1.6, 1.2	Fox
13	E 429	9.33, 3.7, 1.6, 1.2	Amc/Fox

Amp: Ampicillin, Col: Colistin, Cot: Cotrimoxazole, Gen: Gentamicin, Nal: Nalidixic-acid, Nit: Nitrofurantoin, Tet: Tetracycline, Strep: Streptomycin, Amc: Amoxicillin-clavulanic acid, Atm: Aztreonam, Caz: Ceftazidime, Cro: Ceftriaxone, Fox: Cefoxitin

Table 7: Plasmid profile of cephalosporin-resistant isolates (GEL 1)

Lanes	Isolate code	Plasmid bands (Approx.) (kb)	Cephalosporin resistance phenotypes
M	-	-	-
1	E 649	7.9	Amc/Cro/Caz
2*	E 645	-	Amc/Atm/Fox/Caz/Cro
3	E 643	4.32	Amc/Atm/Fox/Caz
4	E 630	7.92, 4.35	Amc/Atm/Fox/Caz/Cro
5	E 608	5.59, 4.32	Amc/Atm/Fox/Caz/Cro
6	E 593	-	Amc/Atm/Fox/Caz/Cro
7	E 563	7.92, 5.30	Amc/Fox/Caz/Cro
8	E 543	7.92	Amc/Atm/Fox/Caz/Cro
9	E 410	7.92	Amc/Atm/Fox/Caz

Amp: Ampicillin, Col: Colistin, Cot: Cotrimoxazole, Gen: Gentamicin, Nal: Nalidixic-acid, Nit: Nitrofurantoin, Tet: Tetracycline, Strep: Streptomycin, Amc: Amoxicillin-clavulanic acid, Atm: Aztreonam, Fox: Cefoxitin, Cro: Ceftriaxone, ceftazidime

more specifically to beta lactam antibiotics. The major findings on the presence of single or multiple plasmids among extended spectrum β-lactamase producing *E. coli* and those isolates that showed resistance to cephalosporins correlate with the findings from such reports (Winokur *et al.*, 2001; Daniels *et al.*, 2007). More specifically, Winokur *et al.* (2001) described plasmids recovered from food animals which confer resistance against ampicillin and some cephalosporins. The presence of multiple resistance plasmids among enteric bacteria from human and animal sources as observed in this present study has been established to have similar clonalities when they are found in the similar environments (Oppegaard *et al.*, 2001). Other enteric bacteria, particularly *Salmonella* isolated from cattle have also been confirmed to carry cephalosporin-resistance plasmids (Aarestrup *et al.*, 2004). Daniels *et al.* (2007), described the clonality of plasmids carrying beta lactamase genes among *Salmonella enterica* and *E. coli* recovered from cattle. This study has shown that generally, plasmids were transferred at relatively high

frequency as revealed by the calculated frequencies of transfer of plasmids detected in some of the representative isolates. All the representative ESBL producing isolates in this study carried plasmids and it was observed that these isolates transferred plasmids at the highest frequency of conjugation compared with the other categories of isolates under study that also carried plasmids. The results obtained from the mating experiment carried out in this study agree with other studies that equally determined the rate of transfer of plasmids among bacteria. In a study to determine the conjugative transfer of ESBL genes among *Salmonella* recovered from poultry and poultry meat, plasmid borne ESBL genes were confirmed to be located on plasmids in *Salmonella* were successfully conjugated with plasmid free recipient that subsequently conferred resistance to the bacteria (Hasman *et al.*, 2005). Furthermore, conjugative transfer of resistant plasmids does not occur only in *E. coli*, other reports have also shown that such plasmids can also be transferred among other bacteria in the Enterobacteriaceae, particularly *Salmonella*, *Enterobacter* and *Shigella* (Dionisi *et al.*, 2009).

It has been noted that enteric bacteria transfer antibiotic resistance plasmids through a fundamentally similar process of conjugation, irrespective of the environments in which such bacteria are found (Garza-Ramos *et al.*, 2007). All the extended spectrum β -lactamase producing and cephalosporin-resistant isolates that carried plasmids in this present research were isolated from cattle and transconjugants were also derived from all isolates at different frequencies of conjugation.

CONCLUSION

In accordance with the results obtained in this present study, several studies have also indicated the derivation of transconjugants as a result of conjugative transfer of plasmids that carry genes that code for extended spectrum β -lactamases (Smet *et al.*, 2009). The high prevalence of resistance to cephalosporins observed in this study could be due to overdependence of the antibiotics for use in animal management in the study location (Singer *et al.*, 2008; Wagner *et al.*, 2008). The implications of the transfer of antibiotic resistance plasmids to other related bacteria are of immense magnitude to environmental and public health. The high prevalence of cephalosporin-resistance among the faecal isolates of *E. coli* from cattle as observed in this present study may be attributed to the extensive transfer of plasmids that code for antibiotic-resistance to other related bacteria within the same environment as confirmed by the mating experiments. The

high frequency of transfer of antibiotic resistance plasmids among enteric bacteria connotes that such bacteria could easily be disseminated into the immediate environment through faecal shedding and such bacteria could contaminate the food chain which is a crucial route through which such bacteria could infect humans through intake of food and drinking water that are contaminated with a sizeable population of antibiotic-resistant enteric bacteria (Witte, 1998). Stringent regulatory regimes for the use of cephalosporins in animals must be enforced to curb the high incidence of cephalosporin resistance in animals, particularly cattle.

REFERENCES

- Aarestrup, F.M., H. Hasman, I. Olsen and G. Sørensen, 2004. International spread of bla_{CMY-2}-mediated cephalosporin resistance in a multi-resistant *Salmonella enterica* serovar heidelberg isolate stemming from the importation of a boar by denmark from Canada. *Antimicrob. Agents Chemother.*, 48: 1916-1917.
- Birnboim, H.C. and J. Doly, 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.*, 7: 1513-1523.
- Daniels, J.B., D.R. Call and T.E. Besser, 2007. Molecular epidemiology of bla_{CMY-2} plasmids carried by *Salmonella enterica* and *Escherichia coli* isolates from cattle in the pacific northwest. *Applied Environ. Microbiol.*, 73: 8005-8011.
- Daniels, J.B., D.R. Call, D. Hancock, W.M. Sischo, K. Baker and T.E. Besser, 2009. Role of ceftiofur in selection and dissemination of bla_{CMY-2}-mediated cephalosporin resistance in salmonella enterica and commensal *Escherichia coli* isolates from cattle. *Applied Environ. Microbiol.*, 75: 3648-3655.
- Dionisi, A.M., C. Laucarli, S. Owczarek, I. Luzzi and L. Villa, 2009. Characterization of plasmids-borne quinolone resistance gene qnrB19 in *Salmonella enterica* serovar typhimurium. *Antimicrob. Agents Chemother.*, 53: 4019-4021.
- Garza-Ramos, U., E. Martinez-Romero and J. Silva-Sanchez, 2007. SHV-type Extended-spectrum β -lactamase (ESBL) are encoded in related plasmids from enterobacteria clinical isolates from Mexico. *Salud Publica Mexico*, 49: 415-421.
- Hasman, H., D. Mevius, K. Veldman, I. Olesen and M.F. Aarestrup, 2005. Beta-lactamases among Extended Spectrum Beta Lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in the Netherlands. *J. Antimicrob. Chemther.*, 56: 115-121.

- Livermore, D.M., R. Canton, M. Gniadkowski, P. Nordmann and G.M. Rossolini *et al.*, 2007. CTX-M: Changing the face of extended spectrum β -lactamases in Europe. *J. Antimicrob. Chemother.*, 59: 165-174.
- Madec, J., S. Martin, P. Libereton and T. Rambaud, 2008. Prevalence of fecal carriage of acquired expanded-spectrum cephalosporin resistance in *Enterobacteriaceae* strains from cattle in France. *J. Clin. Microbiol.*, 46: 1566-1567.
- Oppegard, H., T.M. Steinum and Y. Wasteson, 2001. Horizontal transfer of a multi-drug resistance plasmid between coliform bacteria of human and bovine origin in a farm environment. *Applied Environ. Microbiol.*, 67: 3732-3734.
- Singer, R.S., S.K. Patterson and R.L. Wallace, 2008. Effects of therapeutic ceftiofur administration to dairy cattle on *Escherichia coli* dynamics in the intestinal tract. *Applied Environ. Microbiol.*, 74: 6956-6962.
- Smet, A., A. Martel, D. Persoons, J. Dewulf and M. Heyndrickx *et al.*, 2009. Comparative analysis of extended-spectrum- β -lactamase-carrying plasmids from different members of *Enterobacteriaceae* isolated from poultry, pigs and humans: Evidence for a shared β -lactam resistance gene pool. *J. Antimicrob. Chemother.*, 63: 1286-1288.
- Tenover, F.C., 2006. Mechanisms of antimicrobial resistance in bacteria. *Am. J. Med.*, 119: 3-10.
- Wagner, B.A., B.E. Straw, P.J. Fedorka-Cray and D.A. Dargatz, 2008. Effect of antimicrobial dosage regimen on salmonella and *Escherichia coli* isolates from feeder Swine. *Applied Environ. Microbiol.*, 74: 1731-1739.
- Whichard, J.M., K. Joyce, P.D. Fey, J.M. Nelson, F.J. Angulo and T.J. Barrett, 2005. Beta-lactam resistance and *Enterobacteriaceae*, United States. *Emerg. Infect. Dis.*, 11: 1464-1466.
- Winokur, P.L., D.L. Vonstein, L.J. Hoffman, E.K. Uhlenhopp and G.V. Doern, 2001. Evidence for transfer of CMY-2 AmpC β -lactamase plasmids between *Escherichia coli* and *Salmonella* isolates from food animals and humans. *Antimicrob. Agents Chemother.*, 45: 2716-2722.
- Witte, W., 1998. Medical consequence of antibiotic use in agriculture. *Science*, 279: 996-997.
- Zioga, A., J.M. Whichard, S.D. Kotsakis, L.S. Tzouvelekis, E. Tzelepi and V. Miriagou, 2009. CMY-31 and CMY-36 cephalosporinases encoded by ColE1-like plasmids. *Antimicrob. Agents Chemother.*, 53: 1256-1259.