

Effect of Dietary Humic Acid on the Oxidative Status in Liver, Meat and Eggs of Quails

¹H. Durmaz, ²M. Avci, ³S. Selek and ³H. Celik

¹Department of Food Hygiene and Technology, Faculty of Veterinary Medicine,
Harran University, Sanliurfa, Turkey

²Department of Animal Nutrition Faculty of Veterinary Medicine,
Harran University 63300 Sanliurfa, Turkey

³Department of Biochemistry, Faculty of Medicine, Harran University, Sanliurfa, Turkey

Abstract: This study was carried out to examine the influence of dietary humic acid on the oxidative status in liver, meat and eggs of quail. A total of 260 Japanese quails, one week of age, were randomly assigned to one control and three experimental groups. The quails were fed with a basal diet (H0), the basal diet supplemented with 360 mg kg⁻¹ (H1), 480 mg kg⁻¹ (H2) or 600 mg kg⁻¹ (H3) humic acid. After 9 weeks of feeding, the all quails were slaughtered. Liver, meat and eggs were analyzed for antioxidant and oxidant parameters. Liver samples of H1, H2 and H3 groups showed a higher degree of oxidant parameters (indicated by a lower antioxidative capacity) than those of H0 samples. However, the antioxidant and oxidant parameters in the meat and eggs samples were not affected by adding humic acid. Our findings indicated safety of humic acid for meat and eggs of quail and support the use of humic acid as a food additive.

Key words: Quail, humic acid, oxidant, antioxidant, parameters, additive

INTRODUCTION

In recent years, it has been observed that humates included in feed and water of poultry promote growth. Humic acids stabilize the intestinal flora and thus ensure an improved utilization of nutrients in animal feed. This leads to an increase in live weight of the animal without increasing the amount of feed given to the animal (Bailey *et al.*, 1996; Shermer *et al.*, 1998). However, humic acid is also believed to be a semiquinonetype radical (Rex, 1960). In addition, humic acid has been shown to be a toxic factor for many mammalian cells, including chondrocytes and erythrocytes (Liang *et al.*, 1998, 1999; Cheng *et al.*, 1999). However, to our knowledge, effect of humic acid on the oxidant and antioxidant potential of liver, meat and eggs of quails have not been studied. The objective of the present study was to investigate the effect of supplementation of humic acid at different levels on antioxidant-oxidant status in the liver, meat and eggs of quails.

MATERIALS AND METHODS

Quails, diets and slaughter: In this research, a total of 260 Japanese quails (*Coturnix coturnix japonica*) at one

weeks of age were used. The birds were randomly assigned to one control and three experimental groups based on their initial body weight, comprising five replicates of 13 birds each. They were fed a basal diet (H0), the basal diet supplemented with either 360 (H1), 480 (H2) or 600 (H3) mg kg⁻¹ of humic acid. Ingredients and chemical compositions of the basal diets are shown in Table 1. Chemical compositions of the diets were analyzed using the international procedures of AOAC (1994). The study was lasted 9 weeks. At the end of the feeding period, 15 birds were selected at random from each treatment group. Prior to slaughtering, the birds were held without feed for 10 h and then they were slaughtered by a neck cut. The birds were eviscerated manually, washed and allowed to drain 10 min. After eviscerating, carcasses were frozen at -20°C until further use. After eggs were collected daily, whole eggs were wrapped in Parafilm® and stored at 4°C for later analysis.

Measurement of antioxidative status: Total Antioxidant Capacity (TAC) levels were determined using a novel automated colorimetric measurement method developed by Erel (2004). Catalase (CAT) activity was assayed using a method described by Goth (1991).

Table 1: Ingredients and chemical analyses of diets fed to quails (g kg⁻¹)

Ingredients	H0	H1	H2	H3		H0	H1	H2	H3
Maize	554.50	553.55	552.00	550.95	Chemical analysis				
Soybean meal	364.00	363.00	363.50	364.00	Crude protein	241	240	240	241
Fish meal	51.00	51.00	51.00	51.00	Calcium	8.1	8.0	8.1	8.1
Vegetable oil	11.00	11.00	11.00	11.00	Total phosphorus	7.2	7.1	7.2	7.1
DCP	3.50	3.50	3.50	3.50	Calculated values				
Ca	11.00	10.70	11.00	10.80	ME (MJ kg ⁻¹)	12.129	12.121	12.129	12.155
Salt	2.50	2.50	2.50	2.50	Lysine	13.8	13.8	13.8	13.8
Vit and Min ^a	2.50	2.50	2.50	2.50	Met+cys	8.0	8.0	8.0	8.0
Humic acid (16%)	--	2.25	3.00	3.75					
Total	1000	1000	1000	1000					

^aVitamin and mineral premix provided the following per kg diet: Vitamin A, 12500 IU; Vitamin D3, 1500 IU; Vitamin E, 31.25 mg; Vitamin K3, 3.75 mg; Vitamin B1, 2.5 mg; Vitamin B2, 7.5 mg; Niacin 25 mg; Cal. D-pantothenate 10 mg; Vitamin B6, 5mg; Vitamin B12, 0.019 mg; Folic acid 1 mg; Choline chloride 250 mg; Mn 100 mg; Fe 75 mg; Zn 75 mg; Cu 6.25 mg; Co 0.25 mg; I, 1.25 mg; Se 0.19 mg

Table 2: Effect of humic acid on the oxidative status in the liver and meat*

	Liver				Meat			
	TAC	LPO	TOS	CAT	TAC	LPO	TOS	CAT
H0	1.20±0.04 ^a	1.49±0.44 ^{bc}	1.54±0.30 ^c	106.42±4.12	0.71±0.03	0.55±0.18	0.79±0.14	11.38±03.36
H1	1.03±0.12 ^{ab}	1.40±0.29 ^c	2.30±0.14 ^b	98.72±4.23	0.65±0.10	0.56±0.15	0.88±0.32	10.57±01.49
H2	1.02±0.21 ^{ab}	2.01±0.52 ^b	2.82±0.71 ^b	94.12±5.90	0.67±0.08	0.60±0.20	0.85±0.04	08.31±01.89
H3	0.96±0.13 ^b	2.56±0.30 ^a	3.61±0.28 ^a	95.60±8.00	0.59±0.05	0.79±0.18	0.91±0.10	08.21±03.57
P	*	**	**	*	ns	ns	ns	ns

* TAC: mmol Trolox Eqv./L; LPO: µmol H₂O₂ Eqiv/L; TOS: µmol H₂O₂ Eqiv/L; CAT: U/L

Table 3: Effect of humic acid on the oxidative status in the eggs*

	Egg white			Egg yolk		
	TAC	LPO	TOS	TAC	LPO	TOS
H0	0.06±0.04	0.16±0.04	0.32±0.07	0.35±0.07	1.51±0.17	2.50±0.21
H1	0.10±0.02	0.22±0.04	0.28±0.11	0.34±0.09	1.41±0.21	2.86±0.29
H2	0.11±0.03	0.20±0.07	0.33±0.10	0.39±0.09	1.60±0.17	2.66±0.15
H3	0.13±0.02	0.25±0.04	0.40±0.10	0.40±0.05	1.43±0.28	2.87±0.42
P	ns	ns	ns	ns	ns	ns

* TAC: mmol Trolox Eqv./L; LPO: µmol H₂O₂ Eqiv/L; TOS: µmol H₂O₂ Eqiv/L

Measurement of oxidative status: Total Oxidant Status (TOS) were determined by using a new automated colorimetric method developed by Erel (2005). Levels of Lipid Peroxidation (LPO) were determined using an automated method described by Khelifa and Steghens (Khelifa and Steghens, 2004).

RESULT AND DISCUSSION

The oxidant and antioxidant status in liver and meat samples are shown in Table 2. Humic acid supplementation significantly affected the levels of oxidant (p<0.01) and antioxidant (p<0.05) parameters in livers. The levels of TOS and LPO in livers were highest in the H3 samples, followed by the H2, H1 and H0 samples. The TAC and CAT values of H1, H2 and H3 samples were significantly higher than H0 samples liver tissue. Oxidant levels and antioxidant enzyme activities have been evaluated in several studies. Although conflicting results have been reported (Ramachandran and Iyer, 1984; Jansson *et al.*, 1985), it is generally accepted that oxidative status is increased through

increase in oxidant levels and/or decrease in antioxidant enzyme capacities. In our study we indicated that antioxidant levels decreased when oxidant levels increased in liver. On the other hand, the levels of TOS, LPO, TAC and CAT in meat and eggs from H1, H2 and H3 groups were not different compared with those from H0 group (Table 2 and 3).

There is limited information about the effect of dietary humic acid on oxidant and antioxidant status in meat and eggs for discussion of the results. In only one study, Aksu *et al.* (2005) reported there is the effect of dietary humate on the lipid oxidation in broilers meat. In our study, meat and eggs of quails were not affected by humic acid supplementation. Ho *et al.* (2003) reported that the increase in lipid peroxidation has an almost linear relationship with the increase in humic acid concentration from 0 to 50 µg mL⁻¹ in liver. Humic acid concentrations in the tissues were not determined in this study. However, it might be suggested that humic acid-fed quails were unable to absorb and transfer more humic acid in their meat and eggs than in their liver.

CONCLUSION

The results of our study indicate safety of humic acid for meat and eggs and support the use of humic acid as a food additive. However, further studies are required to determine the humic acid concentration in tissue from humic acid-fed quails for oxidant status.

REFERENCES

- Aksu, M.I., M. Karaoglu, M. Kaya, N. Esenbuga and M. Macit, 2005. Effect of dietary humate on the pH, TBARS and microbiological properties of vacuum-and aerobic-packed breast and drumstick meats of broilers. *J. Sci. Food Agric.*, 85: 1485-1491.
- AOAC, 1994. Official Methods of Analysis of the Association of Official Chemists. 14th Edn., Inc. Arlington. Erel, O.A., 2004. Novel automated method to measure total antioxidant response against potent free radical reactions. *Clin. Biochem.*, 37: 112-119.
- Bailey, C.A., K.E. White and S.L. Donke, 1996. Evaluation of Menefee Humate™ on the performance of broilers. *Poult. Sci. (Abstr.)*, 75(Suppl. 1): 84.
- Cheng, M.L., H.Y. Ho, D.T. Chiu and F.J. Lu, 1999. Humic acid-mediated oxidative damages to human erythrocytes: A possible mechanism leading to anemia in blackfoot disease. *Free Radic. Biol. Med.*, 27: 470-477.
- Erel, O.A., 2005. A new automated colorimetric method for measuring total oxidant status. *Clin. Biochem.*, 38: 1103-1111.
- Goth, L.A., 1991. Simple method for determination of serum catalase activity and revision of reference range. *Clin. Chim. Acta*, 196: 143-152.
- Ho, K.J., T.K. Liu, T.S. Huang and F.J. Lu, 2003. Humic acid mediates iron release from ferritin and promotes lipid peroxidation *in vitro*: A possible mechanism for humic acid-induced cytotoxicity. *Arch. Toxicol.*, 77: 100-109.
- Jansson, L.T., M.V. Perkkio, W.T. Willis, C.J. Refino and P.R. Dallman, 1985. Red cell superoxide dismutase is increased in iron deficiency anemia. *Acta Haematol.*, 74: 218-221.
- Khelifa, A. and J.P. Steghens, 2004. Plasma lipid hydroperoxides measurement by an automated xylenol orange method. *Anal. Biochem.*, 325: 58-163.
- Liang, H.J., C.L. Tsai and F.J. Lu, 1998. Oxidative stress induced by humic acid solvent extraction fraction in cultured rabbit articular chondrocytes. *J. Toxicol. Environ. Health*, 54: 477-489.
- Liang, H.J., C.L. Tsai, P.Q. Chen and F.J. Lu, 1999. Oxidative injury induced by synthetic humic acid polymer and monomer in cultured rabbit articular chondrocytes. *Life Sci.*, 65: 1163-1173.
- Ramachandran, M. and G.Y. Iyer, 1984. Erythrocyte membrane lipid peroxidation in iron deficiency anemia. *Experientia*, 40: 173-174.
- Rex, R.W., 1960. Electron paramagnetic resonance studies of stable free radicals in lignins and humic acids. *Nature*, 188: 1185-1186.
- Shermer, C.L., K.G. Maciorowski, C.A. Bailey, F.M. Byers and S. Ricke, 1998. Caecal metabolites and microbial populations in chickens consuming diets containing a mined humate compound. *J. Sci. Food Agric.*, 77: 479-486.