

Exploitation of Ethanol Extract of *Adenium obesum* Stem Bark as a Potent Organic Piscicide

¹Samson Enejo Abalaka, ¹Muhamad Yakassai Fatihu,
¹Najume Doguwar Gingiya Ibrahim and ²Suleiman Folorunsho Ambali
¹Department of Veterinary Pathology,
²Department of Veterinary Pharmacology and Toxicology,
University of Ahmadu Bello, Zaria, Nigeria

Abstract: The possible use of ethanol extract of *Adenium obesum* stem bark as a potent organic piscicide, against unwanted predatory animals and/or fish in fish ponds prior to the stocking of the desired fish was investigated via the exposure of the hardy air-breathing *Clarias gariepinus* to the extract over a 96 h exposure period. *Clarias gariepinus* of 265.50±4.03 g mean weight and 32.85±0.16 cm mean total length were exposed to 0.00, 6.25, 7.50, 8.20, 8.80 and 9.30 mg L⁻¹ of the extract in a static acute toxicity bioassay after performing a range finding test. The extract contained some toxicologically bio-active substances. Changes in the fish culture water quality were within acceptable limits. The exposed fish showed signs of restlessness and hyperactivity along with signs of respiratory and nervous compromise, as well as the observed mortality in some of the exposed fish whose appearance and intensity were concentrations and exposure period-dependent. However, no signs of toxicity were observed in the unexposed control fish. The median lethal concentration of the extract in the exposed fish over the 96 h exposure period was 7.15 mg L⁻¹. Therefore, at this or much higher concentrations, the extract will be a potent organic piscicide that can be used to eradicate unwanted predatory animals, especially fish from ponds prior to the stocking of desired fish.

Key words: *Adenium obesum*, *Clarias gariepinus*, toxicity, median lethal concentration, pond management, piscicidal plant

INTRODUCTION

There has been a global demand for fish within the last 30 years from 45 million metric ton (mt) in 1973 to about 91 million mt in 1997 (Delgado *et al.*, 2003) and this is further expected to rise to about 183 million mt by year 2030 (Ye, 1999). This is because fish has become one of the most important sources of available animal protein (Jan *et al.*, 2012) that may be the sole accessible and/or affordable sources of animal protein for poor households in urban and semi-urban areas (Bene and Heck, 2005). This is most important in developing countries where fish is comparatively cheaper to either beef or chicken or mutton which is even usually in short supply (Omoniyi, 1995; Amisah *et al.*, 2008). The increased global demand has led to increased fishing efforts in wild aquatic environments, especially in the developing countries. However, the global landings of wild-caught fish are thought to be nearing and/or exceeding their sustainable levels resulting in a worldwide resort to fish farming or aquaculture (Faturoti, 1999; Jourie, 2006; Omitoyin, 2007).

Aquaculture is the fastest growing food sector in the world, accounting for an estimated 43% of all fish consumed by humans globally (Allsopp *et al.*, 2008). However, aquaculture growth is still very slow in Africa as it is currently faced with numerous challenges (FAO, 2011; Peteri *et al.*, 1992; Masser *et al.*, 1999; Omitoyin, 2005). Notable amongst these challenges is the issues of predatory animals like tadpoles, frogs, leeches and unwanted fish species that competes with and feed on the fry and fingerlings of stocked fish (Olaifa *et al.*, 2008; Fafioye, 2012), including competition for available oxygen (Kumar, 1992). The use of synthetic compounds to clear out these predatory animals and/or fish has been a common practice amongst aquaculturists (Marking, 1992) but synthetic compounds cause heavy environmental pollution (Mondal *et al.*, 2007) with harmful residues in aquatic environments (Koesomadinata, 1980; Cagauan and Arce, 1992). That is why attention is now focused on the use of piscicidal plants or plants that are poisonous to fish in place of these synthetic compounds (Marston and Hostettmann, 1985; Dahiya *et al.*, 2000). This is because

piscicidal plants are easily bio-degraded, less severe and are comparatively safer toward humans and the environment (Kela *et al.*, 1989; Stalin *et al.*, 2008).

Adenium obesum is a known piscicidal plant (Adamu *et al.*, 2005; Oyen, 2008). The plant is distinct with swollen base, twisted branches and characteristic showy flowers which gives it the name Desert rose (McLaughlin and Garofalo, 2002; Bawden-Davies, 2010). The plant is slow growing but long lived (McLaughlin and Garofalo, 2002) and produces sticky clear or white latex (Rowley, 1983; Oyen, 2008). Although, the plant occurs throughout the Sahel region of Africa, spreading downward towards Central Africa and Northward into Arabia (Plaizier, 1980; Arbonnier, 2004), it is found worldwide where it is usually cultivated for ornamental purposes (Oyen, 2008; Hastuti *et al.*, 2009).

Even though, *Clarias gariepinus* has Pan-African distribution (De Graaf and Janssen, 1996), it is absent from the Maghreb, Upper and Lower Guinea and Cape Provinces (FAO, 2011). The fish is hardy (Olaifa *et al.*, 2003) and can tolerate harsh aquatic conditions (Hogendoorn, 1979; Eyo and Ezechie, 2004). Therefore, any piscicidal plant that kills this fish species might kill some of these unwanted predatory animals and/or fish before the stocking of the desired fish. This is in addition to the apparent dearth of information on the toxicity of *A. obesum* stem bark to exposed fish, especially the air-breathing *C. gariepinus* which has created the need for this study. The research aimed to ascertain the toxicity of the ethanol extract of *A. obesum* stem bark in the exposed adult *C. gariepinus*, as a possible tool for the eradication of unwanted animals and/or fish from ponds prior to the stocking of desired fish.

MATERIALS AND METHODS

Plant collection and preparation: *Adenium obesum* were collected from the open fields of Rurum town, Rano Local Government Area, Kano State, Nigeria within the months of January to April, 2011. These were authenticated at the Herbarium, Department of Biological Sciences, Ahmadu Bello University (ABU), Zaria, Nigeria with Voucher No. 1386. The barks were removed, sun-dried and pounded into powder ready for the extraction process.

A total of 3.95 kg of the pulverized *A. obesum* stem bark was macerated in 21 L of ethanol (96.0% vol. Sigma-Aldrich® Inc., St. Louis, MO 63178, USA) based on the method of Bentley (1977) and Ghani (1990) over a 72 h period. The filtrates were then concentrated to dryness in evaporation dishes by allowing the ethanol to evaporate at room temperature until constant weights were obtained as described by Abu-Dahab and Afifi (2007).

Phytochemical screening: A preliminary phytochemical screening of the ethanol extract of *A. obesum* stem bark for tannins, steroids and triterpens, glycosides, saponins, flavonoids and anthraquinones (Trease and Evans, 1983) and alkaloids (Harbone, 1998) was carried out.

Physicochemical analysis: Fish culture water quality parameters was monitored and recorded prior to fish introduction and daily throughout the 96 h exposure period. The modified method of Winkler-Azide (Lind, 1979; APHA, 1985) was used to determine the dissolved oxygen content of the fish culture water. Similarly, the pH, temperature, electrical conductivity and the total dissolved solids were determined using the Hanna Combo portable hand instrument (Hi 98129, Hanna Instruments, Mauritius).

Fish toxicity bioassay: Adult *Clarias gariepinus* were purchased from a commercial catfish farm (Fannasson Investments Limited, Kano, Nigeria). The fish were authenticated at the Fishery Section, Department of Biological Sciences, ABU, Zaria, Nigeria. The fish were acclimatized for 21 days before the start of the experiment under natural day and night photo-periods. Pond water was changed completely once in every 3 days. The experimental fish were fed to satiation twice daily with 6 mm Coppens® fish feed for aquaculture (Coppens International bv., 5700 AM Helmond, Holland). The fish were observed for disease conditions and mortality throughout the acclimatization period as recommended in the OECD (1992) guideline No. 203. The exposed fish were not fed 24 h prior to and during the 96 h exposure period in order to prevent interference from stomach contents and wastes in the fish culture water (Smith *et al.*, 2007; Olufayo, 2009). The exposed and the unexposed (control) fish were monitored with prompt recordings of all observations at 24 (0, 3, 6 and 12 h), 48, 72 and 96 h during the exposure period.

Clarias gariepinus (265.50±4.03 g mean weight and 32.85±0.16 cm mean total length) were used for the static acute toxicity bioassay over the 96 h exposure period (OECD, 1992) in triplicates. This was after a preliminary concentration range finding test was performed (Fafioye, 2001) to determined 5 different ethanol extract concentrations of *A. obesum* stem bark that was used (0.0, 6.25, 7.50, 8.20, 8.80 and 9.30 mg L⁻¹). The median lethal concentration (LC₅₀) of the ethanol extract of *A. obesum* stem bark in the exposed *C. gariepinus* over the 96 h exposure period was established based on the Probit and logit analysis method (Finney, 1971). Fish mortality represented extract toxicity. The exposed fish were considered dead if there were no visible opercula movements or no visible reactions when touched (OECD, 1992).

Statistical analysis: Data were expressed as mean (\pm SEM) and subjected to ANOVA and Tukey's multiple comparison test for statistical significance at $p < 0.05$ using GraphPad software programme (GraphPad Prism, version 4.0, San Diego, California, USA).

RESULTS AND DISCUSSION

The ethanol extract of *A. obesum* stem bark had no significant ($p > 0.05$) effects on the physicochemical parameters of the fish culture water over the 96 h exposure period as shown in Table 1. Although, the observed changes were not concentration dependent, the extract caused decreases in the temperature, pH and the dissolved oxygen but increases in total dissolved solids and the electrical conductivity of the fish culture water with increasing extract concentrations compared to the respective controls. The preliminary phytochemical screening of the extract revealed the presence of tannins, alkaloids, saponins, steroids and triterpens, glycosides, flavonoids, cardiac glycosides but no anthraquinones as shown in Table 2.

The exposed fish showed signs of toxicity in the forms of restlessness, erratic movements and repeated attempts to jump out of the fish culture water in addition to excessive mucus secretion. Frequent opercula movements and air gulping with snouts exposures were also exhibited by the exposed fish. Other signs of toxicity were those of increasing states of motionlessness and the adoption of different postures with sudden darts, swirling movements and loss of balance, including the observed mortality in some of the exposed fish. These signs were directly related to the concentrations of the extract. None of these signs of toxicity were observed in the unexposed control fish. The LC_{50} of the ethanol extract of *A. obesum* stem bark in the exposed *C. gariepinus* over the 96 h exposure period was established to be 7.15 mgL^{-1} with a 96.95% degree of fits (R^2) as shown in Fig. 1. Mean mortality increased significantly ($p < 0.05$) with increased extract concentrations.

The observed changes in the physicochemical parameters of the fish culture water were all within the acceptable limits for the growth and survival of *C. gariepinus* (Viveen *et al.*, 1985; Peteri *et al.*, 1992;

Hogendoorn, 1979; Chapman, 2000). The preliminary phytochemical screening of the extract revealed the presence of some toxicologically active bio-active substances (Da Rocha *et al.*, 2001; Bent and Ko, 2004). This is because saponins which are directly absorbed through the gills, haemolyses erythrocytes (Elpel, 2000), due to their ability to lower the surface tension between the aqueous and the lipid phases of the membranes of erythrocytes resulting in the emulsification of their lipid components with Na^+ and water influx and K^+ efflux until they ruptures to release their haemoglobin contents (Hostettmann and Marston, 1995). The toxicity might have also been through the impairment of oxygen consumption in the exposed fish as saponins are also reported to lower the surface tension of reconstituted extracts with the formation of colloidal substances within them (Geidam *et al.*, 2007; Armstrong, 2008). This is in addition to the fact that saponins are also known to cause structural damages in the respiratory and intestinal epithelia of exposed fish (Roy *et al.*, 1990).

The toxicity of tannins might be through their protein coagulative property (Davidson, 1979), especially gill epithelia, thereby causing respiratory failure or asphyxiation in the exposed fish (Idris, 2012). Tannins are also known to bind and inhibit endogenous proteins like the digestive enzymes (Kumar and Singh, 1984) in addition to causing intestinal damage and interference with iron absorption (Butler, 1989). Impaired oxygen consumption via the inhibition of oxidative phosphorylation involving the blockage of mitochondrial enzyme, NADH ubiquinone reductase has been ascribed to alkaloids in exposed fish (Bocek, 1984; Tiwari and Singh, 2003). This agreed with the reported inhibition of pyridine-linked substrates oxidation suggestive of interference with oxidative phosphorylation in ticks exposed to the aqueous extract of *A. obesum* stem bark (Mgbojikwe, 2000). The observed signs of toxicity might have also been due to excessive inhibition of the Sodium-Potassium pump (Na^+ -ATPase pump) caused by the cardiac glycosides thereby causing a variety of severe arrhythmias resulting in blocked cardiac activity, decreased cardiac output and death in some of the exposed fish (Knight and Walter, 2001).

Table 1: The physicochemical parameters of the fish culture water for the 96 h exposure period


Extract conc. (mgL^{-1})	Physicochemical parameters				
	Temperature ($^{\circ}\text{C}$)	pH	Dissolved oxygen (mg L^{-1})	Total dissolved solids (ppm)	Electrical conductivity (μScm^{-1})
9.30	24.34 \pm 0.14	7.20 \pm 0.06	3.15 \pm 0.80	348.90 \pm 6.08	697.40 \pm 11.61
8.80	24.47 \pm 0.13	7.16 \pm 0.04	3.96 \pm 0.67	352.80 \pm 6.77	706.10 \pm 13.33
8.20	24.57 \pm 0.15	7.14 \pm 0.04	3.59 \pm 0.62	349.80 \pm 7.39	697.80 \pm 12.09
7.50	24.57 \pm 0.17	7.16 \pm 0.04	3.81 \pm 0.65	347.90 \pm 5.70	699.70 \pm 11.74
6.25	24.63 \pm 0.16	7.17 \pm 0.04	4.22 \pm 0.47	345.60 \pm 4.81	688.70 \pm 9.850
0.00 (Control)	24.65 \pm 0.18	7.22 \pm 0.04	4.56 \pm 0.63	341.80 \pm 7.61	680.30 \pm 14.70

Conc. = Concentration

Table 2: Preliminary phytochemical screening of the ethanol extract of *Adenium obesum* stem bark

Parameters	Tests	Extract solution	
		Observations	Indication
Saponins	Frothing	Honey comb forth	+
Tannins	Ferric chloride	Green precipitate	+
Alkaloids	Meyer	Creamy precipitate	+
Glycosides	Fehling's solution	Brick red precipitate	+
Cardiac glycoside	Kella-Killiant	Purple brown inter-phase	+
Flavonoids	Sodium hydroxide	Yellow colour	+
Carbohydrates	Molisch	Purple precipitate	+
Steroids and triterpenes	Lieberman-Buchard	Reddish brown color	+
Anthraquinones	Borntrager	Transparent	-

+ = Present; - = Absent

Fig. 1: Log of concentration of the ethanol extract of *Adenium obesum* stem bark and the probit values in *Clarias gariepinus* over the 96 h exposure period

The initial agitations and restlessness characterized by erratic movements and repeated attempts of the fish to jump out of the culture water were manoeuvres to escape from the toxic fish culture water. This is because fish responds to perceived noxious contamination by avoiding the area containing the chemical (Kane *et al.*, 2005) in attempts to prevent the continuous absorption of the toxicant present within the fish culture water. Similar hyperactivity and uncoordinated movements were reported in ticks exposed to the aqueous extract of *A. obesum* stem bark (Mgbojikwe, 2000).

The increased opercula movements were due to decreased efficiency of oxygen uptake and transport and/or increased metabolic rate (Thomas and Rice, 1975). The observed frequent air gulping and snout exposures were attempts to free gill respiration for aerial respiration in order to limit continuous gill contact with the toxic aquatic environment. This is an aquatic surface respiration in response to low oxygen that predisposes affected to easy catch and/or predation (Reebs, 2009). The observed nervous disorders might be due to brain cytochrome c oxidase inhibition leading to cytotoxic

hypoxia with resultant changes in the electrical activity of the brain associated with the maintenance of equilibrium (Rao and Rao, 1987). It might have also been due to acetylcholinesterase inhibition leading to acetylcholine accumulation at synaptic junctions with subsequent hyperstimulation (Halappa and David, 2009). This is because Mgbojikwe (2000), reported acetylcholinesterase activity inhibition in ticks exposed to the aqueous extract of *A. obesum* stem bark.

The presence of mortality in some of the exposed fish was indicative of the toxicity of the extract in the exposed fish as reinforced by the established LC₅₀ value of 7.15 mgL⁻¹. The decrease in mean mortality with increase exposure period showed the acute nature of the toxicity. The subsequent decline might be due to the bio-transformation of the toxic bio-active constituents of the extract over time (Kela *et al.*, 1989; Fafioye, 2005), especially as the extract concentrations were not maintained daily throughout the exposure period. The ethanol extract of *A. obesum* stem bark was highly toxic to the exposed *C. gariepinus* because the lower the LC₅₀, the higher the toxicity or sensitivity to test organisms and vice versa (Buikema *et al.*, 1982) meaning that at higher LC₅₀, greater concentrations are required to the produce 50% mortality in test organisms and vice versa (Hilmy *et al.*, 1985; Eisler and Gardner, 1973). The high toxicity might be the reason behind the global use of the powdered roots or a decoction of the bark and the leaves of the plant to poison fish (Arbonnier, 2004; Adamu *et al.*, 2005; Oyen, 2008; Bawden-Davis, 2010).

CONCLUSION

The ethanol extract of *A. obesum* stem bark was highly toxic to the exposed *C. gariepinus*. The utilization of the LC₅₀ or much higher concentrations of the extract will make it a potent organic piscicide that can be used to eradicate predatory animals, especially unwanted fish from ponds prior to the stocking of the desired fish.

ACKNOWLEDGEMENTS

The immense contributions of Mr. A. Umar and Mr. S.M. Meslam are well appreciated.

REFERENCES

- APHA, 1985. Standard Method for the Examination of Water and Wastewater. 15th Edn., American Public Health Association, Washington, DC., USA., pp: 413-426.
- Abu-Dahab, R. and F. Afifi, 2007. Antiproliferative activity of selected medicinal plants of Jordan against a breast adenocarcinoma cell line (MCF7). *Scient. Pharm.*, 75: 121-136.

- Adamu, H.M., O.J. Abayeh, M.O. Agho, A.L. Abdullahi, A. Uba, H.U. Dukku and B.M. Wufem, 2005. An ethnobotanical survey of Bauchi State herbal plants and their antimicrobial activity. *J. Ethnopharmacol.*, 99: 1-4.
- Allsopp, M., P. Johnston and D. Santillo, 2008. Challenging the aquaculture industry on sustainability: Technical overview. Greenpeace Research Laboratories Technical Note 01/2008. http://www.greenpeace.to/publications/Aquaculture_Report_Technical.pdf.
- Amisah, S., D. Adjei-Boateng and D.D. Afianu, 2008. Effects of bamboo substrate and supplementary feed on growth and production of the African catfish, *Clarias gariepinus*. *J. Applied Sci. Environ. Manage.*, 12: 25-28.
- Arbonnier, M., 2004. Trees, Shrubs and Lianas of West African Dry Zones. CIRAD, MARGRAF and MNHN, Wageningen, The Netherlands, ISBN-13: 9782876145795, pp: 161.
- Armstrong, W.P., 2008. Major types of chemical compounds in plants and animals part II. Phenolic compounds, glycosides and alkaloids. <http://waynesword.palomar.edu/chemid2.htm>.
- Bawden-Davies, J., 2010. The *Adenium* species. http://www.how.com/about_6721472_adenium-species.html.
- Bene, C. and S. Heck, 2005. Fish and food security in Africa. *NAGA WorldFish Center Q.*, 28: 8-13.
- Bent, S. and R. Ko, 2004. Commonly used herbal medicines in the United States: A review. *Am. J. Med.*, 116: 478-485.
- Bentley, A.O., 1977. *Bentley's Textbook of Pharmaceutics*. 8th Edn., Bailliere Tindall, London, ISBN: 10: 0702003913, pp: 177-180.
- Bocek, B.R., 1984. Ethnobotany of Costanoan Indians, California, based on collections by John P. Harrington. *Econ. Bot.*, 38: 240-255.
- Buikema Jr., A.L., B.R. Niederlehner and J. Cairns Jr., 1982. Biological monitoring part IV: Toxicity testing. *Water Res.*, 16: 239-262.
- Butler, L.G., 1989. Effects of Condensed Tannins on Animal Nutrition. In: *Chemistry and Significance of Condensed Tannins*, Hemingway, R.W. and J.J. Karchesy (Eds.). Plenum Press, New York, pp: 391-402.
- Cagauan, A.G. and R.G. Arce, 1992. Overview of Pesticides Use in Rice-Fish Farming in South East Asia. In: *Rice-Fish Research Development in Asia*, Dela Cruz, C.R., C. Lightfoot, B.A. Coasta-Pierce, V.R. Carangal and M.P. Bimbao (Eds.). International Centre for Living Aquatic Resources Management (ICLARM), Manila, Philippines, ISBN-13: 9789711022884, pp: 217-233.
- Chapman, F.A., 2000. Farm-Raised Channel Catfish. Department of Fisheries and Aquatic Sciences, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville.
- Da Rocha, A.B., R.M. Lopes and G. Schwartzmann, 2001. Natural products in anticancer therapy. *Curr. Opin. Pharmacol.*, 1: 364-369.
- Dahiya, S., A. Kuar and N. Jain, 2000. Studies on diflouridation of water with Eichornia plant. *J. Environ. Biol.*, 21: 337-340.
- Davidson, G., 1979. *Beverages, Herbs and Species, Human Nutrition and Dietetics*. 7th Edn., Churchill Livingstone, London, UK., pp: 200-201.
- De Graaf, G. and J.A.L. Janssen, 1996. Artificial reproduction and pond rearing of the African catfish *Clarias gariepinus* in Sub Saharan Africa. FAO Fisheries Technical Paper 362.
- Delgado, C.L., N. Wada, M.W. Rosegrant, S. Meijer and M. Ahmed, 2003. Fish to 2020: Supply and demand in changing global markets. WorldFish Center Technical Report 62, International Food Policy Research Institute, Washington, DC. and WorldFish Centre, Penang, Malaysia.
- Eisler, R. and G.R. Gardner, 1973. Acute toxicology to an estuarine teleost of mixture of cadmium, copper and zinc salts. *J. Fish. Biol.*, 5: 131-142.
- Elpel, T.J., 2000. *Botany in a Day: Thomas J. Elpel's Herbal Field Guide to Plant Families*. 4th Edn., HOPS Press, Pony, Montana, USA., ISBN-13: 9781892784070, Pages: 196.
- Eyo, J.E. and C.U. Ezechie, 2004. The effects of rubber (*Havea brasiliensis*) seed meal based diets on diets acceptability and growth performance of *Heterobranchus bidorsalis* (male) x *Clarias gariepinus* (Female) hybrid. *J. Sustainable Trop. Agric. Res.*, 10: 20-25.
- FAO, 2011. Cultured aquatic species information programme: *Clarias gariepinus* (Burchell, 1822). Fisheries and Aquaculture Department, Food and Agriculture Organization of the United Nations.
- Fafioye, O.O., 2001. Lethal and sub-lethal effect of extracts of *Parkia biglobosa* and *Rahpia vinifera* on some fresh water fauna. Ph.D. Thesis, University of Ibadan, Nigeria.
- Fafioye, O.O., 2005. Plants with piscicidal activities in SouthWestern Nigeria. *Turk. J. Fish. Aqua. Sci.*, 5: 91-97.
- Fafioye, O.O., 2012. Acute and sub-acute toxicities of five plant extracts on white tilapia, *Oreochromis niloticus* (Trewavas). *Int. Res. J. Agric. Sci. Soil Sci.*, 2: 525-530.

- Faturoti, E.O., 1999. Fisheries potentials and investments opportunities in Nigeria. Proceedings of the Fisheries Society of Nigeria (FISON) Investors Workshop, April 22, 1999, National Institute for Oceanography and Marine Research (NIOMR), Lagos, Nigeria -.
- Finney, D.J., 1971. Probit Analysis. Cambridge University Press, London, UK., pp: 123-125.
- Geidam, M.A., E. Dauda and H.G. Hamza, 2007. Effects of aqueous stem-bark extract of *Momordica balsamina* Linn on some serum enzymes in normal and ethanol fed rats. J. Boil. Sci., 7: 397-400.
- Ghani, A., 1990. Introduction to Pharmacognosy. 1st Edn., Ahmadu Bello University Press Ltd., Zaria, Nigeria, pp: 109-145.
- Halappa, R. and M. David, 2009. Behavioural responses of the freshwater fish, *Cyprinus carpio* (Linnaeus) following sublethal exposure to chlorpyrifos. Turk. J. Fish. Aquat. Sci., 9: 233-238.
- Harbone, J.B., 1998. Phytochemical Methods: A Guide to Modern Techniques of plant 3rd Edn., Chapman and Hill, London, pp: 279-280.
- Hastuti, D., Suranto and P. Setyono, 2009. Variation of morphology, karyotype and protein band pattern of adenium (*Adenium obesum*) varieties. Nusantara Biosci., 1: 78-83.
- Hilmy, A.M., M.B. Shabana and A.Y. Daabees, 1985. Bioaccumulation of cadmium: Toxicity in *Mugil cephalus*. Comp. Biochem. Physiol. C Comp. Pharmacol., 81: 139-144.
- Hogendoorn, H., 1979. Controlled propagation of Africa catfish, *Clarias lazera* (C & V).1. Reproductive biology and field experiment. Aquaculture, 17: 323-333.
- Hostettmann, K. and A. Marston, 1995. Saponins: Chemistry and Pharmacology of Natural Products. Cambridge University Press, Melbourne, Australia, ISBN-13: 9780521329705, Pages: 548.
- Idris, U.J., 2012. Toxicity of the methanol extracts of *Jatropha gassypifolia* L. (Bellyache) and *Trophosia vogelii* Hook (Fish bean) on African catfish, *Clarias gariepinus* (Burchell). Ph.D. Thesis, Ahmadu Bello University, Zaria, Nigeria.
- Jan, U., M. Shah, T. Manzoor and S.A. Gaine, 2012. Seasonal and monthly variations of protein content in the muscle of fish *Schizothorax esocinus*. Recent Res. Sci. Technol., 4: 5-7.
- Jourie, J.J., 2006. A practical investigation into catfish (*Clarias gariepinus*) farming in the Vaalharts irrigation scheme. Master's Thesis, University of the Free State, South Africa.
- Kane, A.S., J.D. Salierno and S.K. Brewer, 2005. Fish Models in Behavioural Toxicology: Automated Techniques, Updates and Perspectives. In: Methods in Aquatic Toxicology, Ostrander, G.K. (Ed.). Vol. 2, Taylor and Francis, Boca Raton, FL., USA., ISBN-13: 9781566706643, pp: 550-590.
- Kela, S.L., R.A. Ogunsusi, V.C. Ogbogu and N. Nwude, 1989. Screening of some Nigerian plants for molluscicidal activity. Revue d'Elevage et de Medicine Veterinaire des Pays Tropicaux, 42: 195-202.
- Knight, A.P. and R.G. Walter, 2001. A Guide to Plant Poisoning of Animals in North America. Teton NewMedia, Jackson, MI., USA., ISBN-13: 9781893441118, pp: 57-75.
- Koesomadinata, S., 1980. Pesticide as a Major Constraint in Integrated Agriculture-Aquaculture Farming System. In: Integrated Agriculture-Aquaculture Farming System, Pullin, R.S.V. and Z.H. Shehadeh (Eds.). International Centre for Living Aquatic Resources Management (ICLARM), Manila, Philippines, pp: 45-51.
- Kumar, R. and M. Singh, 1984. Tannins: Their adverse role in ruminant nutrition. J. Agric. Food Chem., 32: 447-453.
- Kumar, D., 1992. Fish Culture in Undrainable Ponds: A Manual for Extension. Food and Agriculture Organization of the United Nations, Rome, Italy, ISBN-13: 9789251031391, Pages: 239.
- Lind, O.T., 1979. A Handbook of Limnological Methods. 1st Edn., C.V. Mosby Co., St. Louis, USA., Pages: 199.
- Marking, L.L., 1992. Evaluation of toxicants for the control of carp and other nuisance fishes. Fisheries, 17: 6-13.
- Marston, A. and K. Hostettmann, 1985. Plant molluscicides. Phytochemistry, 24: 639-652.
- Masser, M.P., D. Steinbach and B. Higginbotham, 1999. Catfish pond for recreation. Texas Agriculture Extension Service (B-1319: 6/99), The Texas A&M University System, USA., pp: 1-12.
- McLaughlin, J. and J. Garofalo, 2002. Desert rose (*Adenium obesum*). Fact-Sheet No. 17, Miami-Dade County/University of Florida Cooperation Extension Services, USA.
- Mgbojike, L.O., 2000. Acaricidal properties of the aqueous stem bark extract of *Adenium obesum*. Ph.D. Thesis, University of Jos, Jos, Nigeria.
- Mondal, D., S. Barat and M.K. Mukhopadhyay, 2007. Toxicity of neem pesticides on a fresh water loach, *Lepidocephalichthys guntea* (Hamilton Buchanan) of Darjeeling district in West Bengal. J. Environ. Biol., 28: 119-122.

- OECD, 1992. OECD guideline for testing of chemicals: Fish, acute toxicity test. No. 203, Organisation for Economic Cooperation and Development. <http://www.oecd.org/chemicalsafety/risk-assessment/1948241.pdf>.
- Olaifa, F.E., A.K. Olaifa and O.O. Lewis, 2003. Toxic stress of lead on *Clarias gariepinus* (African catfish) fingerlings. Afr. J. Biomed. Res., 6: 101-104.
- Olaifa, F.E., O.O. Oyetooyan and R.A. Hamzat, 2008. Acute toxicity of ethanol extracts of cocoa bean shell on *Sarotherodon galilaeus* juveniles. J. Fish. Int., 3: 56-60.
- Olufayo, M.O., 2009. Haematological characteristics of *Clarias gariepinus* (Burchell, 1822) juveniles exposed to *Derris elliptica* root powder. Afr. J. Food Agric. Nutr. Dev., 9: 920-933.
- Omitoyin, O.B., 2005. Problems and prospects of fish feed production in Nigeria. Proceedings of the USAID Aquaculture Marketing Stakeholder Forum, December 13, 2005, University of Ibadan Conference Centre, Ibadan, Nigeria, pp: 3-.
- Omitoyin, O.B., 2007. Introduction to Fish Farming in Nigeria. Ibadan University Press, Nigeria, ISBN-13: 9789781214271, pp: 1-50.
- Omoniyi, I.T., 1995. Hybridization in *Oreochromis niloticus* (Trewaras) and *Sarotherodon galilaeus* (Linnaeus): An assessment of morphometrics and the response of F1 fry to formulated diets. Ph.D. Thesis, University of Ibadan, Ibadan, Nigeria.
- Oyen, L.P.A., 2008. *Adenium obesum* (Forssk.) Roem. & Schult. In: Plant Resources of Tropical Africa 11 (1): Medicinal Plant 1, Schmelzer, G.H., A. Gurib-Fakim, R.R.J. Arroyo, C.H. Bosch, A. de Ruijter and M.S.J. Simmonds (Eds.). PROTAFoundation/Backhuys Publishers/CTA, Wageningen, Netherlands, ISBN-1 3: 9789057822049, pp: 6-49.
- Peteri, A., S. Nandi and S.N. Chowdhury, 1992. Manual on seed production of African catfish (*Clarias gariepinus*). BGD/87/045/92/22, Field Document (Draft), Institutional Strengthening in the Fisheries Sector, Bangladesh.
- Plaizier, A.C., 1980. A revision of *Adenium* Roem and Schult. and of *Diplorhynchus* Welw. ex Fic. & Hiern (Apocynaceae). Publication No. 80-12, Wageningen, Netherlands, pp: 1-40.
- Rao, K.R.S.S. and J.C.S. Rao, 1987. Independent and combined action of carbaryl and phenthoate on snakehead, *Channa punctatus* (Bloch). Curr. Sci., 56: 331-332.
- Reebs, S.G., 2009. Oxygen and fish behaviour. Universite de Moncton, Canada, pp: 1-13.
- Rowley, G.D., 1983. The *Adenium* and *Pachypodium* Handbook. British Cactus and Succulent Society, London, UK., ISBN-13: 9780902099074, Pages: 95.
- Roy, P.K., J.D. Munshi and H.M. Dutta, 1990. Effect of saponin extracts on morphology and respiratory physiology of an air breathing fish, *Heteropneustes fossilis* (Bloch). J. Freshwater Biol., 2: 135-145.
- Smith, C.J., B.J. Shaw and R.D. Handy, 2007. Toxicity of single walled carbon nanotubes to rainbow trout (*Oncorhynchus mykiss*): Respiratory toxicity, organ pathologies and other physiological effects. Aquat. Toxicol., 82: 94-109.
- Stalin, S.I., S. Kiruba and S.S.M. Das, 2008. A comparative study on the toxicity of a synthetic pyrethroid, deltamethrin and a neem based pesticide, azadirachtin to *Poecilia reticulata* Peters 1859 (Cyprinodontiformes: Poeciliidae). Turk. J. Fish. Aquat. Sci., 8: 1-5.
- Thomas, R.E. and S.D. Rice, 1975. Increased opercular rates of pink salmon (*Oncorhynchus gorbuscha*) fry after exposure to the water-soluble fraction of prudhoe bay crude oil. J. Fish. Res. Board Can., 32: 2221-2224.
- Tiwari, S. and A. Singh, 2003. Control of common freshwater predatory fish, *Channa punctatus*, through *Nerium indicum* leaf extracts. Chemosphere, 53: 865-875.
- Trease, G.E. and W.C. Evans, 1983. Pharmacognosy. 12th Edn., Bailliere Tindall, Philadelphia, USA., pp: 338-628.
- Viveen, W.J.A.R., C.J.J. Rither, P.G.W.J. Van Oordt, J.A.L. Janssen and E.A. Huisman, 1985. Practical manual for the culture of the African catfish (*Clarias gariepinus*). Directorate General for International Technical Corporation, Hague, Netherlands, pp: 93.
- Ye, Y., 1999. Historical consumption and future demand for fish and fishery products: Exploratory calculations for the years 2015/2030. FAO Fish., Circ., Rome, Italy, Vol. 946. pp: 2.