

Contaminant Source as Factor of Soil Heavy Metal Toxicity and Bioavailability to Plants

Olayinka I. Nwachukwu

Department of Soil Science and Agroclimatology,

Michael Okpara Univeristy of Agriculture, Umudike, P.M.B. 7267, Umuahia, Nigeria

Abstract: Mine waste and sewage sludge treated soil were used as contaminants in a greenhouse pot experiment and the effect of the contaminants on ryegrass growth, soil metal levels and plant metal uptake was evaluated over a period of 16 weeks. Soils were amended with bone meal, compost, peat, coir or lime phosphate and amendments led to immobilization of Pb, Cu and Zn. Biomass yield was significantly reduced in contaminated soil, with greater reduction in mine waste contaminated soil. EDTA extractable soil metal and metal uptake by ryegrass was significantly higher in mine waste contaminated soil than in sewage sludge treated soil.

Key words: Heavy metal, source, mine waste, sewage sludge, bioavailability

INTRODUCTION

Soil physicochemical properties are adversely affected by high concentrations of heavy metals, rendering contaminated soils unsuitable for crop production (Hernandez-Allica *et al.*, 2007; Udom *et al.*, 2004). It has been documented that soils polluted by industrial effluents or wastes tend to exhibit a reduced porosity, have poor hydraulic conductivity and increased acidity (Hernandez-Allica *et al.*, 2007). The source of metals in soil may be due to the parent material or may be anthropogenic. Metals can also be transported from the soil into the groundwater and crops growing on contaminated soil (Sharma *et al.*, 2008).

Two devastating sources of heavy metal contamination in modern times is from industrial activities linked to mining (ADB, 2001) from the long-term use of sewage sludge for agricultural purposes (Fuentes *et al.*, 2006). Metals in the soil environment may exist as free metal ions and soluble metal complexes in the soil solution, occupy exchange sites, precipitated as insoluble compounds, be bound to inorganic or organic soil fractions, or occur as metals in the structure of silicate minerals (Basta *et al.*, 2005; Roberts, 2003). The bioavailable fraction is composed of free ions and those bound on exchangeable sites, while those bound to organic matter can also be rendered available on decomposition (Yin *et al.*, 2002). The mobility of trace metals, their bioavailability and related ecotoxicity to plants, depend strongly on their specific chemical forms or ways of binding (Basta *et al.*, 2005) and also on the nature of soil or material which serves as carrier of the contaminant material (Voutsas *et al.*, 1996).

Sewage sludge has been used successfully as a source of adding nutrients in the soil (Poykio *et al.*, 2007), at the same time it increases organic matter content; raises soil pH and thus, render the metals less soluble/bioavailable. However, depending on their characteristics the long-term use of sludge can cause heavy metal accumulation in soils (Alvarenga *et al.*, 2008; Fuentes *et al.*, 2006). Remediation of contaminated soil is possible by amending such soils with agricultural materials, as soil metals are adsorbed onto the amendment materials, immobilising such metals and thus, reducing their soil and plant availability (Abbaspour *et al.*, 2008; Nwachukwu and Pulford, 2008). The objective of the study was to evaluate the effect of contaminant source on soil and plant metal concentrations.

MATERIALS AND METHODS

Mine waste from Tyndrum and sewage sludge treated soil from Stoke Bardolph, both in Scotland, were used as contaminants in a greenhouse pot experiment in 2004. The effects of the contaminants on ryegrass growth and uptake of Pb, Cu and Zn was evaluated over a period of 16 weeks. Three sets of pot experiments were set up concurrently, using sharp sand as the major component of the growth medium. Mine waste and sewage sludge treated soil were manually added at the rates of 2.5 and 25%, respectively, while amendments bone meal, general purpose compost, peat, coir and lime phosphate were applied as 10% of soil weight. Each experiment was set up as a 5×2×4 CRD. Total metal analysis in contaminants and amendments was by aqua regia digestion, extractable soil metal was determined using EDTA, while leaf metal was

Table 1: Chemical properties of contaminants and amendments before crop growth (Nwachukwu, 2007; Nwachukwu and Pulford, 2008)

Material	pH	OM (%)	CEC cmol K kg ⁻¹	Pb (ug g ⁻¹)	Cu (ug g ⁻¹)	Zn (ug g ⁻¹)
Bone meal	6.9	32	19	61	23	188
Compost	6.4	77	62	30	23	49
Peat	3.6	95	197	23	12	22
Coir	5.6	93	120	28	33	56
Phosphate	2.8	9	33	43	33	252
Mine waste	5.9	18	8	65593	2947	25083
Sewage	6.9	29	60	1024	895	2302

OM = Organic Matter; pH and metal values (n = 10); CEC and OM% (n=5)

determined by hot nitric acid digestion. Initial metal concentrations in contaminated soils were above normal levels as stated by Ross and Kaye (1994). All metals were read on the AAS.

Experiment I: Sand was amended with the 5 materials and left to incubate for 5 days, following which each pot was sown with approximately 1 g of ryegrass seed (*Lolium perenne*) and harvested twice, first after 8 weeks of growth and again after a further 8 weeks of growth.

Experiment II: Sand was first manually contaminated with mine waste or sewage, amended with bone meal, compost, peat, coir and lime phosphate and incubated for 16 weeks.

Experiment III: Sand was manually contaminated with mine waste or sewage, amended with bone meal, compost, peat, coir and lime phosphate and incubated for 5 days before planting with ryegrass.

Total Pb, Cu and Zn concentrations were much higher in mine waste than the sewage sludge, while metal concentrations in the amendments were negligible (Table 1).

All statistical analyses were carried out with the programme MINITAB version for windows, analysed using a general linear model of ANOVA. Where, F-test was significant, Tukey's test at (p<0.05) was used to determine LSD for the comparison of means.

RESULTS AND DISCUSSION

Buffering capacity of contaminants: The response of the different agricultural materials to pH changes was evaluated by subjecting them to increasing amounts of HCl or NaOH. Mine waste was readily subject to changes in pH at the slightest increase in amounts of acid or alkali added, but the sewage treated soil was resistant to pH changes in the presence of either acid or alkali (Fig. 1). This showed that the inorganic mine waste had a very low buffering capacity, while sewage treated soil (organic) on the other hand, had a high buffering capacity, typical of organic materials (Fig. 1).




Fig. 1: Buffering capacity of mine waste and sewage soil contaminants ryegrass biomass in contaminated soil

Ryegrass biomass was significantly higher in the sewage sludge treated soil had better yield when compared to mine waste contaminated soil (Fig. 2), which may be explained by the fact that sewage would be nutrients to the soil. Metal induced leaf chlorosis was observed in plants growing in both contaminated soils, but significantly greater in mine waste contaminated soil (Fig. 2). Addition of amendments led to a significant increase in biomass yield in both the mine waste and sewage spiked soil when compared to that of the non amended soil.

Metal uptake of ryegrass from amendments and contaminants only (experiment I): In comparing, the relative effect of metal present in amendments (experiment I) and that in the contaminants to the total metal taken up into ryegrass leaves, it was observed that the contribution of metals from the amendments to overall metal uptake from contaminants was negligible (Fig. 3). The very low concentrations of Pb, Cu and Zn in the amendments was however, available to the ryegrass plants, confirming that it is impossible to completely exclude metal uptake by plants, even when such plants are growing in non contaminated soil (Friesl *et al.*, 2006).

Metal uptake by ryegrass in contaminated soil: After 8 weeks of plant growth, metal uptake in mine waste spiked soil was significantly higher than in the sewage contaminated one (Fig. 4), taking initial metal concentrations into consideration.




Fig. 2: Ryegrass in non-amended metal 'spiked' soil after 4 weeks of growth (G-0 = mine waste contaminated soil + no amendment added; SB-0 = sewage treated soil + no amendment added)




Fig. 3: Comparative contributions of amendments and contaminants to the uptake of Pb, Cu and Zn by ryegrass after 8 weeks of growth

For instance, Zn concentrations in both the mine waste treated and sewage sludge treated 'made soils' was




Fig. 4: Total Pb, Cu and Zn in ryegrass leaves after 8 weeks of growth in contaminated soil (error bars represent \pm standard deviation of 4 replications)

approximately the same ($25000 \mu\text{g g}^{-1}$) having used mine waste at 1% and sewage soil at 10%, yet plant uptake of Zn was much higher in the mine waste treated soil (Fig. 4). Zn uptake from mine waste ranged from $200\text{--}550 \mu\text{g g}^{-1}$, but only from $120\text{--}180 \mu\text{g g}^{-1}$ from sewage sludge. This indicates that Zn was more readily available from the inorganic mine waste and less from the organic sewage treated soil within the first 8 weeks. Amendments in the mine waste spiked soil led generally to a decrease in uptake of Pb and Zn when compared to non amended soil, while in the sewage treated soil, the impact of amendments was significant on Pb and Cu uptake (Fig. 4).

At 16 weeks, metal uptake from mine waste remained significantly higher than that from sewage sludge however, metal uptake from both contaminants had increased above that observed at 8 weeks, indicating greater solubility of the metals over time (Fig. 5).

This confirms the findings of Karaca (2004) and Clemente *et al.* (2006) and it is thought that this may be due to the re-release of metals which complexed with organic matter during the earlier phase of an experiment (Madrid *et al.*, 2006).

Bioavailability of metals from mine waste and sewage sludge: Using a bioavailability index or factor which was essentially a ratio of plant metal concentration to soil metal, Pb and Zn from mine waste contamination was consistently more bioavailable than that from the sewage contamination in non amended soils. There was no




Fig. 5: Total Pb, Cu and Zn in ryegrass leaves after 16 weeks of growth in contaminated soil (error bars represent \pm standard deviation of 4 replications)

Table 2: Effect of contaminant source on Pb, Cu and Zn bioavailability

Contaminant	Bioavailability factor (BF)		
	Pb	Cu	Zn
Mine waste	19.0 \pm 3.3b	34.1 \pm 5.2ab	56.7 \pm 5.5b
Sewage sludge	6.7 \pm 0.8a	27.6 \pm 2.2a	29.4 \pm 5.5a
LSD	2.6	9.7	7.9

Means \pm S.D. (n = 4); Values within a column followed by the same letter are not significantly different at $p < 0.05$

significant difference in Cu bioavailability from either contaminant (Table 2). The trend was the same in soils amended with agricultural materials (Kidd *et al.*, 2007; An *et al.*, 2004).

$$BF = \frac{\text{mg HM (kg plant leaves)}^{-1} \times 100}{\text{Total content HM (mg kg soil}^{-1})}$$

where:

BF = Bioavailability Factor.
 HM = Heavy Metal.

Sewage sludge has been known to present greater assimilation of the metals when compared to inert material like sand, probably due to the presence of anionic groups, which favors adsorption and complexation processes (Barros *et al.*, 2007; Srivastava *et al.*, 2005). Thus, there is slower metal release from soils treated with sewage (Pampura *et al.*, 2007; Jamali *et al.*, 2006). Metals from mine waste on the other hand is readily available in soil (Kiikkila, 2002).

CONCLUSION

The origin or source of contamination influences soil metal availability and plant uptake, as metals due to mine waste contamination (inorganic) was more readily extractable and available for plant uptake than that from the sewage treated soil (organic). Conversely, metal immobilisation by addition of amendments is greater in soils contaminated by an organic source. For instance, Zn immobilisation was greater in the organic sewage treated soil whereas the same metal was more easily taken up by plants growing in soil contaminated by the inorganic mine waste. Soils subjected to the 2 contaminant sources varied in terms of metal availability, concentration or immobilisation, regardless of similar initial contaminant pH values of 6.4 and 6.9. This suggests that contaminant source may take precedence over pH as dominant factor influencing metal availability in soils where pH values are similar.

Amendments were effective in immobilizing lead, copper and zinc to a certain extent in the contaminated soils, even though the initial metal concentrations in both contaminated soils were at toxic levels. Finally, the findings of the study confirm that where negligible concentrations of metals occur in soils or agricultural materials, plants will take up such metals whenever present, even in non contaminated soil. The evaluation of sewage sludge toxicity by chemical characterisation and biological testing is therefore, extremely important for screening the suitability of sludge for land application.

ACKNOWLEDGEMENT

The author wishes to thank Michael Okpara University of Agriculture Umudike, Nigeria, for granting academic leave to pursue this research and the Commonwealth Scholarship Commission in the UK for funding the study.

REFERENCES

Abbaspour, A., M. Kalbasi, S. Hajrasuliha and A. Fotovat, 2008. Effect of organic matter and salinity on ethylenediaminetetraacetic acid-extractable and solution species of cadmium and lead in 3 agricultural soils. *Commun. Soil Sci. Plant Anal.*, 39: 983-1005.

ADB, 2001. Agricultural biotechnology, poverty reduction and food security. Manila, the Philippines: Asian Development Bank.

Alvarenga, P.M., A.P. Goncalves, R.M. Fernandes, A.P. de Varennes, E.C. Duarte, G. Vallini and C.F. Cunha-Queda, 2008. Effect of organic residues and liming materials on metal extraction from a mining-contaminated soil. *Bioremed. J.*, 12 (2): 58-69.

- An, Y.J., Y.M. Kim, T.I. Kwon and S.W. Jeong, 2004. Combined effect of copper, cadmium and lead upon *Cucumis sativus* growth and bioaccumulation. *Sci. Total Environ.*, 326 (1-3): 85-93.
- Barros, A.J.M., V.D. Shiva Prasad Leite and A.G. Souza, 2007. Biosorption of heavy metals in upflow sludge columns. *Bioresour. Technol.*, 98 (7): 1418-1425.
- Basta, N.T., J.A. Ryan and R.L. Chaney, 2005. Trace element chemistry in residual-treated soil: Key concepts and metal bioavailability. *J. Environ. Quality*, 34 (1): 49-63.
- Clemente, R., A. Escolar and M.P. Bernal, 2006. Heavy metals fractionation and organic matter mineralisation in contaminated calcareous soil amended with organic materials. *Bioresour. Technol.*, 97 (15): 1894-1901.
- Friesl, W., J. Friedl, K. Platzler, O. Horak and M.H. Gerzabek, 2006. Remediation of contaminated agricultural soils near a former Pb/Zn smelter in Austria: Batch, pot and field experiments. *Environ. Pollut.*, 144 (1): 40-50.
- Fuentes, A., M. Llorens, J. Saez, M.I. Aguilar, A.B. Perez-Marin, J.F. Ortuno and V.F. Meseguer, 2006. Ecotoxicity, phytotoxicity and extractability of heavy metals from different stabilised sewage sludges. *Environ. Pollut.*, 143: 355-360.
- Hernandez-Allica, J., C. Garbisu, O. Barrutia and J.M. Becerril, 2007. EDTA-induced heavy metal accumulation and phytotoxicity in cardoon plants. *Environ. Exper. Bot.*, 60 (1): 26-32.
- Jamali, M.K., T.G. Kazi, M.B. Arain, H.I. Afridi, N. Jalbani and R.S. Adil, 2006. The correlation of total and extractable heavy metals from soil and domestic sewage sludge and their transfer to maize (*Zea mays* L.) plants. *Toxicol. Environ. Chemist.*, 88 (4): 619-632.
- Karaca, A., 2004. Effect of organic wastes on the extractability of cadmium, copper, nickel and zinc in soil. *Geoderma*, 122 (2-4): 297-303.
- Kidd, P.S., M.J.D.J. Dominguez-Rodriguez and C. Monterroso, 2007. Bioavailability and plant accumulation of heavy metals and phosphorus in agricultural soils amended by long-term application of sewage sludge. *Chemosphere*, 66 (8): 1458-1467.
- Kiikkila, O., 2002. Remediation Through Mulching with Organic Matter of Soil Polluted by a Copper-nickel Smelter. University of Helsinki, F.O.A.A.F. (Ed.). Finnish Forest Research Institute. Academic Disserta. *Environ. Protect. Sci.*, pp: 57.
- Madrid, F., A.S. Romero, L. Madrid and C. Maqueda, 2006. Reduction of availability of trace metals in urban soils using inorganic amendments. *Environ. Geochem. Health*, 28 (4): 365-373.
- Nwachukwu, O.I., 2007. The use of organic materials in the remediation of soils contaminated by lead, copper and zinc. 236: Ph. D Thesis, University of Glasgow, United Kingdom.
- Nwachukwu, O.I. and I.D. Pulford, 2008. Comparative effectiveness of selected adsorbant materials as potential amendments for the remediation of lead, copper-and zinc-contaminated soil. *Soil Use Manage.*, 24 (2): 199-207.
- Pampura, T., J.E. Groenenberg, S. Lofts and I. Priputina, 2007. Validation of transfer functions predicting Cd and Pb free metal ion activity in soil solution as a function of soil characteristics and reactive metal content. *Water, Air Soil Pollut.*, 184 (1/4): 217-234.
- Pöykkiö, R., H. Nurmesniemi and R.L. Keiski, 2007. Environmental risk assessment of heavy metal extractability in a biosludge from the biological wastewater treatment plant of a pulp and paper mill. *Environ. Monitoring and Assessment*, 128 (1/3): 153-164.
- Roberts, D.A.C.S.A.D.L.S., 2003. Zinc Speciation in Contaminated Soils Combining Direct and Indirect Characterization Methods. In: Selim, H.M. (Ed). *Geochemical and Hydrological Reactivity of Heavy Metals in Soils*. Boca Raton: Lewis Publishers, pp: 187-227.
- Ross, S.M. and K.J. Kaye, 1994. The meaning of metal toxicity in soil-plant systems. In: Ross, M. (Ed.). *Toxic metals in soil-plant systems*. John Wiley and Sons.
- Sharma, R.K., M. Agrawal and F.M. Marshall, 2008. Heavy metal (Cu, Zn, Cd and Pb) contamination of vegetables in urban India: A case study in Varanasi. *Environ. Pollut.*, 154: 254-263.
- Srivastava, P., B. Singh and M. Angove, 2005. Competitive adsorption behavior of heavy metals on kaolinite. *J. Colloid Interface Sci.*, 290 (1): 28-38.
- Udom, B.E., J.S.C. Mbagwu, J.K. Adesodun and N.N. Agbim, 2004. Distributions of zinc, copper, cadmium and lead in a tropical ultisol after long-term disposal of sewage sludge. *Environ. Int.*, 30 (4): 467-470.
- Voutsas, D., A. Grimanis and C. Samara, 1996. Trace elements in vegetables grown in industrial areas in relation to soil and air particulate matter. *Environ. Pollut.*, 94: 325-335.
- Yin, Y.J., C.A. Impellitteri, S.J. You and H.E. Allen, 2002. The importance of organic matter distribution and extract soil: Solution ratio on the desorption of heavy metals from soils. *Sci. Total Environ.*, 287 (1-2): 107-119.