



Morbidity Pattern of Children Admitted to a Paediatric Intensive Care Unit of India

Bandy Sahoo, Sibabratta Patnaik, Reshmi Mishra and Mukesh Kumar Jain

Department of Pediatrics, Kalinga Institute of Medical Sciences, Bhubaneswar, Odisha 751015, India

Key words: Clinical profile, mortality, PICU, innovative facilities, developing countries

Abstract: This study was carried out to assess the morbidity pattern and outcome of children in a PICU (Paediatric Intensive Care Unit) of a tertiary care hospital in Odisha. In developing countries, there are scarce data on paediatric critical care. This makes the modification of practices to improve outcome, difficult. A retrospective study of the demography, clinical profile, diagnosis, treatment and outcome of children admitted to the PICU of Kalinga Institute of Medical Sciences from January 2014 to December 2015 was done. A total of 848 children were admitted of which 61.3% were males and 38.7% females. The leading cause of admission was infectious diseases (20.7%), respiratory disease (19.1%) and central nervous system diseases (14.3%). Multiorgan Dysfunction Syndrome (MODS) and co-morbidity were present in 25 and 22%, respectively. The proportion of deaths among patients admitted to the PICU was 4.1%. Children with MODS and co-morbidity had higher mortality. The overall mortality rate in our PICU was low. We conclude, a well-equipped intensive care unit with modern and innovative facilities leads to a good outcome.

Corresponding Author:

Bandy Sahoo

Department of Pediatrics, Kalinga Institute of Medical Sciences, Bhubaneswar, Odisha 751015, India

Page No.: 80-83

Volume: 15, Issue 4, 2020

ISSN: 1816-3319

International Journal of Tropical Medicine

Copy Right: Medwell Publications

INTRODUCTION

The care of critically ill children remains one of the most demanding and challenging aspects in the field of pediatrics. Pediatric Intensive Care Unit (PICU) aims at promoting early intervention and quality care with an objective of achieving good results and better prognosis. This can be achieved by well-equipped and well-staffed intensive care units^[1,2]. But despite all measures, ICU is one of the sites where medical errors are most likely to occur because of the complexity of the diseases and multiple interventions^[3,4].

With the advancement in intensive care facilities, there is a dramatic increase in survival of critically ill

children^[5,2]. In critical care medicine, Intensive Care Unit (ICU) results can be assessed on the basis of outcome such as mortality rate or survival. Evaluation of the outcome of medical interventions can assess the efficacy of treatment. This helps in better decision making, improving quality of care and modifying the future management if required. Mortality of patients depends on many factors such as demographic variables, clinical characteristic, associated co-morbidities, infrastructure and availability of adequate staffs.

Pediatric intensive care is an emerging concept in the eastern part of India. This is the only PICU in Odisha with trained Pediatric Intensivists working round the clock. Data of PICU in this part of the country is not available

till date. Therefore, PICU data were analyzed to find out the pattern of diseases and outcome at our centre which would help in modifying practices if necessary, leading to better management and outcome of critically ill children.

MATERIALS AND METHODS

This study was a retrospective record based study which reviewed the admissions into the PICU of a tertiary care center in Odisha for a period of 2 years from January 2014 to December 2015. The hospital has a well-equipped ten-bedded PICU which admits pediatric patients upto 18 years of age, from both medical and surgical subspecialties.

PICU records of all admissions, transfers out, discharges and deaths were analyzed. Data collected on patients included age, gender, diagnosis, duration of stay in the unit and outcome. The outcome was classified as transfers to the main pediatric wards, discharges, Discharges Against Medical Advice (DAMA) and death. All patients in the unit were treated according to the written standard protocol. Relevant investigations, including hemoglobin, total and differential blood counts, electrolytes, urea, creatinine, blood glucose, blood culture and arterial blood gas were done at admission. Blood tests were repeated subsequently whenever required. Cerebrospinal fluid analysis was done for suspected central nervous system infections. Treatment was started as per the protocol. Antibiotic therapy was modified whenever necessary depending upon the culture and sensitivity pattern. Vasopressors were used for patients in shock or poor perfusion (Fig. 1).

Suspected sepsis cases (with culture negative) and proven sepsis cases with culture positive body fluid or positive viral marker were included in infectious disease. Suspected sepsis cases included those patients who had Systemic Inflammatory Response Syndrome. (Tachycardia, tachypnoea, temperature 38.5°C or <36°C, abnormal leukocyte count or >10% band cells.) Patients with tropical diseases (malaria/typhoid/dengue/scrub typhus) were also included in this group.




Fig. 1: Gender distribution

RESULTS

During the period of the study, a total of 848 patients was admitted into the PICU with male and female children being 61.3 and 38.7%, respectively. Maximum number of patients belonged to the age group of 1 month to 1 year (47.4%) followed by age group of 1-5 years (24.2%). The mean Length of Stay (LOS) in the PICU was 3.7±2.5 days (range, 0-28 days). The three most common disease categories admitted were infectious diseases (20.7%) followed by respiratory diseases (19.1%), central nervous system diseases (14.3%), cardiovascular diseases (10.8%), gastrointestinal diseases (7%), surgical problems (4.7%) hematological (4%), renal (3.3%), poisonings (1.4%) and others (14.3%). 596(70.3%) patients improved and were transferred to the paediatric wards, 205(24%) were discharged directly from PICU, 35(4.1%) died and 12(1.4%) left against medical advice. Seventy two (8.5%) children received mechanical ventilation, among which 45(62.5%) improved, 25(34.7%) died and 2(2.8%) children left against medical advice (Fig. 2).

A total number of 782 culture samples were sent out of which 35(4.5%) samples were positive. Out of the 56 cases of sepsis, 15(26.8%) had positive blood culture. Staphylococcus sepsis was the commonest blood stream infection. Out of the 72 patients who were mechanically ventilated, 4(5.5%) had endotracheal secretions culture positive. The rate of nosocomial infection was 4.7% with acinetobacter from endotracheal secretions and staphylococcus aureus from central venous catheter as common isolates.(nosocomial infection is to be defined) Bloodstream infection from central venous catheter (mostly placed in femoral vein) and ventilator associated pneumonias were the main sites of infections in patients with nosocomial infections.

Thirty five (4.1%) patients died during the period, consisting of nineteen (59.3%) males and thirteen (40.6%) females. Leading causes of death in this study were sepsis with MODS (n = 11), encephalitis (n = 7), pneumonia (n = 7) and congenital heart diseases (n = 5). Maximum deaths 13 (37.1%) occurred in the age group 1 month to 1 year. Mortality analysis in relation to different diseases is presented in Table 1 and 2.




Fig. 2: Age distribution of patients

Table 1: Distribution as per morbidity pattern

Condition	No. of patients	Total admissions (%)
Infections/Sepsis	176	20.7
Respiratory	162	19.1
Neurological	122	14.3
Cardiovascular	92	10.8
Gastrointestinal	60	7
Surgical	40	4.7
Hematological	34	4
Renal	28	3.3
Poisoning	12	1.4
Others	122	14.3
Total	848	100.0

Table 2: Mortality according to disease

Diseases	No. of cases	Death n = 35 (%)
Sepsis	56	11(31.4)
Pneumonia	86	7(20)
Encephalitis	51	7(20)
Congenital heart disease	23	5(14.3)
Leukemia	7	2(5.7)
Aplastic anemia	5	1(2.8)
Hepatic encephalopathy	1	1(2.8)
Retinoblastoma	1	1(2.8)

Common respiratory illnesses included pneumonia (n = 86, 53%), acute bronchiolitis (n = 40, 24.7%) and bronchial asthma (n = 32, 19.75%). Encephalitis (n = 51, 41.8%) followed by seizure disorders (n = 35, 28.7%) and meningitis (n = 11, 9%) were the common central nervous system diseases requiring PICU care. Other conditions requiring PICU admissions included acute gastroenteritis (n = 13, 76.4%), congenital heart diseases (n = 8, 53.3%), and poisoning (n = 4, 36.3%).

DISCUSSION

The PICU is a special unit primarily concerned with the care of patients with critical illness and demands a broad-based knowledge to achieve good outcome^[6]. Advances in pediatric sub-specialties including the critical care medicine have improved the survival of sick children. During the 24 months study period, 848 children were admitted to the 10 bedded PICU (432 and 416 in each year) which is comparable to other tertiary level PICUs in the country^[7]. Majority of the patients were males (61.3%) a finding similar to that of Shah^[8] and 32% were infants as recorded by Haque and Bano^[9, 10]. This study revealed that infectious diseases (20.7%), respiratory diseases (19.1%) and central nervous system diseases (14.3%) were the major causes of admission into the PICU. Shah *et al.*^[8] reported respiratory illness (33%) as the commonest indication for admission^[11], reported cardiovascular disease (41.1%) as the commonest indication for admission in their series while a study from Pakistan found post cardiac surgery (34%) to be the most common condition^[9, 10]. This shows that paediatric intensive care admissions vary in different countries and one should be aware of the prevalent conditions to develop the facilities and prepare treatment protocols accordingly.

Overall mortality in this study was 4.1%, giving an ICU survival rate of 95.9%. This value is higher than documented by Shah *et al.*^[8] with the mortality rate (2.1%) and Choi *et al.*^[12] with the mortality rate (2.6%) for a five-bed PICU in a general hospital in Hong Kong. It is, however, less than an overall mortality of 6.7 and 16.7% recorded in India by Khilnani *et al.*^[7] and Bellad *et al.*^[13], respectively. The reported mortality varied from 9.8-35% in different series by other authors^[14, 9, 10, 15]. About 31.4% of non-survivors had multiorgan failure which is lower as compared to the study by Khilnani *et al.*^[7] (49.5%). The average LOS (3.7±2.5 days) in PICU of the present study is similar to that of Abhulimhen-Iyoha *et al.*^[11] but in contrast to mean duration of 4.52±2.6 days reported by Khilnani *et al.*^[7]. The absence of a high-dependency unit at our center led to the admission of some patients who were not ill enough to remain in PICU. To enhance cost-effective management of patients and avoid unnecessary stretch of the ICU stay this situation needs to be addressed. This was one of the limitations of our study. The other limitation was inability to assess the severity scoring.

Based on our observation, it appears that care of patients in our PICU is somewhat similar to other tertiary level PICUs in the country. The low mortality rate in our PICU could be due to medical college affiliation with better resources availability, 24-h physician coverage, highly trained-nurses, good nurse-patient ratio and presence of 4 trained paediatric intensivists. Pearson *et al.*^[16] have suggested that the availability of full-time trained paediatric intensivists can deliver care of high quality and with much higher efficiency than without them in PICUs.

CONCLUSION

We conclude that the demographic profile of patients including age, sex, source of admission and co morbidities follow a varied pattern in different PICU patients worldwide. The low mortality rate indicates optimal quality management of our patients. A well-equipped intensive care unit with modern and innovative facilities along with the availability of fulltime trained paediatric intensivists made a significant impact on the outcome of critically ill children in our PICU.

REFERENCES

- Earle, M., O.M. Natera, A. Zaslavsky, E. Quinones and H. Carrillo *et al.*, 1997. Outcome of pediatric intensive care at six centers in Mexico and Ecuador. *Crit. Care Med.*, 25: 1462-1467.

02. Young, M.P. and J.D. Birkmeyer, 2000. Potential reduction in mortality rates using an intensivist model to manage intensive care units. *Eff. Clin. Pract. ECP.*, 3: 284-289.
03. Cullen, D.J., B.J. Sweitzer, D.W. Bates, E. Burdick, A. Edmondson and L.L. Leape, 1997. Preventable adverse drug events in hospitalized patients: A comparative study of intensive care and general care units. *Crit. Care Med.*, 25: 1289-1297.
04. Pronovost, P., A.W. Wu, T. Dorman and L. Morlock, 2002. Building safety into ICU care. *J. Crit. Care*, 17: 78-85.
05. Tilford, J.M., P.K. Roberson, S. Lensing and D.H. Fiser, 1998. Differences in pediatric ICU mortality risk over time. *Crit. Care Med.*, 26: 1737-1743.
06. Downes, J.J., 2007. Development of Paediatric Critical Care Medicine-How Did we Get Here and Why?. In: *Paediatric Critical Care Medicine: Basic Science and Clinical Evidence*, Wheeler, D., H. Wong and T. Shanely (Eds.), Springer, London, UK., pp: 3-32.
07. Khilnani, P., D. Sarma, R. Singh, R. Uttam, S. Rajdev, A. Makkar and J. Kaur, 2004. Demographic profile and outcome analysis of a tertiary level pediatric intensive care unit. *Apollo Med.*, 1: 161-166.
08. Shah, G.S., B.K. Shah, A. Thapa, L. Shah and O.P. Mishra, 2014. Admission patterns and outcome in a pediatric intensive care unit in Nepal. *J. Adv. Med. Res.*, 4: 4939-4945.
09. Haque, A. and S. Bano, 2009a. Clinical profile and outcome in a paediatric intensive care unit in Pakistan. *J. Coll. Phys. Surg. Pak.*, 19: 534-535.
10. Haque, A. and S. Bano, 2009b. Improving outcome in pediatric intensive care unit in academic hospital in Pakistan. *Pak. J. Med. Sci.*, 25: 605-608.
11. Abhulimhen-Iyoha, B.I., S.K. Pooboni and N.K.K. Vuppali, 2014. Morbidity pattern and outcome of patients admitted into a pediatric intensive care unit in India. *Indian J. Clin. Med.*, Vol. 5, 10.4137/IJCM.S13902
12. Choi, K.M., D.K. Ng, S.F. Wong, K.L. Kwok, P.Y. Chow and C.H. Chan, 2005. Assessment of the pediatric index of mortality (PIM) and the pediatric risk of mortality (PRISM) III score for prediction of mortality in a paediatric intensive care unit in Hong Kong. *Hong Kong Med. J.*, 11: 97-103.
13. Bellad, R., S. Rao, V.D. Patil and N.S. Mahantshetti, 2009. Outcome of intensive care unit patients using pediatric risk of mortality (PRISM) score. *Indian Pediatr.*, 46: 1091-1092.
14. Singhal, D., N. Kumar, J.M. Puliyl, S.K. Singh and V. Srinivas, 2001. Prediction of mortality by application of prism score in intensive care unit. *Indian Pediatr.*, 38: 714-719.
15. Costa, G.D.A., A.F. Delgado, A. Ferraro and T.S. Okay, 2010. Application of the Pediatric Risk of Mortality Score (PRISM) score and determination of mortality risk factors in a tertiary pediatric intensive care unit. *Clin.*, 65: 1087-1092.
16. Pearson, G., F. Shann and D. Field, 1997. Should pediatric intensive care be centralized? Trent versus Victoria. *Lancet*, 349: 1213-1217.