

## Effects of Different Rations on the Slaughter and Carcass Characteristics of Ak Keçi (White Goat) Male Kids

<sup>1</sup>Apkin Kor, <sup>2</sup>Mehmet Ertugrul and <sup>3</sup>Serhat Arslan

<sup>1</sup>University of Yuzuncu Yil, Faculty of Agriculture, Department of Animal Science, 65080 Van-Turkey; <sup>2</sup>University of Ankara, Faculty of Agriculture, Department of Animal Science, 06110, Ankara, Turkey; <sup>3</sup>University of Yuzuncu Yil, Faculty of Agriculture, Department of Animal Science, 65080 Van-Turkey

**Abstract:** Aim of this study was to determine the slaughter and carcass characteristics of Akkeçi (White Goat) male kids which were fed on a standard mixture and vitamin–mineral enriched barley. Flank joints ratio in left half carcass weight was  $13.62 \pm 0.734\%$  at only barley group and  $11.69 \pm 0.367\%$  at standard group. Only flank joints between the two groups were found to be significantly different. At the end of this study, it was determined that differences between barley and standard mixture rations on slaughter data and carcass characteristics in Akkeçi male kids were insignificant.

**Key words:** Dressing percentage, omental and mesenteric fat, carcass joints

### Introduction

In Turkey, goat rising has an extensive characteristic due to the social and economical conditions of the farmers, natural conditions and environmental factors which are supplied to goat production. As a result of this extensive farming and the restricted genotypic capacities of our native goat races, yields per goat are extremely low. However, goat stocks across the different parts of Turkey still have an important place in both living of many farmers and in Turkey's economy.

Although fattening performance of goats is not as enough as other ruminants, there has been increasing interest towards farming of goats because of both their increasing importance in meat production and their high lactation capacity per body weight. Therefore, it would be appropriate to choose the best goat farming methods in order to positively affect this interest.

In general information about goat farming and its products is limited in spite of all efforts. Knowledge accumulation by doing various researches will help to both goats farming to be done more rationally and to realize the value of goat products.

Aim of this study was to determine the slaughter and carcass characteristics of Akkeçi (White Goat) male kids which were fed with a standard mixture and vitamin–mineral enriched barley.

### Materials and Methods

A total of 16 heads Akkeçi male kids which were weaned at 2.5 month-old age and then fed with different mixtures given at Table 1 for 75 days were obtained from the Research and Experiment Farm of Ankara University and constituted the animal materials.

Table 1: Composition of mixtures and their analytical values

Materials	Rations	
	Barley (%)	Standard (%)
Barley	96.65	32.30
Maize	--	27.40
Soybean pulp	--	8.00
Cotton seed pulp	--	6.00
Sunflower seed pulp	--	12.20
Wheat bran	--	10.45
CaCO <sub>3</sub>	1.80	2.80
DCP	0.70	--
Salt	0.50	0.50
Preliminary vitamin mixture	0.25	0.25
Preliminary mineral mixture	0.10	0.10
Total	100.00	100.00
<b>Analytical values</b>		
Protein (CP, %)	11.66	17.66
Energy (NB/kg)	724.87	669.89

Male kids were separated into two groups, each has 8 kids. One of the groups was fed on vitamin-mineral enriched barley; the other was fed with standard mixture ad-libitum. Moreover, kids were fed on minced clover as 1% of their body weight twice a day. Fresh and clean water was supplied to the kids during the trial.

Live measurements of all kids were taken before slaughter (Ertugrul, 1996). Hot carcass, hearth, lungs-livers, spleen, omental fat, hide, four feet and head were weighted soon, after slaughter. Carcasses were then placed in a chiller operating at 4 °C for 24 hours. After chilling cold carcasses were measured and weighted (Hankins, 1959), then separated into joints according to a proposed standard method for goats in Mediterranean Countries (Colomer-Rocher *et al.*, 1987). Each joint were also separately weighted.

Means of groups were compared with 't-test' in using SAS (1998).

## Results and Discussion

Pre-slaughter body and 24h after slaughter carcass measurements of kids are given at Table 2.

As it seen Table 2, there is a significant similarity between kids fed on barley and standard mixture in terms of body and carcass measurements ( $P > 0.05$ ). Values obtained in this study were in line with those other studies (Kor and Ertugrul, 2000; Kor, 1997; Eker *et al.*, 1976; Elicin *et al.*, 1976; Akman *et al.*, 1991; Daskiran, 1992 and Keles, 1995).

Slaughter characteristics and its proportion at slaughterhouse and hot carcass weight data are given in Table 3 and Table 4.

In this study, slaughter characteristics of male kids fed with two different rations were not significantly different. On the other hand, omental and mesenteric fat, one of the important slaughter parameters, was higher in the subjects fed with standard mixture than that of barley group. This is in favor of male kids fed with barley mixture because omental fat is not utilized with carcass. Moreover, slaughter characteristics determined in this study were in agree with those of other researches (Kor, 1997; Guney *et al.*, 1984 and Yargici and Yener, 1991). However, omental and mesenteric fat in those of this study was lower than those of some studies ( Bayraktaroglu *et al.*, 1988 and Hogg *et al.*, 1992), and was higher than those of some studies ( Kor and Ertugrul, 2000 and Kor, 1991). There were no significant differences between two groups of kids fed on different rations on the proportion of slaughter by-products to body weight and hot carcass weight.

Carcass composition of 24hrs chilled (at +4 °C) kid carcasses of research materials are seen at Table 5.

As seen at Table 5, there were no significant differences between two groups of kids fed on different rations on carcass composition and weights of left-half carcass joints. However, similar to omental and mesenteric fat, there was more accumulation of kidney knob and channel fat in kids fed on standard mixture.

Carcass loss which was calculated as  $100 * (\text{cold carcass weight} - \text{hot carcass weight}) / \text{cold carcass weight}$  was reported as  $4.00 \pm 0.668$  % for kids fed with barley mixture and  $3.60 \pm 0.190$  % for kids fed with standard mixture. Although this was not significantly different, the low carcass loss in later group was due to lower omental and kidney-pelvis fat amount of them.

Table 2: Body and carcass measurements (cm)

	Barley Group	Standard Group
	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$
<b>Body measurements (n = 8)</b>		
Wither height	57.25 ± 0.790	56.00 ± 0.790
Body length	58.13 ± 1.080	58.31 ± 1.080
Heart girth depth	24.38 ± 0.350	24.94 ± 0.350
Heart girth width	15.13 ± 0.470	15.75 ± 0.470
Heart girth circumference	70.38 ± 1.350	71.00 ± 1.350
Leg circumference	61.38 ± 1.350	62.38 ± 1.350
<b>Carcass measurements</b>		
Leg depth	9.13 ± 0.160	9.00 ± 0.160
Leg width	5.63 ± 0.240	5.88 ± 0.240
Leg length	24.50 ± 0.380	24.50 ± 0.380
Rump width	14.25 ± 0.530	14.63 ± 0.530
Heart girth depth	23.50 ± 0.247	24.13 ± 0.247
Heart girth width	15.75 ± 0.547	15.50 ± 0.547
Shoulder width	14.25 ± 0.371	14.44 ± 0.371
Carcass length	59.25 ± 0.873	58.50 ± 0.873

Table 3: Slaughter characteristics

Characteristics (n = 8)	Barley Group	Standard Group
	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$
Slaughter weight (kg)	26.05 ± 0.990	27.85 ± 0.990
Hot carcass weight (kg)	13.01 ± 0.667	13.86 ± 0.667
Dressing percentage (%)	49.89 ± 0.950	49.50 ± 0.950
Head weight (kg)	1.75 ± 0.092	1.98 ± 0.092
Pelt weight (kg)	1.95 ± 0.120	1.98 ± 0.120
Four feet weight (g)	712.50 ± 34.395	787.50 ± 34.395
Omental fat weight (g)	376.90 ± 47.050	494.40 ± 47.050
Heart, lungs, liver weight(kg)	1.03 ± 0.035	1.03 ± 0.035
Spleen weight (g)	36.25 ± 2.950	38.75 ± 2.950

Table 4: Proportions of slaughter by-products to body and hot carcass weight (%)

Characteristics	In Body Weight		In Hot Carcass	
	Barley Group	Standard Group	Barley Group	Standard Group
	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$
Head	6.72 ± 0.167	7.07 ± 0.167	--	--
Four feet	2.74 ± 0.093	2.84 ± 0.093	--	--
Pelt	7.49 ± 0.348	7.06 ± 0.348	--	--
Omental fat	1.45 ± 0.145	1.77 ± 0.145	2.93 ± 0.291	3.56 ± 0.291
Heart, lungs, liver	3.96 ± 0.121	3.72 ± 0.121	7.97 ± 0.350	7.55 ± 0.350
Spleen	0.14 ± 0.008	0.14 ± 0.008	0.28 ± 0.017	0.28 ± 0.017

Table 5: Various carcass characteristics and weights of left-half carcass joints

Characteristics (n = 8)	Barley Group	Standard Group
	$\bar{X} \pm S\bar{X}$	$\bar{X} \pm S\bar{X}$
Cold carcass weight (kg)	12.48 ± 0.640	13.36 ± 0.640
Cold dressing percentage (%)	47.87 ± 0.903	47.73 ± 0.903
Carcass loss (%)	4.04 ± 0.490	3.60 ± 0.490
Testicle weight (couple) (g)	160.00 ± 22.420	213.13 ± 22.420
Kidney weight (couple) (g)	78.13 ± 2.900	81.88 ± 2.900
Kidney-knob and pelvis fat (g)	330.00 ± 43.765	405.63 ± 43.765
Left-half carcass weight (kg)	6.05 ± 0.230	6.23 ± 0.230
Carcass joints weights (kg)		
Shoulder	1.28 ± 0.067	1.36 ± 0.067
Hind leg	1.81 ± 0.095	1.93 ± 0.095
Neck	0.57 ± 0.084	0.69 ± 0.084
Flank	0.82 ± 0.049	0.73 ± 0.049
Back loin	1.30 ± 0.120	1.50 ± 0.120

Proportions of carcass joints are presented at Table 6 in order to explain the distribution of these joints.

Only flank joints between the two groups found to be significantly different ( $P < 0.05$ ).

Carcass composition of this study was similar to those of most other researchers, but higher than some researchers (Kor and Ertugrul, 2000; Keles, 1995; Guney and Cayan, 1987 and Ertugrul, 1994).

In various studies were stated that muscles amounts of carcass in sheep and goats are correlated with *musculus longissimus dorsi* (MLD) area which is taken between 12<sup>th</sup> and 13<sup>th</sup> ribs, and MLD areas of 20 kg weighted various goat genotypes from 4.03 to 16.12 cm<sup>2</sup> (Devendra and Owen, 1983). In Turkey, MLD areas were found to be ranged 8.36 to 10.05 cm<sup>2</sup> (Kor and Ertugrul, 2000; Kor, 1997; Akman *et al.*, 1991 and Daskiran, 1992). MLD area of this study was in the range of Devendra and Owen (1983), but was higher than those of other studies (Kor and Ertugrul, 2000; Kor, 1997; Akman *et al.*, 1991 and Daskiran, 1992).

This study was conducted to determine the slaughter data and carcass characteristics of Akkeçi male kids fed with two rations with different content and cost. Findings support that ration with barley can supply adequate performance for lamb feeding (Alarслан, 1993; Celik, 1993 and Isik *et al.*, 1987). However, there was no study

Table 6: Proportions of carcass joints (%)

Characteristics (n = 8)	Barley Group $\bar{X} \pm S_{\bar{X}}$	Standard Group $\bar{X} \pm S_{\bar{X}}$
In left-half carcass weight		
Shoulder	21.12 ± 0.280	21.86 ± 0.280
Hind leg	29.88 ± 0.380	31.02 ± 0.380
Neck	9.57 ± 1.190	10.91 ± 1.190
Flank	13.62 ± 0.580	11.69 ± 0.580
Back loin	23.53 ± 1.570	24.08 ± 1.570
In cold carcass weight		
Testicle (couple)	1.27 ± 0.169	1.61 ± 0.169
Kidney (couple)	0.63 ± 0.028	0.62 ± 0.028
Kidney-knob and pelvis fat	2.63 ± 0.280	3.04 ± 0.280
MLD area (cm <sup>2</sup> )	11.28 ± 0.880	10.51 ± 0.880

\* P &lt; 0.05

on the effects of barley ration to slaughter data and carcass characteristics in kids.

In conclusion, it was determined that differences between barley and standard mixture rations on slaughter and carcass characteristics in Akkeçi male kids were not statistically different. Moreover, results obtained in this study were in agreement with the other studies worked on slaughter and carcass characteristics. Therefore it would be appropriate and economic for farmers to use vitamin-mineral added barley mixture instead of standard mixture.

## References

- Akman, N., M. Ertugrul, A. Tatayoglu, A. Kor and A.U. Yavuzer, 1991. Slaughter and Carcass Characteristics of Angora Goat (in Turkish). *Journal of Lalahan Livestock Research Institute*, 31: 3-4.
- Alarслан, O.F., 1993. A Comparative Study on the Effects of Lamb Fattening Ration and Barley Ration on Fattening Performance, Wool Yield and Some Carcass Characteristics of Lambs. Ankara University, Faculty of Agricultural. Publication, 1298.
- Bayraktaroglu, E., N. Akman and E. Tuncel, 1988. Effects of Early Castration on Slaughter and Carcass Characteristics in Crossbred Saanen x Kilis Goats. *Small Ruminant Res.*, 1:189-194.
- Colomer-Rocher, F., P. Morand-Fehr and A.H. Kirton, 1987. Standard Methods and Procedures for Goat Carcass Evaluation Jointing and Tissue Separation. *Livestock Prod. Sci.*, 17:149-157.
- Celik, B., 1993. Effects of Urea addition to Barley as a Protein Source to Fattening Performance of Lambs (in Turkish). Ankara University, Graduate School of Natural and Applied Sciences MSc.(Agric) Thesis, (Unpublished).
- Daskiran, I., 1992. Fattening Performance and Carcass Characteristics of Weaned Angora Male Kids. Ankara University, Graduate School of Natural and Applied Sciences MSc. (Agric) Thesis.
- Devendra, C. and J.E. Owen, 1983. Quantitative and Qualitative Aspects of Meat Production From Goats. *World Animal Review*, 47:19-29.
- Eker, M., E. Tuncel, Y. Askin and S.M. Yener, 1976. Studies on Live Weight and Body Development of Saanen x Kilis Goats. *Annual of Ankara Univ. Agri. Faculty*, 26:164-175.
- Elicin, A., F. Tepe and E. Tuncel, 1976. Adaptation Studies of Saanen x Kilis Dairy Goats to the Condition of Antalya Agricultural Research Institute. II. Live Weight, Body Measurements and Growth Rate. *Annual of Ankara University, Agricultural Faculty*, 26:107-120.
- Ertugrul, M., 1994. Slaughter and Carcass Characteristics of Angora Male Kids Intensively Fed at Weaning Period and Extensively Raised (in Turkish). Ankara University, Faculty of Agricultural Publication, 1392.
- Ertugrul, M., 1996. Applications of Small Ruminant Raising. Second Edition. Ankara University, Faculty of Agricultural Publication, 1446.
- Guney, O. and O. Cayan, 1987. The Fattening Performance and Carcass Characteristics of Hair Male Kids Under Intensive Feeding Conditions. *Philoetions Symposium on the Evaluation of Mediterranean Sheep and Goat, Fonte-boa*, 23-25 Portugal.
- Guney, O., F. Eroglu and O. Bicer, 1984. Effects of Castration on Fattening Performance and Carcass Characteristics in Kids (in Turkish). *Journal of Natural Sciences, TUBITAK*. D1: 8.
- Hankins, O.G., A.M. Gaddis and W.I. Sulzbacher, 1959. Meat Research Techniques Pertinent to Animal Production Research. *Techniques and Procedures in Animal Production Research*. American Society of Animal Production. 194-221.

- Hogg, B.W., G.J.K. Mercer, B.J. Mortimer, A.H. Kirton and D.A. Duganzich, 1992. Carcass and Meat Quality Attributes of Commercial Goats in New Zealand. *Small Ruminant Res.*, 8: 243-256.
- Isik, N., O.F. Alarslan and D. Gormus, 1987. Effects of Barley Ration Supplemented with Different Amounts of Cotton Seed and Wheat Pulps on Anatolian Merinos Lambs.(in Turkish). *Annual of Ankara University, Agricultural Faculty*, 38.
- Keles, M.A., 1995. Fattening Performance and Carcass Characteristics of Hair Goat Male Kids Fed on at Weaning Period (in Turkish). Ankara University, Graduate School of Natural and Applied Sciences MSc. (Agric). Thesis.
- Kor, A. and M. Ertugrul, 2000. Some Slaughtering- Carcass Characteristics and Meat Composition of Akkeçi Male Goats. *Ankara. Univ. J. Agri. Sci.*, 6:86-91.
- Kor, A., 1997. The Effects of Castration at Different Ages of Male Kids of Various Genotypes on the Fattening Performance, Carcass Characteristics and Meat Quality. Ankara University, Graduate School of Natural and Applied Sciences Ph.D. (Agric). Thesis.
- Kor, A., 1991. The Growth Performance and Carcass Characteristics of Damascus x Hair Goat (Fi) Crossbred Male Kids (in Turkish). Cukurova University, Graduate School of Natural and Applied Sciences MSc. (Agric) Thesis.
- SAS, 1998. SAS Statistics Software, SAS. Inc., Cary, NC., 27513.
- Yargici, S. and S.M. Yener, 1991. Effects of Early Weaning on Fattening Performance, Growth and Some Reproduction Characteristics of Akkeçi Goats. *Journal of Faculty of Agriculture Akdeniz University.*, 4:39-53.