

Alterations in the Pituitary- Thyroid Axis in Camel *Camelus dromedarius* Infected by Larvae of Nasal Bot Fly *Cephalopina titillator*

¹Ghada M. El Bassiony, ²Othman A. Al Sagair, ³Ezzat S. El Daly and ⁴Asmaa M. El Nady

¹Entomology Department, Faculty of Science, Cairo University, Giza, Egypt.

²Medical Microbiology Department, Faculty of Science, El Qassim University, El Qassim, KSA

³Department of Zoology, Faculty of Science, Helwan University, Cairo, Egypt

⁴Animal Health Research Institute, Agricultural Research Center, Dokki, Giza, Egypt

Abstract: The present study used the Chemiluminescent Microparticle Immunoassay (CMIA) to examine the alterations in pituitary-thyroid function, in camels infected with third instar larvae of *Cephalopina titillator*, by measuring the levels of triiodothyronine (T3) and thyroxine (T4) of the thyroid gland and level of thyroid stimulating hormone (TSH) of the pituitary gland. The results indicated that the infection of camels by larvae of the nasal bot fly caused a case of hypothyroidism. This was evidenced from the declines recorded in blood levels of T3 and T4. Also the data obtained point to the occurrence of a parallel decrease in the level of blood TSH. It appears that the depressed release of TSH during infection with *C. titillator* together with the subsequent decline in T3 and T4 levels from thyroid gland, might reflect the direct effect of infection on pituitary gland, and suggest decreased synthesis of T3 and T4 in response to larval infection of the nasal bot fly.

Key words: *Cephalopina titillator*- *Camelus dromedarius*- T3- T4- TSH

Introduction

The camel nasal bot fly, *Cephalopina titillator* (subfamily: *Oestrinae*) is the only species in its genus (Sharma, 1992). It develops in the nasal cavities of camels and shows a high degree of host specificity. High level of infestation causes congestion of nasal cavity with mucous, severe inflammatory and degenerative changes, leading to extensive damage of nasopharyngeal tissues and the formation of lymphoid nodules at the site of larval attachment in the pharynx. The affected camels may snort, show abnormal behaviour and nervous symptoms including restlessness, cramp, excitation, convulsions, difficulties in breathing and may lead to death following complications due to secondary infections (Hussein *et al.*, 1982 and 1983; Higgins, 1985; Derhalli *et al.*, 1989 and Musa *et al.*, 1989).

In the previous studies, no results have linked *C. titillator* -infected camels with serum thyroid hormone concentrations (T3 and T4) or serum thyroid stimulating hormone (TSH) level. On the other hand, it was reported that *Trypanosoma congolense* infection rapidly impaired the function of the thyroid gland in goats as defined by considerable plasma thyroxine (T4) decrease (Mutayoba *et al.*, 1988) and hypothyroidism in cattle (Abebe and Eley, 1992). Experimental infection of rabbits with *Trypanosoma congolense* induced a rapid decline in both T3 and T4 (Lomo *et al.*, 1996). Similarly, studies on human sleeping sickness showed a significant decrease in 3, 5, 3'-triiodothyronine (T3) and (T4) without TSH variation (Boersma *et al.*, 1989). Al-Qarawi *et al.* (2001) indicated that *Trypanosoma evansi* caused a significant case of hypothyroidism evidenced as a decrease in the thyroid triiodothyronine (T3) and thyroxine (T4) blood levels associated with a parallel decrease in the level of blood pituitary thyrotrophic hormone (TSH).

The objective of the present investigation was to measure the levels of T3, T4 and TSH in the camel *Camelus dromedarius* during the infection with larvae of *Cephalopina titillator*.

Materials and Methods

Experimental Animals: For this study, adult male and female camels, *Camelus dromedarius* were used. Animals were obtained from abattoir of El-Basatein, Cairo, Egypt

Groups of Animals Under Investigation: Camels were divided into two main groups. Each group consists of 7 animals

Group 1: This group is the control animals. They were free from any larval instars of the studied insect, *Cephalopina titillator*

Group 2: The animals of this group were infected by the larvae of *C. titillator* with third instar larval burdens of 10 – 15 (mean 13) per each camel

Blood and Tissue Sampling: Animals were sacrificed by cervical dislocation. Blood was collected in separated clean centrifuge tubes, allowed to coagulate and serum was separated by centrifugation at 3000 r.p.m. for 20 min. Serum specimens were quickly kept frozen at - 20°C until needed for analysis of thyroid hormones (triiodothyronine, T3 and thyroxine, T4) and thyroid stimulating hormone (TSH). The levels of T3, T4 and TSH in blood serum were determined using the Chemiluminescent Microparticle Immunoassay (CMIA). CMIA kits were provided by Abbott Laboratories Diagnostics Division, Abbott Park, IL 60064, USA.

Statistical Analysis: The results are reported as the mean \pm S.E. Differences between means were determined by using one way analysis of variance (ANOVA) and F-test followed by Tukey's multiple range test using the Statistical Package for the Social Sciences (SPSS) version 10. Non- significant level was taken at $P > 0.05$.

Table 1: Changes in levels of triiodothyronine (T3), thyroxine (T4) and thyroid stimulating hormone (TSH) in serum of camels infected with third instar larvae of *Cephalopina titillator*.

Parameter	Groups	Mean ± S.E	% of Change
Triiodothyronine (T3)	Non-infected (Control)	120.0 ± 20.80	- 78.52 ***
	Infected	25.78 ± 7.08	
Thyroxine(T4)	Non-infected (Control)	11.62 ± 1.33	- 23.58 *
	Infected	8.88 ± 1.89	
Thyroidstimulatinghormone(TSH)	Non-infected(Control)	0.014 ± 0.003	- 14.29
	Infected	0.012 ± 0.002	

Numbers of camels in each group = 7
 Non-significant at P>0.05.
 *** Very highly significant at P<0.0001.
 * Significant at P<0.01

Absolute value is measured by ng/ml for T3; mg/dl for T4 and μ U/ml for TSH.
 * Significant at P<0.01.
 Each bar represents a percentage of change value.
 *** Very highly significant at P<0.0001

Results and Discussions

Results of thyroid hormones (T3 and T4) and thyroid stimulating hormone (TSH) in blood serum of non-infected (control) and infected camels are shown in Table (1) and graphically illustrated in Fig. 1.

Triiodothyronine (T3): The response of thyroid hormones for the action of third larval instar of *C. titillator* infection shows a very highly significant decrease (F=146.35, P<0.0001) in the level of T3 in serum. The value (expressed as percentage of change) was -78.52%.

Thyroxine (T4): The analysis of T4 level in serum of infected camels showed a value of -23.58%. (expressed as percentage of change). This value indicates the occurrence of a significant decrease (F=6.07, P<0.01).




Fig. 1: Effect of infection with 3rd instar larvae of *Cephalopina titillator* on triiodothyronine (T3), thyroxine (T4) and thyroid stimulating hormone (TSH) in serum of camels

Thyroid stimulating hormone (TSH): Studies were undertaken to examine the TSH level of serum in response to third larval instar of *C. titillator* infection. The results revealed the occurrence of a non-significant decrease (F=0.40, P>0.05) of -14.92%.

Although the literature contains many references to the effect of different parasites on the pituitary- thyroid axis in mammals (Ikede and Losos, 1975; Mutayoba *et al.*, 1988; Boersma *et al.*, 1989; Abebe and Eley, 1992; Abebe *et al.*, 1993; Lomo *et al.*, 1996; Haroun *et al.*, 2000 and Al-Qarawi *et al.*, 2001), no direct estimates of the effect of nasal bot fly *C. titillator* on serum T3, T4 and TSH in camels have been made before.

The present results indicated that larvae of *C. titillator* caused a significant case of hypothyroidism evidenced as decrease in the thyroid triiodothyronine (T3) and thyroxine (T4) blood levels associated with a parallel decrease in the level of blood pituitary thyrotropic hormone (TSH). These data suggest that the depressed release of TSH during infection together with the subsequent decline in T3 and T4 levels from thyroid gland, might reflect the direct effect of infection on pituitary and thyroid gland. In support of these observations, Al-Qarawi *et al.* (2001) indicated that decreases in the levels of T3, T4 and TSH of blood in camels infected with *Trypanosoma evansi* were recorded.

In hypothyroid animals the synthesis of long chain fatty acids from glucose and other carbohydrate precursors, as well as their oxidation, were reduced (Bray and Goodman, 1968). Since both of these processes require movements of metabolites across the mitochondrial membrane, it has been postulated that the metabolic defect in hypothyroidism may reside in the mitochondrial membrane (Bray and Goodman, 1968). Lipogenesis as well as lipolysis were reported to be largely affected by thyroid hormones (Fisher and Bull, 1967)

Roncari and Murthy, 1975): T3 and T4 have pronounced physiological effects in the control of respiration and energy metabolism as well as in the biogenesis of the mitochondria. Their production is important in conditions such as disease that leads to alter metabolic

status. The thyroid gland is one of the endocrine glands, which is affected during parasite infection. In our study, the infection was indicated by a significant reduction in the level of T3 and T4 blood concentrations. This impairment in thyroid function could be explained on the basis of a direct or/and an indirect (through the pituitary gland) effect of the larvae on the thyroid gland. The direct effect may be mediated by biologically active factors of parasite origin such as phospholipase A (Tizard *et al.*, 1978), protease (Lonsdale-Eccles and Grab, 1986) and peptidase (Knowles *et al.*, 1989) thereby causing thyroid dysfunction. On the other hand, indirect pathological changes which occur during parasites infection can lead to pituitary dysfunction (Ikede and Losos, 1975 and Mutayoba, *et al.*, 1988). The presence of larvae of *C. titillator* within the nasopharyngeal cavity and their causes for abnormal movement of the head, bleeding from the nostrils and meningitis in camels (Hussein *et al.*, 1982; AL-Ani *et al.*, 1991 and Bekele, 2001), might alter the pattern of pituitary TSH production and thereby prevent normal thyroid function. Proteases of parasitic origin have been implicated in the gonadotrophic dysfunction in rats infected with *T. Brucei Brucei* (Humblart *et al.*, 1990). Thus, an indirect antithyroid effect could have been mediated through; activation of epithelial cells, monocytes and macrophages in the immune system during infection releasing cytokines which activate and down regulate the endocrine system. The results of De Souza *et al.* (1989) and Honegger *et al.* (1991) suggested that IL-1 may be involved in the paracrine and/or autocrine regulation of pituitary function. The two molecular forms of IL-1: IL-1 α and IL-1 β affect the CNS and the endocrine system (Schmidt and Tocci, 1990), in addition, IL-1 and IL-6 (Sapolsky *et al.*, 1987; Battman *et al.*, 1989 and Tsagarakis *et al.*, 1989) are known to be stimulants of the hypothalamic-pituitary axis while TNF (cachectin) is known to inhibit the secretion of pituitary hormones such as growth hormone (Walton and Cronin, 1989) and TSH (Pang *et al.*, 1989). Pang *et al.* (1989) implicated TNF in causing impairment of hypothalamic-pituitary-thyroid function in rats and suggested that TNF is one of the mediators responsible for alterations in the thyroid function. On the other hand, thyroid-stimulating hormone concentrations decline following subcutaneous injection of recombinant human IL-1 β in rats (Dubois *et al.*, 1988). The weakness that has been pointed to in the study implemented by Abebe and Eley (1992) and Al-Qarawi *et al.* (2001) in their failure to study the pituitary TSH output has been our target to strengthen the present study and achieve our conclusion which indicated a dysfunction of the servomechanism in the pituitary-thyroid axis during the state of larvae of *C. titillator* infection in camel. The alteration in the T3 and T4 dynamics in the study of Al-Qarawi *et al.* (2001), in the camels infected with *Trypanosoma evansi* was not consistent with the results of Lomo *et al.* (1996) in the rabbits infected with *Trypanosoma congolense*. The depression in the T3/T4 ratio in the previous study of Al-Qarawi *et al.* (2001), may be due to the release of T3 by the thyroid gland under the influence of small amounts of pituitary hormones (Nejad *et al.*, 1973). According to the above developments, it is evident that infection with larvae of *C. titillator* would tend through their multiple functions to cause decrease in thyroid hormones synthesis which might explain the decrease in T3 and T4 during the experiment. Also the concomitant decrease in blood TSH level might as well as stimulate the decline of the T3 and T4 in blood.

لم ج ل ي ف ي ق ي ر د ل ا - ة ي م ا خ ن ل ا د غ ل ا ر و ح م ي ف ت ا ر ي غ ث ل ا
 ة ب ا ص ل ا ة ج ي ت ن س ي ر ا د ي م و ر د س ي ل ي م ا ك
 ا ن ي ب و ل ا ق ي س ل م ج ل ا ف ا غ ن ة ب ا ب ذ ت ا ق ر ي ب
 ر و ت ا ل ي ت ي ت
 ء ا م س ا و ³ ي ل ا د ل ا ت ز ع ، ² ر ي ق ي ص ل ا ن ا م ث ع ، ¹ ا ي ن و ي س ب ل ا د ا ع
 ل ا م ك ⁴

- 1- ر ص م - ا ز ي ق ل ا - ا ن و ا ق ل ا ا ع ر ا ج م و ل ع ل ا ي ل ك - ت ا ر ش ح ل ا د ا م س ق
- 2- ي ب و ر ل ا د ل م م د ل ا م ي س ق ل ا م ي س ق ل ا ا ع ر ا ج م و ل ع ل ا ي ل ك - ا ي ب ط ل ا ة ي ق ي ر د ل ا ء ا ي ا ل ا م س ق
- 3- ر ص م - ا ن و ا ق ل ا - ن ا و ل ح ا ع ر ا ج م و ل ع ل ا ي ل ك - ت ا ر ش ح ل ا ر ا و ي ح ل ا م ل م س ق
- 4- ر ص م - ا ز ي ق ل ا - ي ق ل ا - ا ي ع ا ر ز ل ا ت و ر ح ل ا ز ه ر م - ن ا و ي ح ل ا ا ع م ش و ا ب د د م

د غ ل ا ر و ح م ي ف ت ا ر ي غ ث ل ا ت ا ر ي غ ث ل ا ت ح ب ل ا ا ذ ه ي ف ة ب ا و د ل ا ت ل و ا ن ت
 ل م ج ل ا ف ا غ ن ة ب ا ب ذ ت ا ق ر ي ب ل ا ا ل ا ر و ط ل ا ب ة ب ا ص ل ا ء ا ن ش ا ة ي ق ي ر د ل ا - ة ي م ا خ ن ل ا
 ي ا ر ت ل ا ن م ل ك ي و ت س م س ا ي ق ب ل ك ل ذ ر و ت ا ل ي ت ي ت ا ن ي ب و ل ا ف ي س
 ا ض ي ا و م د ل ا ي ف ي ق ي ر د ل ا د غ ل ل (T4) ن ي س ك و ر ي ث ل ا و (T3) ن ي ن و ر ي ث و د ي ا
 (TSH) ة ي ق ي ر د ل ا د غ ل ل ش ا ح ل ا ة ي م ا خ ن ل ا د غ ل ا ن و م ر ه
 ت د ا ل م ج ل ا ف ا غ ن ة ب ا ب ذ ت ا ق ر ي ب ل ا م ج ل ا ة ب ا ص ل ا ن ا ن ع ج ئ ا ت ن ل ا ت ف ش ك
 د غ ل ا ب ن ي ص ا ل ا ن ي ن و م ر ه ل ا ن م ل ك ي و ت س م ي ف ض ر ا ف ا خ ن ا ت و د ح ي ل ا
 T3 و T4 ا ن و م ر ه ا م و م د ل ا ي ف ي ق ي ر د ل ا
 ي ف ي م ا خ ن ل ا د غ ل ل TSH ن و م ر ه ي ف ص ر ق ن ت و د ح ج ئ ا ت ن ل ا ت ن ي ب ا ض ي ا
 م د ل ا
 ة ي ق ي ر د ل ا د غ ل ل ش ا ح ل ا ن و م ر ه ل ا ق ا ل ط ن ا ي ف ص ر ق ن ل ا ن ا ج ئ ا ت ن ل ا ر ه ط ت
 ي ف ب ح ا ص ل ا م ص ر ق ن ل ا م ج ل ا ي ف ء ر ش ح ل ا ت ا ق ر ي ب ة ب ا ص ل ا ء ا ن ش ا TSH
 د غ ل ا ل ع ة ب ا ص ل ل ر ش ا ب م ل ر ي ث ا ت ل ا س ن ك ع ي ا م ب ر T4 و T3 ي و ت س م
 ر و ط ل ا ب ة ب ا ص ل ا ن ا ا ض ي ا ج ئ ا ت ن ل ا ح ر ت ق ت ا م ك ، ة ي ق ي ر د ل ا د غ ل ا و ة ي م ا خ ن ل ا
 T3 ن م ل ك ن ي و ك ت ي ف ص ر ق ن ب ب س ت ل م ج ل ا ف ا غ ن ة ب ا ب ذ ت ا ق ر ي ب ل ا ا ل ا
 و T4

References

Al-Qarawi A. A., H. Abdel-Rahman and S. A. El-Mougy. 2001. Impairment in the pituitary-thyroid axis of the *Camelus dromedarius* infected with *Trypanosoma evansi*. Dtsch. Tierarztl. Wschr., 108: 172-174.
 Battman A., A. T. Singh and S. Solomon. 1989. The immuno-hypothalamic pituitary adrenal axis. Endocrinological Reviews, 10: 92-112.

- Boersma A., M. Hublart and F. Boutignon. 1989. Alterations in thyroid function in patients with *Trypanosoma brucei gambiense* infection. Transactions of the Royal Society of Tropical Medicine and Hygiene, 83: 208-209.
- Higgins A. J., 1985. The camel in health and disease. 4. Common ectoparasites of the camel and their control. Br. Vet. J., 141: 197-216.
- Mutayoba B. M., H. B. O'Hard-Iriri, and S. Gombe. 1988. Trypanosome-induced depression of plasma thyroxine levels in prepubertal and adult female goats. Acta Endocrinologia (Copenhagen), 199: 21-26.
- Ikeke B. O. , and G. J. Losos.,1975. Pathogenesis of *Trypanosoma brucei* infection in sheep. Hypophyseal and other endocrine lesions. J. Comparative Pathology, 85: 37-44.
- Roncarì D. A. K. and V.K. Murthy. 1975. Effect of thyroid hormones on enzymes involved in fatty acids and glycerolipid synthesis. J. Biol.Chem., 250: 4134-4138
- Lonsdale-Eccles D. J. and Grab, 1986. Cysteine proteinases and their inhibitors, ed. V. Turk, P. 189, Berlin, Walter de Gruyter
- De Souza E. B. , E. L. Webser, D. E. Grigoriadis and E. D. Tracey, 1989. Corticotrophin releasing factor (CRE) and interleukin-1 (1L-1) receptors in the brain-pituitary immune axis. Psychopharmacology Bulletin, 25: 299-305
- Haroun E. M., M. Magzoub, O. M. Mahmoud and O. H. Omer, 2000 Some studies on trypanosomiasis in Najdi camels in AlQassim region. 20th Annual Meeting of the Saudi Biological Society, 1-3, King Fas
- Al-Ani F. K. , W. A. Khamas and M. R. Al-Shareefi, 1991. Camel nasal myiasis: clinical, epidemiological and pathological studies in Iraq. Indian Journal of Animal Sci., 61: 576-578
- Derhalli F. S. , G. A. Zakia, E. L. El-Ramadan and A. M. Abdel Gwad, 1989. Serological diagnosis of *Cephalopina titillator* infestation of camel. J. Egypt. Vet. Med. Ass., 49: 541-548
- Bray G. A. and H. M. Goodman, 1968. Metabolism of adipose tissue from normal and hypothyroid rats. Endocrinol., 82: 865
- Abebe G. and R. M. Eley, 1992 .Trypanosome induced hypothyroidism in cattle. British Veterinary J., 148, 63-70
- Abebe G., M. K. shaw and R. M. Eley, 1993. *Trypanosoma congolense* in the microvasculature of pituitary gland of experimentally infected Boran cattle (*Bos indicus*): Veterinary Pathology, 30: 401-409
- Knowles G. , G. Abebe and S. J. Black, 1989. Detection of parasite peptidase in the plasma of heifers infected with *Trypanosoma congolense*. Molecular Biochemistry and Parasitology, 34: 25-34
- Nejad I. F. , J. A. Bollinger, M. Mitrick, and S. Reichlin, 1973. Importance of T3 (triiodothyronine) secretion in altered states of thyroid function in the rat: cold exposure, subtotal thyroidectomy and hypophysectomy. Transactions of the Association of American Physicians, 85: 295-308
- Tizard I. , K .H. M. Nielsen, J. R. Seed, J. E. Hall, 1978. Biologically active products from African trypanosomes. Microbiological Reviews, 42: 661-681
- Honegge J. r, A. Spagnoli, R. Durso, P. Navarra, S. Tsagarakis, G. M. Besser and A. B. Grossman, 1991. Interleukin-1 β modulates the acute release of growth hormone-releasing hormone and somatostatin from rate hypothalamus in vitro, whereas tumor necrosis factor and interleukin-6 have no effect. Endocrinology, 129: 1275-1282
- Dubois J. M., J. M. Dayer, C. A. Siegrist-Kaiser and A. G. Burger, 1988. Human recombinant interleukin-1 β decreases plasma thyroid hormone and thyroid stimulating hormone levels in rats. Endocrinology, 123: 2175-2181
- Fisher J. N. and E. G. Bull. 1967. Studies on the metabolism of adipose tissue. xx. The effect of thyroid studies upon oxygen consumption and lipolysis. Biochemistry (wash) 6: 637-647
- Sharma L. K., 1992. Efficacy of ivermectin against nasal bots in camels (*Camelus dromedarius*). Indian Veterinary Journal, 69: 835-836.
- Hussein M. F. , F. M. El-Amin, N. T. El-Taib and S. M. Easmaeil, 1982. The pathology of nasopharyngeal myiasis in Saudia Arabian camels (*Camelus dromedarius*). Vet. Parasitol., 9: 253-259
- Hussein M. F. , H. A. R. Hassan, H. K. Bilal, S. M. Younis, A. A. R. Al-Motlaq and M. A. Al-Sheikh, 1983. *Cephalopina titillator* (Clark, 1797) infection in Saudi Arabian camels. Zentralblatt Fur Veterinarmedizin, B30: 553-558
- Humblart M. , D. Tetaert, D. Crolx, F. Boutignon, F. Degand, A. Boersma, 1991. Gonadotrophic dysfunction produced by *Trypanosoma brucei* in the rat. Acta Tropica, 4: 177-184
- Musa M. T. , M. Harrison, A. M. Ibrahim and T. O. Taha, 1989. Observations on Sudanese camel nasal myiasis caused by the larvae of *Cephalopina titillator*. Revue d'Elevage et de Medecine Veterinaire des Pays Tropicaux, 42: 27-31
- Walton P. E. and J. M. Cronin, 1989. Tumor necrosis factor inhibits growth hormone secretion from cultured anterior pituitary cells. Endocrinology, 125: 925-929
- Lomo P. O. , D. W. Makawiti and V. N. Konji. 1996. Thyroid status and adenosine triphosphatase activity in experimental *Trypanosoma congolense* infection in rabbits. British Veterinary J., 152: 659-667
- Sapolsky R. , C. Rivier, G. Yamamoto, P. Plotsky, and W. Vale, 1987. Interleukin-1 stimulates the secretion of hypothalamic corticotrophin-releasing factor. Sci., 238: 522-526
- Tsagarakis S. , G. Gillies, L. H. Rees, M. Besser, A. Grossman. 1989. Interleukin-1 directly stimulates the release of corticotrophin releasing factor from rat hypothalamus. Neuroendocrinology, 49: 98-101
- Schmid T. A. t and M. J. Tocci. 1990. "Interleukin-1" in peptide growth factors and their receptors 1. Spom, M.B.; Roberts, A.B. (eds): Springer Verlag, New York, p: 473
- Bekele T., 2001. Studies on *Cephalopina titillator*, the cause of "Sengale" in camels (*Camelus dromedarius*) in semi-arid areas of Somali State, Ethiopia. Tropical Animal Health and Production, 33: 489-500
- Pang X. P. , J. M. Hershman, C. J. Mirell, and A. E. Pekary. 1989. Impairment of hypothalamic-pituitary-thyroid function in rats treated with human recombinant tumor necrosis factor (Cachectin). Endocrinology, 125: 76-84