

## Some Descriptive Characteristics Concerning Craniometric Measurement in Kangal Dog

<sup>1</sup>Irfan Daskiran and <sup>2</sup>Fatin Cedden

<sup>1</sup> Department of Animal Science, Faculty of Agriculture, Yüzüncüyıl University, Van, Turkey

<sup>2</sup> Department of Animal Science, Faculty of Agriculture, Ankara University, Ankara, Turkey

**Abstract:** In this study, a total of 38 adult female and male Kangal dogs were used for craniometric measurements. Average facial width was found  $13.09 \pm 0.202$  mm for females,  $14.84 \pm 0.197$  mm for males ( $p < 0.01$ ). Average facial and skull index were found as  $122.22 \pm 3.709$  mm,  $55.213 \pm 0.984$  mm for females,  $134.26 \pm 3.470$  mm and  $60.14 \pm 1.080$  mm for males, respectively. Body weights for females and males were  $34.27 \pm 0.944$  kg and  $43.43 \pm 1.330$  kg. The ratio of skull length to facial width was 1.0:0.55 for females, 1.0:0.60 for males. In conclusion, Kangal dogs can be considered in dolichocephalic group, according to the value of skull length/ facial width ratio.

**Key words:** Kangal dog, Craniometric measurement, descriptive characteristics

### INTRODUCTION

Sheep rearing concerns the largest part of animal production sector in Turkey, due to 26 millions of sheep<sup>[1]</sup>. Sheep flocks are usually kept out-door in pasture based feeding during grazing period. In many sheep breeding areas, the basic problem for safety of flocks are predators. It is possible to protect sheep herd against predators about up to 50-80% of safety in grazing areas<sup>[2]</sup>.

The Kangal dog is one of the famous Anatolian sheep dog with its size, courage and mastiff-like character.

The dogs being raised from Kangal district of Sivas province in Central Anatolia have typical and most appreciable characteristics type of Kangal dog.

The earliest apparition of Kangal like sheep dogs was at 2000-1800 before Christ, during Hittite civilization in Anatolia<sup>[3]</sup>. It was mentioned of a dog strain which was very similar to Kangal dog of today, in ancient Tibetan reliefs from the period 4000 before Christ<sup>[4]</sup>. It is thought to be that ancient settlers migrated from Asia, might have brought their sheep and hunt dogs to Anatolia. Especially, Central Asian tribes migrated in 1000 after Christ, might have played a particular role in apparition of different sheep dogs in Anatolia.

The objective of this study is to investigate some skull measurement characteristics in Kangal sheep dogs.

### MATERIALS AND METHODS

**Material:** In this study, a total of 22 female and 16 male Kangal dogs were used. The dogs were 1-5 years-old and

bred in Ulas State Farm, situated in Sivas province, Turkey. Dogs were fed daily with cereal-based mixture and weekly slaughtering houses remainders.

**Method:** Craniometric measurements were taken by using the method described by Stockard<sup>[5]</sup>. Craniometric measurements and indices were obtained as follows (Fig. 1 and 2):

**Skull length:** Rostral point of sutura interincisiva-protuberantia occipitalis externa

**Maximum zygomatic (facial) width:** Left arcus zygomaticus-right arcus zygomaticus

**Facial length:** Frontal midpoint-rostral point of sutura interincisiva

**Skull index:** Maximum zygomatic width x 100/Skull length

**Facial index:** Maximum zygomatic width x 100/Facial length

Data were analysed by Minitab package programme<sup>[6]</sup>.

### RESULTS

Average skull measurements of female and male Kangal dogs are shown in Table 1. Average skull measurement values, such as facial and skull length and skull indices do not vary significantly according to gender of adult dogs, except for facial width which was found

Table 1: Average skull measurements in female and male Kangal dogs ( $\bar{X} \pm S\bar{X}$ )

| Characteristics      | Female (n:22) | Male (n:16)   |
|----------------------|---------------|---------------|
| Facial length(cm)    | 10.841±0.243  | 11.156±0.298  |
| Facial width(cm)     | 13.091±0.202  | 14.844±0.197* |
| Skull length(cm)     | 23.773±0.308  | 24.750±0.374  |
| Facial index         | 122.220±3.709 | 134.260±3.470 |
| Skull index          | 55.213±0.984  | 60.140±1.080  |
| Body weight(kg)      | 34.227±0.944  | 43.410±1.330  |
| General skull index  | 57.288±0.823  |               |
| General facial index | 127.290±2.740 |               |

P<0.01

Table 2. Average skull measurements by ages in Kangal dogs ( $\bar{X} \pm S\bar{X}$ )

| Characteristics   | Age groups    |               |               |               |               |
|-------------------|---------------|---------------|---------------|---------------|---------------|
|                   | 1<br>n:4      | 2<br>n:13     | 3<br>n:5      | 4<br>n:6      | 5<br>n:10     |
| Facial length(cm) | 10.625±0.239  | 11.077±0.383  | 10.900±0.400  | 10.667±0.357  | 11.200±0.436  |
| Facial width(cm)  | 14.000±0.408  | 13.346±0.402  | 13.900±0.557  | 13.583±0.327  | 14.500±0.365  |
| Skull length(cm)  | 23.250±0.479  | 24.038±0.369  | 24.200±1.020  | 23.833±0.477  | 24.950±0.519  |
| Facial index      | 132.090±6.030 | 122.100±5.510 | 127.900±5.050 | 128.090±5.290 | 131.330±6.170 |
| Skull index       | 60.240±1.670  | 55.620±1.720  | 57.500±1.110  | 57.020±1.040  | 58.330±1.880  |
| Body weight(kg)   | 38.250±2.690  | 35.770±1.850  | 37.800±3.580  | 36.170±1.220  | 42.350±2.160  |




Fig. 1. Lateral view of Kangal dog skull and measurement points

- a: Rostral point of sutura interinciviva
- b: Frontal midpoint
- c: Protuberantia occipitalis externa




Fig. 2: Dorsal view of Kangal dog skull and measurement points

13.09 ± 0,202 mm for females, 14.84 ± 0.197 mm for males (p<0.01) respectively. Average facial and skull index were

found as 122,22±3,709 mm and 55.213±0,984 mm for females, 134.26±3.470 mm and 60.14±1.080 mm for males respectively. Average body weights for females and males were found as 34.27±0.944 kg and 43.41±1.330 kg, respectively.

The distribution of average skull measurements by ages, are presented in Table 2. These values do not vary significantly between age groups in Kangal dog.

## DISCUSSION

In a study concerning skull typology, Kangal dogs were considered mesaticephalic, like German Shepherd dogs<sup>[7]</sup>. In same study, average facial and skull lengths in Kangal dogs of 1, 2 and 3 years-old were found as 13.31 and 26.38, 31.34 and 26.61, 13.45 and 28.99 cm, respectively. Average values of facial length were found lower in our study. Onar *et al.*,<sup>[8]</sup> have obtained average values in macerated adult Kangal skulls as follows: Skull length, 25.87±13.375 cm; facial length, 13.00±5.404 cm; skull index, 50.29±1.033 and facial index 99.62±3.891. In our study, the ratio of skull length to facial width was 1.0:0.55 for females, 1.0: 0.60 for male Kangal dogs. Zietzschmann<sup>[9]</sup> had reported that dogs described in dolichocephalic group have the ratio of skull length to facial width as 1.0:0.60-0.65. Onur *et al.*,<sup>[8]</sup> have found this ratio as 1.0: 0.50. Same authors have considered Kangal dog as dolichocephalic race, according to the values of skull measurement which were found noticeably higher than these obtained from mesocephalic breeds. Therefore, the values obtained from our study were very close to the other researchers findings. The facial and skull indices in Kangal dogs of 1, 2 and 3 years-old were found 92.8 and 46.8, 96.5 and 48.4, 108.7 and 50.4, respectively<sup>[2]</sup>. The values of facial index were found higher in the study

conducted by Yildiz *et al.*,<sup>[2]</sup> compared to our findings. Generally, a higher value of facial length was not appreciated by Kangal dog breeders, because of a harmony between facial and skull length is preferred in dogs. In practice, the preference of Kangal breeders is as 1/3 for facial/skull length ratio in Kangal dogs.

#### REFERENCES

1. Anonymous, 2001. Tarımsal Yapı ve Üretim. T.C. Bařbakanlık. D.İ.E. Yay. Ankara
2. Green, J.S. and R.A. Woodruff, 1985. Summary of Livestock guarding dog research at the U.S. Sheep Exp. Station. Sheep Production.
3. Kırmızı, E., 1994. Türk Çoban Köpeđinin Tariřesi. Türk Veteriner Hekimleri Dergisi, 1, 6.
4. Sefton, F., 1969. Complete Dog Guide, Pet Library, The Netherlands, 11-28.
5. Stockard, C.R., 1941. The genetic and Endocrinic Basis for Differences in Form and Behavior. American Anatomical Memoirs. No: 19, Philadelphia: Wistar Institute, 1-775.
6. Minitab, 1993. Minitab Reference Manuel Released 9 for Windows, July 1993 by Minitab inc.
7. Yıldız, B., O. Yılmaz, A. Serbest and H. Kırbıyk, 1993. Türk Çoban ve Alman Kurt Köpeklerinin Bař Ölçümleri Üzerinde Arařtırma. U. Ü. Vet. Fak. Dergisi, 1, 12.
8. Onar, V., S. Ozcan and G. Pazvant, 2001. Skull Typology of Adult Kangal Dogs. Anat. Hist. Embr. Blackwell Wissenschafts-Verlag, Berlin. 30: 41-48.
9. Zietzschmann, O., 1943. I. Das Skeletsystem. Der passive Bewegungsapparat. In: Handbuch der Vergleichenden Anatomie der Haustiere, 18 Auflage (W. Ellenberger and H. H. Baum) Berlin. 114-116