

Seroprevalence Studies on EGG Drop Syndrome-76 Virus in Southren Part of India (Kerala)

¹ A.M. Arun, ²Mini, ³Leo Joseph and ⁴V. Jayaprakasan

^{2,4}Department of Microbiology, College of Veterinary and Animal Sciences, Mannuthy, Thrissur, India

³University Poultry and Duck Farm, College of Veterinary and Animal Sciences, Mannuthy, Thrissur, India

Abstract: The present seroprevalence of EDS-76 by HI test revealed widespread distribution of EDS-76 antibodies in layer chicken in Kerala, India. Out of 615 sera samples tested by HI test, 58 samples found to be positive for EDS-76 virus antibodies. The overall prevalence of EDS-76 antibodies in Kerala was 9.43 % The attempts for virus isolation from clinical materials were failed.

Key words: Seroprevalence, EGG drop syndrome-76, haemagglutination inhibition test

INTRODUCTION

EGG drop syndrome-76 is an economically important viral disease of laying hens, characterized by a drop in EGG production and EGGshell defects and the virus can be transmitted through eggs^[1]. Egg drop syndrome-76 virus belongs to the family *Adenoviridae* and is classified under the genus, *Aviadenovirus*. Mohanty *et al.*^[2] conducted a serological survey of layer flocks in different parts of India during 1980 to 1983 and reported EDS-76 as a widespread condition associated with fall in EGG production. In Kerala antibodies against EDS-76 were found in pigeons and house crows^[3] duck and chicken^[4,3] by Haemagglutination (HI) test. The present study was conducted to estimate the seroprevalence of EDS-76 among layer chicken in Kerala.

The EDS-76 virus (strain 127 of UK), which was maintained in Department of Microbiology, College of Veterinary and Animal Sciences, Thrissur, Kerala was used as a reference virus. The EDS-76 virus grown in embryonated duck EGGs was used as antigen for raising hyperimmune serum. The antisera were prepared as per the method of Mohanty *et al.*^[2] with the slight modifications. Antisera were heat inactivated at 56° C for 30 min and stored at -20° C.

Sera samples were collected randomly from layers at nine organized layer poultry farms situated in various districts of Kerala. A total of 615 sera samples were collected randomly from apparently healthy layers, irrespective of age. Sera samples were screened for EDS-76 by HI test and the test was carried out as per the method^[5].

Table 1: Seroprevalence of EDS-76 virus in chicken in kerala

Farm	Total number of samples screened	HI	
		Number positive	Percentage positive
RPF*, Malampuzha	28	3	10.71
RPF*, Kodapanakunnu	20	2	10.00
KSPDC**, Kodapanakunnu	52	3	5.77
RPF*, Koovapadi	68	9	13.23
UPF*, Mannuthy	105	15	14.28
RPF*, Chathamangalam	65	8	12.31
RPF*, Mundayad	80	6	7.5
RPF*, Manarkad	95	6	6.31
Central Hatchery, Chengannoor	98	4	4.08
Department of Microbiology	4	2	50
Total	615	58	9.43

* Regional poultry farm ** Kerala state poultry development corporation
 # University Poultry Farm

The present investigation revealed the presence HI antibodies against EDS-76 virus in Kerala. The HI antibody titre of 8 (3 log₂) or above was considered as positive in the present study. A total of 58 out of 615 samples found to be positive for EDS-76. The overall prevalence of EDS-76 antibodies in Kerala was 9.43% as against 14.91 reported earlier^[4]. The farm wise incidence of seropositive samples ranged from 4.08 to 14.28% and is shown in Table 1.

Attempts for isolation of EDS-76 virus were not successful. Since no vaccination was carried out in most

Corresponding Author: V. Jayaprakasan, College of Veterinary and Animal Sciences, Kerala Agricultural University Mannuthy, Thrissur, Kerala, 680651, India

of the farms, the failure of virus isolation and the presence of antibodies indicated inapparent infection of EDS-76 virus among layer chicken in Kerala. The present study suggested the widespread infection of EDS-76 in chicken in Kerala and vaccination of the chicken is warranted.

REFERENCES

1. McFerran, J.B., R.M. McCracken, E.R. McKillop, M.S. McNulty and D.S. Collins, 1978. Studies on a depressed EGG production syndrome in Northern Ireland. *Avian Pathol.*, 7: 35-47.
2. Mohanty, G.C., K.C. Verma, H.K. Pradhan and R. Kumar, 1984. EGG drop syndrome-76 (EDS-76) in India, Seroprevalence of EDS-76 virus infection in poultry flocks. *Indian J. Poultry Sci.*, 19: 15-18.
3. Sulochana, S. and D. Sudharma, 1987. Seroprevalence of EGG drop syndrome virus infection in domestic and free flying birds in Kerala. *Kerala J. Vet. Sci.*, 18: 83-88.
4. Priya, P.M., G.K. Nair, M. Mini and V. Jayaprakas, 2001. Comparison of haemagglutination inhibition test and indirect elisa for detecting EDS-76 antibodies in chicken and ducks. *Indian J. Poultry Sci.*, 36: 326-328
5. McFerran, J.B., H.M. Rowley, M.S. McNulty and L.J. Montgomery, 1977. Serological studies on flocks showing depressed EGG production. *Avian Pathol.*, 6: 405-413.