

Sero-Surveillance of Hemorrhagic Septicemia in Cattle and Buffalo in Malakand District of NWFP, Pakistan

Asadullah Khan, ⁴Riaz Ahmad, ²Umar Saddique, ²Hamayun Khan,
¹Mohammad Zubair and ³Younas Mohammad

¹Office of the Deputy Director Artificial Insemination, Directorate of Livestock and Dairy Development Peshawar

²Department of Animal Health, NWFP Agricultural University Peshawar

³Civil Veterinary Hospital Julagram District Malakand Agency

⁴London Business Accademy London, UK

Abstract: The present study showed the Sero-surveillance of Hemorrhagic septicemia in cattle and buffalo in Malakand division of NWFP, Pakistan. The average GMT was recorded against hemorrhagic septicemia in buffaloes in the range of 4.12 to 46.98, while those in cattle were recorded in the range of 4.45 to 46.40. In young buffalo calves, incidence rate, mortality and morbidity rate was observed as 22.25, 21.19 and 95.25%, respectively. In adult buffalo morbidity, mortality and case fatality rate was 5.49, 1.65 and 30%, respectively. In study of young cattle calves morbidity, mortality and case fatality was recorded as 3.94, 1.77 and 45%, respectively. While in study of adult cattle, morbidity, mortality and case fatality rates were recorded as 2.51, 0.39 and 15.79%, respectively. During present investigation the incidence rate, mortality and case fatality due to Hemorrhagic septicemia was found greater in the young calves as compared to the adult in both buffalo and cattle.

Key words: Sero-surveillance, hemorrhagic septicemia, cattle, buffalo, malakand

INTRODUCTION

Hemorrhagic Septicemia (HS) is an acute pasteurellosis caused by particular serotypes of *Pasteurella multocida* and manifested by an acute and highly fatal septicemia principally in cattle and water buffaloes; the latter are thought to be more susceptible. It has been reported in bison, camels, elephants, horses and donkeys and there is evidence of its occurrence in yak. HS is a major disease of cattle and water buffalo in Asia, Africa and some countries of southern Europe and the Middle East. Although it may be seen at any time of year, the worst epidemics occur during the rainy season. It is most common in the river valleys and deltas of Southeast Asia among buffaloes used in rice cultivation.

Study on the seroprevalence on Hemorrhagic septicemia has been carried out in other region of the world as well as some area of the Punjab in our country⁽¹⁻⁴⁾.

Keeping in view above mentioned facts the present study was undertaken to collect a data regarding the severity of the Hemorrhagic Septicemia (HS) disease in the region and to know exactly about the strength of Hemorrhagic Septicemia (HS) antigen in the blood of the

animal either vaccinated or unvaccinated. Further more the finding of the present study would be helpful to chalk out the strategy for precise control and prevention of the disease.

MATERIALS AND METHODS

The present study was conducted to find out the seroprevalence of Hemorrhagic septicemia in cattle and buffalo in Malakand district of NWFP, Pakistan.

Experimental animals: Blood sample were collected in a test tube from both ten healthy and ten infected animals (five each vaccinated against hemorrhagic septicemia and five each were un-vaccinated). Six milliliter of blood was collected from each animals and were kept overnight at room temperature in slanting position. The sera was separated through centrifugation 3000 rpm for 10 min and stored at -20°C for further studies.

Laboratory assay: Antibodies against the *Pasteurella multocida* were measured by using classical Indirect Haemagglutination Test (IHA) using human blood O⁽⁵⁾.

Corresponding Author: Asadullah Khan, Office of the Deputy Director Artificial Insemination, Directorate of Livestock and Dairy Development Peshawar

Test procedure: The test was carried out in microtitre plates of Flow Laboratories, each having 96 U shaped wells, arranged in 8 rows and 12 columns designated as A-H and 1-12, respectively. Two fold dilutions of the test sera starting from 1:5 to 1:640 were made in normal saline solution and added in 25 µL amounts to all the wells of plate except those of column 11 and 12 which were maintained as controls. First four wells (A-D) of column 11 were added with known negative serum and last four wells (E-H) with the known positive serum. All the wells of the column 12 were added with normal saline solution. Sensitized RBC's (1%) were added in equal amounts (25 µL) to all the wells of the plate, so that column 12 served as control for the RBC's. The plates were incubated at room temperature for two hours and the observations recorded. Thereafter the plates were kept under refrigeration for overnight shake lightly, allowed to resettle and read again. Results were interpreted as under:

- Positive:** No button formation, clumping occurring in an unordered and ragged pattern.
Negative: Button formation, RBC's clumping in an organized and regular pattern.

CALCULATION OF GMT

The GMT was calculated by the following formula:

$$GMT = n\sqrt{X_1 \cdot X_2 \dots X_n}$$

n = where n is the number of titre
 x = the value of titre

RESULTS

Hemorrhagic Septicemia (HS)

Buffalo and cattle: The total cattle and buffalo population in selected villages/town was 4371. Out of which young male and female livestock were 731 and 1037, while the adult male and female livestock were 416 and 2187, respectively. The total population of the young male and female buffalo was 313 and 442, while the adult male and adult female were 182 and 909, respectively Table 1. The population of the total young male and female cattle was 418 and 595, while the adult male and female cattle were 234 and 1278, respectively Table 1. Table 2 showed the average GMT in vaccinated, unvaccinated, recovered and affected buffaloes in Malakind District in the range of 4.12 to 46.98.

Table 3 showed the average GMT in vaccinated, unvaccinated, recovered and affected cattle in Malakind District in the range of 4.45 to 46.40.

Table 1: Total livestock population in selected villages/town of malakand district

Animals	Male		Female		Total
	Young	Adult	Young	Adult	
Buffalo	313	182	442	909	1846
Cattle	418	234	595	1278	2525
Total	731	416	1037	2187	4371

Table 2: Calculation of Geometric Mean Titre (GMT) against hemorrhagic septicemia in buffaloes through Indirect Haemagglutination Assay (IHA) in various classified areas of district malakand

Animal condition	Town	Big villages	Small villages	Average GMT
Affected	3.99	4.19	4.18	4.12
Recovered	63.99	73.51	55.71	64.41
Unvaccinated	13.92	18.38	12.13	14.81
Vaccinated	55.71	48.48	36.75	46.98

Table 3: Calculation of Geometric Mean Titre (GMT) against hemorrhagic septicemia in cattle Haemagglutination Assay (IHA) in various classified areas of district malakand

Animal condition	Town	Big villages	Small villages	Average GMT
Affected	5.27	3.48	4.59	4.45
Recovered	63.97	55.69	48.48	56.05
Unvaccinated	10.55	9.19	15.99	11.91
Vaccinated	42.22	48.48	48.50	46.4

Table 4: Annual incidence, mortality and case fatality rates due to HS in buffalo and cattle population in 10 randomly selected villages/town of district malakand

Animal	Age group	Pop "n"	AA "n"	DA "n"	MB %	MR %	CF
Buffalo	Young*	313	42	42	13.42	13.42	100
	Adult*	182	12	-	6.59	-	-
	Young**	442	126	118	28.5	26.69	93.65
Total	Adult**	909	48	18	5.28	1.98	37.5
	Young	755	168	160	22.25	21.19	95.23
	Adult	1091	60	18	5.49	1.65	30
Cattle	Young*	418	18	7	4.31	1.67	38.8
	Adult*	234	02	-	0.85	-	-
	Young**	595	22	11	3.69	1.85	50
Total	Adult**	1278	36	6	2.82	0.47	16.6
	Young	1013	40	18	3.94	1.77	45
	Adult	1512	38	06	2.51	0.39	15.79

Pop = Population * = Male, AA = Affected Animal, ** = Female, DA = Died Animal, MB = Morbidity, MR = Mortality, CF = Case Fatality

In young buffalo calves, incidence rate was observed as 22.25% along with 21.19 and 95.25% mortality and case fatality, respectively. In adult buffalo morbidity, mortality and case fatality rate was 5.49, 1.65 and 30%, respectively Table 4.

Total population of young cattle was 1013. Out of which morbidity, mortality and case fatality was recorded as 3.94, 1.77 and 45%, respectively while in study of total population of adult cattle was 1512, morbidity, mortality and case fatality rates were 2.51, 0.39 and 15.79%, respectively Table 4.

DISCUSSION

Hemorrhagic Septicemia (HS): Haemorrhagic Septicaemia (HS) is one of the most important diseases

of bovines in South Asian and Middle Eastern countries. Epidemiological studies in neighboring country was reported over a period of thirteen years (1974-1986) in India indicated that mortality-wise, H.S. was placed first and morbidity-wise, second as compared to four other epizootic diseases namely, foot and mouth disease, rinderpest, anthrax and black quarter^[2]. Many states in India were marked as high risk zones. About 26 outbreaks have been recorded in Punjab State from 1989 to 1990 Saini^[6].

Similarly, Outbreaks of Haemorrhagic Septicaemia (HS) of have been recorded in Srilanka^[7,8], South Asia, The Middle East and Africa^[9]. The morbidity rate reported was 6.40 and the mortality rate was 6.28 per lakh of bovine population^[2], Mortality rates range has a widely from 5 to 90% in India, Nepal and the Philippines in different outbreaks and seasons^[10]. Most of the outbreaks have been managed by medical treatment alone and resulted in poor survival rate.

These findings are accordance with the study of Shahid^[11] who reported mortality (23.13%), morbidity (21.71%) and case fatality (93.89%), respectively. The case fatality of young buffaloes in this survey was 95.23% who observed 95.45% case fatality in young buffaloes. The result obtained in present study regarding the mortality, morbidity and case fatality are in close vicinity with the finding of Sheikh^[12] who carried out study on observation on hemorrhagic septicemia in nine districts of Punjab, Pakistan showed 11% incidence, 9% mortality and 78% case fatality rates of hemorrhagic septicaemia in buffalo, whereas these values were 4, 2.5 and 62% in cattle.

Morbidity, mortality and case fatality in young cattle was 3.94, 1.77 and 45%, respectively, while in adult cattle these values recorded were as 2.51, 0.39 and 15.79%, respectively. These findings are accordance with the study of Shahid^[11] who recorded the morbidity, mortality, case fatality in young cattle was 3.52, 1.76 and 50%, respectively, while in adult cattle these records were 2.1, 0.36 and 17.50%, respectively. As concerned the morbidity of young cattle in this study in partial agreement with result obtained by Ibrahim^[13] who recorded 2.45%, respectively.

The finding observed during the present study regarding the mortality, morbidity and case fatality in both buffaloes and cattle were in agreement with the finding of Saini^[6] who reported 19.8% mortality in buffalo and 23.7% mortality in cattle, respectively. De-Alwis and De-Alwis^[14] reported higher mortality in buffaloes (45.2%) as compared to cattle (15.8%) in Sri Lanka due Hemorrhagic Septicemia (HS).

From this study it is concluded that the occurrence of the disease was higher in buffaloes when compared to cattle in the present study. Hemorrhagic Septicemia (HS) is endemically occurring in cattle and buffalo in region

under investigation and incurred severed economic losses to poor livestock farmer's community of the area.

Further more, the results obtained in present investigation regarding the Sero-surveillance, antibody titer and geometric mean titer in cattle and buffalo against Hemorrhagic Septicemia (HS) could be used as baseline data for the preparation of vaccine in order to control and prevention of the disease.

REFERENCES

1. Molina, H.A., E.De. La. Pena, G. Camer and M.A. Padilla, 1994. An outbreak of haemorrhagic septicaemia among carabaos in Northern Samar. *Philippine J. Vet. Med.*, 31: 27-31.
2. Dutta, J., B.S. Rathore, S.G. Mulick, R. Singh and G.C. Sharma, 1990. Epidemiological studies on occurrence of haemorrhagic septicaemia in Inida. *Ind. Vet. J.*, 67: 893-899.
3. Zaka, S., 1998. Retrospective epidemiological study and clinico-serological observations on H.S. in cattle and buffaloes in district Dera Ghazi Khan during 1996. M.Sc. thesis, C.V.S., Lahore.
4. Zyambo, G.C.N., R.C. de-Rooij, P.D.S. Moorhouse and H.F. Schels, 1986. The Emergence of Haemorrhagic Septicaemia as a Disease of Economic Importance in Zambia. *Proc. 4th Intl. Symp. Vet. Epidemiol. Eco.*, held on at Singapore, pp: 395-397.
5. Bain, R.V.S., M.C.L. de-Alwis, G.R. Carter and B.K. Gupta, 1982. Haemorrhagic septicaemia. FAO Animal Production and Health Paper rome, Italy.
6. Saini, S.S., D.R. Sharma, B.S. Gill, M.S. Kwarts, J. Singh, J.K. Sharma, S.S. Dhillon and R. Ramneek, 1991. Re-emergence of haemorrhagic septicaemia in Punjab. *Ind. J. Anim. Sci.*, 61: 1178-1180.
7. De Alwis, M.C.L. and A.A. Vipulasiri, 1980. A study of haemorrhagic septicaemia in a wild elephant in Srilanka. *Ceylon Vet. J.*, 13: 17-19.
8. Lane, E.P., N.D. Kock, F.W.G. Hill and K. Mohan, 1992. An outbreak of haemorrhagic septicaemia (septicaemia pasteurellosis) in cattle in Zimbabwe. *Tropical Anim. Health and Prod.*, 24: 97-102.
9. F.A.O., 1989. Fao-who-oje Animal Health Yearbook. Rome, Italy.
10. FAO, 1991. Proceedings of the FAO/APHCA workshop on haemorrhagic septicaemia, Feb. Kandy, Srilanka.
11. Shahid, A.S., 1999. Magnitude of health problems of the livestock population, serological findings and cost/benefit ratio at a farmer level. M.Sc. Thesis, C.V.S., Univ. of Agric., Faisalabad.
12. Sheikh, M.A., M. Anzam and A.R. Shakoore, 1996. Observations on haemorrhagic septicaemia in Pakistan livestock. *J. Vet. Med.*, 43: 293-304.

13. Ibrahim, M., 1993. Epidemiological investigation into economically important livestock diseases, district mardan. A M.Sc. Thesis, CVS, Lahore, University of Agriculture, Faisalabad.
14. De-Alwis, M.C.L. and M.C.L. De-Alwis, 1981. Mortality among cattle and buffaloes in Sri Lanka due to haemorrhagic septicaemia. *Trop. Anim. Health and Prod.*, 13: 195-202.