

Incorporation of DHA and EPA Fatty Acids into Broiler Meat Lipids

R. Salamat Doust Nobar, K. Nazeradl, A. Gorbani, H. Aghdamshahriar and J. Gheyasi
Department of Animal Science, Islamic Azad University, Shabestar Branch, Shabestar, Iran

Abstract: One of the most famous omega 3 fatty acids source is canola oil. The aim of this study was to evaluate the effects on omega 3 fatty acids include of Docosahexaenoic (DHA) and Eicosapentaenoic (EPA) content when Canola Oil (CO) was included in broiler chicks rations. Ninety one day old male broiler chicks (Ross- 308) were randomly distributed into 3 treatment: Control (0% CO), 2% CO and 4% CO for 5 week. These diets were isonitrogenous and isoenergetic were given to broiler chickens throughout a 42 days growth period. Eicosapentaenoic and docosahexaenoic acids and other fatty acids were analyzed by gas chromatography. This trial was conducted in completely randomized design. Birds were slaughtered at 56 days of age. Breast and thigh meat samples were separated and frozen at -20°C until to determine as fatty acid profile. Data was analyzed with one way ANOVA and means compared with Duncan test. Results show that using CO with high level of omega 3 fatty acids could influence fatty acid profile and improved meat quality. With increasing dietary canola oil level in diet (from 0-4 g kg⁻¹ diet) omega 3 contents that were significantly (p<0.05) increased in thigh and breast meat. N-3 fatty acids, DHA and EPA content were significant among treatment (p<0.05).

Key words: Broiler, meat, omega-3, DHA, EPA

INTRODUCTION

There are two reasons for the increasing level of polyunsaturation in chicken meat. First, human nutritionists recommend reducing the intake of Saturated Fatty Acids (SFA) because of its relationship with the development of cardiovascular diseases (Krauss *et al.*, 2001). Secondly, the use of animal fats has been reduced approximately in world, in favor of vegetable oils that are more polyunsaturated. Many authors have studied how the inclusion of different fat sources in the broiler's diet affect the proportion of Fatty Acids (FA), mainly Polyunsaturated Fatty Acids (PUFA), in meat (Salamatdoustnobar *et al.*, 2007; Scaife *et al.*, 1994; Hrdinka *et al.*, 1996; Lo'pez-Ferrer *et al.*, 1999a, b) and the amount of fat deposited by the birds (Sanz *et al.*, 1999, 2000; Crespo and Esteve-García, 2001, 2002a, b). However, there are few reports on the effect of increasing levels of dietary PUFA on the amount and type of FA deposited in chicken tissues, especially in the edible portions. An increase in the degree of polyunsaturation of meat may enhance the development of organoleptic problems (Ajuyah *et al.*, 1993; Gonza'lez-Esquerra and Leeson, 2000; Bou *et al.*, 2001) and lead to an increased susceptibility to lipid oxidation (Klaus *et al.*, 1995; Cortinas *et al.*, 2001; Grau *et al.*, 2001a, b). Most of the reports do not express the FA composition of chicken

meat as amount but rather as profile (percentage of total FA). The objective of this trial was to determine the rate of incorporation of dietary fatty acids from feeding a PUFA-rich diet to broiler meat lipids.

MATERIALS AND METHODS

Animals and diets: Experiment was conducted of the Ross 208 strain were obtained from a commercial hatchery (90 one day old male broiler chicks) and were placed in 9 floor pens of 2×2 m with 10 birds per pen. All chicks were fed a starter diet from 0-21 d and were ad libitum access to water and feed. The experimental design consisted in a completely randomized design with 3 treatments [T1 Control (Soybean + corn), T2 (2% CO) and T3 (4% CO)] with three replication. The treatments diets of were isonitrogenous and isoenergetic. Diets were formulated by adding 0, 2 and 4% canola oil to basal diet (corn and soybean meal) that met the requirements recommended by the National Research Council (1994). The control diet, which was not enriched with canola oil and was administered throughout the 21 days of experimental period (starter). The levels of canola oil were replaced with corn in diets during 2 different periods (grower and finisher). Ingredient composition and nutrient analysis for each treatment is described in (Table 1-3). At the age of 8 weeks, all the birds were weighed before being

Corresponding Author: R. Salamat Doust Nobar, Department of Animal Science, Islamic Azad University, Shabestar Branch, Shabestar, Iran

Table 1: Percentage composition of experimental diet in starter period

Ingredients	(%)
Corn	53.50
Soybean	34.00
Canola oil	0.50
Starch	8.00
Wheat bran	0.00
DL-Methionine	0.54
Lysine	0.00
DCP	1.38
Oyster	1.33
Vitamin 1	0.25
Mineral 2	0.25
Salt	0.25
Coccidiostat	0.00
Sand	0.00
	100.00
Calculated nutrient content	
ME kcal kg ⁻¹	2920.00
Crude protein (%)	21.00
Calcium (%)	0.94
Available P (%)	0.43
ME/CP	139.70
Ca/P	2.10

¹: Vitamin content of diets provided per kilogram of diet: vitamin A, D, E and K, ²: Composition of mineral premix provided as follows kg⁻¹ of premix: Mn, 120,000mg; Zn, 80,000 mg; Fe, 90,000 mg; Cu, 15,000 mg; I, 1,600 mg; Se, 500 mg; Co, 600 mg

Table 2: Percentage composition of experimental diets in growth period

Ingredients	Experimental diets		
	T1 ³	T2	T3
Corn	64	60	55
Soybean	27.4	28	27.1
Canola oil	0	2	4
Starch	3.74	2.06	1.22
Wheat bran	1	2	5.5
DL-Methionine	0	0	0
Lysine	0	0	0
DCP	1.13	1.14	1.16
Oyster	1.5	1.48	1.46
Vitamin ¹	0.25	0.25	0.25
Mineral ²	0.25	0.25	0.25
Salt	0.25	0.25	0.25
Coccidiostat	0.15	0.15	0.15
Sand	0.33	2.42	3.66
	100	100	100
Calculated nutrient content			
ME kcal kg ⁻¹	2920	2920	2920
Crude protein (%)	18.2	18.2	18.2
Calcium (%)	0.9	0.9	0.9
Available P (%)	0.35	0.35	0.35
ME/CP	160.1	160.8	160.7
Ca/P	2.5	2.5	2.5

¹: Vitamin content of diets provided per kilogram of diet: vitamin A, D, E and K, ²: Composition of mineral premix provided as follows kg⁻¹ of premix: Mn, 120,000mg; Zn, 80,000 mg; Fe, 90,000 mg; Cu, 15,000 mg; I, 1,600 mg; Se, 500 mg; Co, 600 mg. ³ T1 = 0%

³: T1= 0% canola oil (CO); T2= 2%CO; T3=4% CO

slaughtered and then eviscerated. Weights air-chilled carcasses after cutting off their heads and feet and after removing abdominal fat (considered as the fat extending within the ischium, surrounding the cloaca and adjacent to the abdominal muscle) to obtain ready-to-cook carcasses were recorded.

Table 3: Percentage composition of experimental diets in finisher period

Ingredients	Experimental diets		
	T1 ³	T2	T3
Corn	66.5	57.5	56
Soybean	24.1	25.85	24
Canola oil	0	2	4
Starch	3.81	4.34	1.94
Wheat bran	0	5	6
DL-Methionine	0.44	0.45	0.45
Lysine	0.043	0.015	0.08
DCP	0.89	0.92	0.89
Oyster	1.38	1.36	1.31
Vitamin ¹	0.25	0.25	0.25
Mineral ²	0.25	0.25	0.25
Salt	0.25	0.25	0.25
Coccidiostat	0.15	0.15	0.15
Sand	1.937	1.665	4.43
	100	100	100
Calculated nutrient content			
ME kcal kg ⁻¹	2920	2920	2920
Crude protein (%)	16.5	16.4	16.5
Calcium (%)	0.79	0.79	0.77
Available P (%)	0.3	0.3	0.3
ME/CP	176.8	177.4	176.6
Ca/P	2.6	2.6	2.6

¹: Vitamin content of diets provided per kilogram of diet: vitamin A, D, E and K, ²: Composition of mineral premix provided as follows kg⁻¹ of premix: Mn, 120,000mg; Zn, 80,000 mg; Fe, 90,000 mg; Cu, 15,000 mg; I, 1,600 mg; Se, 500 mg; Co, 600 mg. ³ T1 = 0%

³: T1= 0% canola oil (CO); T2= 2%CO; T3 = 4% CO

In order to reduce variation in the cutting procedure, all dissections were carried out by one operator. After weighing the eviscerated carcass, it was apportioned into commercial cuts as back, 2 leg-thigh, 2 wings and breast (Hudspeth *et al.*, 1973; Orr *et al.*, 1984). Breast was obtained after removing wings by cutting through the shoulder joint at the proximal end of humerus and by cutting through the ribs, thereby separating the breast from the back (excluding skin). The resulting cut pieces (breast meat, wings and thighs with drumsticks) were then weighed. After quartering, breasts and thighs were separated and frozen at -20°C until to determine as fatty acids profile. The lipid composition was determined by gas chromatography (Model 6890N American Technologies Agilent). The composition of meat samples fatty acid of supplemented lipids s (Table 4-7) data were statistically analyzed using one-way ANOVA and means with significant F ratio were compared by Duncan multiple range test.

Gas chromatography of fatty acids methyl esters

Sample preparation

Fatty acids: Total lipid was extracted from breast and thigh according to the method of Folch *et al.* (1957). Approximately 0.5 g of meat weighed into a test tube with 20 mL of (chloroform: methanol = 2:1, vol/vol) and homogenized with a polytron for 5-10 sec at high speed. The BHA dissolved in 98% ethanol added prior to

Table 4: Least square means for fatty acid profiles in broilers breast meat fed canola oil

	Treatments				
	T1	T2	T3	SEM	p>F
C14:0	0.59 ^a	0.51 ^a	0.54 ^a	0.016883	0.1101
C14:1n5	0.11 ^b	0.10 ^b	0.54 ^a	0.009156	<0.0001
C16:0	28.50 ^a	27.01 ^a	22.71 ^b	0.765465	0.0262
C16:1n7	6.47 ^a	6.60 ^a	5.26 ^b	0.176499	0.0218
C18:0	6.60 ^a	6.21 ^a	6.34 ^a	0.18724	0.4380
C18:1n9	33.65 ^a	30.00 ^a	30.81 ^a	0.920539	0.1299
C18:1n7	2.40 ^b	2.93 ^a	2.73 ^{ab}	0.078816	0.0379
C18:2n6cis	15.35 ^a	13.53 ^{ab}	12.33 ^b	0.404577	0.0295
C18:3n3	0.72 ^b	0.75 ^b	0.87 ^a	0.02186	0.0295
C20:0	0.75 ^a	0.24 ^b	0.23 ^b	0.013268	0.0002
C20:5n3	0.37 ^c	1.18 ^b	2.03 ^a	0.040638	0.0002
C20:1n9	0.17 ^c	0.23 ^b	0.31 ^a	0.007092	0.002
C22:6n3	0.61 ^b	0.62 ^b	0.75 ^a	0.01854	0.0228
C22:0	0.93 ^b	1.96 ^a	1.93 ^a	0.050233	0.0011

Table 5: Least square means for fatty acid profiles in broilers thigh meat fed canola oil

	Treatments				
	T1	T2	T3	SEM	p>F
C14:0	0.60 ^a	0.14 ^c	0.17 ^b	0.0122	0.0002
C14:1n5	0.195 ^c	0.87 ^a	0.47 ^b	0.0171	0.0002
C16:0	26.21 ^a	22.37 ^b	21.88 ^b	0.6878	0.0370
C16:1n7	6.20 ^b	7.83 ^a	6.17 ^b	0.1973	0.0149
C18:0	8.280 ^b	8.96 ^{ab}	10.07 ^a	0.2667	0.0393
C18:1n9	35.32 ^a	37.25 ^a	35.76 ^a	1.0534	0.4686
C18:1n7	2.52 ^a	2.48 ^a	2.27 ^a	0.0703	0.1453
C18:2n6cis	13.14 ^a	11.53 ^b	12.0 ^{ab}	0.3565	0.1002
C18:3n3	0.52 ^b	0.66 ^a	0.74 ^a	0.0185	0.0085
C20:0	0.81 ^a	0.54 ^c	0.63 ^b	0.0185	0.0041
C20:5n3	0.34 ^c	1.43 ^b	2.36 ^a	0.0461	0.0002
C20:1n9	0.12 ^b	0.21 ^a	0.14 ^b	0.0041	0.0013
C22:6n3	0.25 ^b	0.50 ^a	0.57 ^a	0.0126	0.0016
C22:0	0.76 ^b	1.94 ^a	1.96 ^a	0.04656	0.0005

Table 6: Least square means for different traits in broilers breast meat fed canola oil

	Treatments				
	T1	T2	T3	SEM	p>F
Satur f.a	37.37 ^a	35.94 ^{ab}	31.76 ^b	0.96900	0.0534
MUFA	42.80 ^a	39.87 ^a	39.65 ^a	0.80700	0.1166
PUFA	16.33 ^a	15.33 ^a	15.11 ^a	0.39000	0.2061
Total n-6	15.35 ^a	13.53 ^b	12.33 ^b	0.40300	0.0292
Total n-3	1.70 ^c	2.55 ^b	3.66 ^a	0.04300	0.0002
EPA	0.37 ^c	1.18 ^b	2.03 ^a	0.040638	0.0002
DHA	0.61 ^b	0.62 ^b	0.75 ^a	0.01854	0.0228

Table 7: Least square means for different traits in broilers thigh meat fed canola oil

	Treatments				
	T1	T2	T3	SEM	p>F
Satur f.a	36.65 ^a	33.94 ^a	34.73 ^a	0.970	0.2709
MUFA	44.28 ^b	48.62 ^a	44.79 ^{ab}	0.910	0.0761
PUFA	13.74 ^a	13.45 ^a	14.81 ^a	0.330	0.1163
Total n-6	13.14 ^a	11.52 ^b	12.00 ^{ab}	0.360	0.1008
Total n-3	1.12 ^c	2.58 ^b	3.56 ^a	0.053	0.0001
EPA	0.34 ^c	1.43 ^b	2.36 ^a	0.0461	0.0002
DHA	0.25 ^b	0.50 ^a	0.57 ^a	0.0126	0.0016

homogenization. The homogenate filtered through a Whatman filter paper into a 100 mL graduated cylinder and 5 mL of 0.88% sodium chloride solution added, stopper and mixed. After phase separation, the volume of lipid layer recorded and the top layer completely siphoned off. The total lipids converted to Fatty Acid Methyl Esters (FAME) using a mixture of boron-trifluoride, hexane and methanol (35:20:45, vol/vol/vol). The FAME separated and quantified by an automated gas chromatography equipped with auto sampler and flame ionization detectors, using a 30 m 0.25 mm inside diameter fused silica capillary column, as described. A (Model 6890N American Technologies Agilent) (USA) Gas chromatography used to integrate peak areas. The calibration and identification of fatty acid peak carried out by comparison with retention times of known authentic standards. The fatty acid results from gas chromatography with Chem Station software analyzed and expressed as weight percentages.

Statistical analyses: Data were analyzed in a complete randomized design using the GLM procedure of SAS version 12 (SAS Inst. Inc., Cary, NC).

$$y_{ij} = \mu + a_i + \epsilon_{ij}$$

Where:

- y_{ij} = All dependent variable
- μ = Overall mean
- a_i = The fixed effect of oil levels (i = 1, 2, 3)
- ϵ_{ij} = The random effect of residual

Duncan multiple range test used to compare means.

RESULTS

We used various rations with different levels of canola oil to study of it effects on the meat fatty acid profiles. Results of breast meat samples quality parameters (Table 4-7). With increasing levels of dietary polyunsaturation caused the higher accumulation of n-3 (C18:2n6cis, C18:3n3, C20:5n3 and C22:6n3) in thigh and breast meat. The n-3 fatty acids content in thigh and breast meat was significantly affected by canola oil and treatment with 4% CO (T3) was higher level of PUFA fatty acids. According to results with increased canola oil level in all experimental diets, content of omega 6 fatty acids have decreased. T3 and T2 treatment was the lowest omega 6 fatty acids in breast and thigh meat respectively. Result show that the omega 36 fatty acids contents in breast meat was higher than thigh meat. The content of omega 3 fatty acid in T2, T3 (C18:3n3, C20: 5n3 and C22:6n3) for breast and thigh meat is higher than T1

(control) treatment and was significant ($p < 0.0001$). EPA and DHA fatty acids content of breast and thigh meat samples significantly affected by dietary polyunsaturation level and were significant ($p < 0.002$) (Table 4). Comparisons show that T3 treatment (4% CO) with 2.03 and 0.75% for breast meat and 2.36 and 0.57% of thigh total fatty acids is best treatment.

DISCUSSION

Depending on dietary polyunsaturation level, EPA and DHA proportions changed for breast and thigh meat samples from 0.37-2.03% and 0.61-0.75% in breast and 0.34-2.36% and 0.25-57% in thigh respectively. Thus, increasing the level of dietary polyunsaturation caused an increase in the accumulation of PUFA in thigh and breast meat lipids, which was according to Sanz *et al.* (1999), results. Furthermore, both tissues accumulated a similar proportion of total omega 3 fatty acids (C18:3n3, C20:5n3, C22:6n3), respectively 1.70, 1.12 and 2.55, 2.58 and 3.66, 3.56 for breast and thigh meat samples. There are contradictory reports on fatty acids content in breast. Some authors showed that the dietary polyunsaturation level of fat does not influence intramuscular lipid content of breast (Scaife *et al.*, 1994; Crespo and Esteve-García, 2001), but Kirchgessner *et al.* (1993) and Ajuyah *et al.* (1991) found a higher fat content in breast muscle with increasing levels of PUFA in the diet that according with this research finding. However, other authors found lower lipid content of breast of chickens fed diets enriched with polyunsaturated oils (Sanz *et al.*, 1999). Such discrepant findings in intramuscular fat content of breast muscles may be attributed to several factors, such as the analytical procedure used to extract fat from samples. Recent studies showed that fat content of tissues in more polyunsaturated treatments was underestimated when lipid contents were analyzed using AOAC (1995) methodology, suggesting total FA content as an estimator of crude fat in highly polyunsaturated samples (Villaverde *et al.*, 2003b). In general, modification of FA composition of intramuscular fat seems to be more limited (Pan and Storlien, 1993; López-Bote *et al.*, 1997). It may be due to the fact that FA in intramuscular fat are used mainly as components of cellular membranes and the cell has to maintain its physical characteristics to ensure fluidity and permeability of different compounds. As expected, when the dietary polyunsaturation level increased, PUFA content in the tissues also increased (Table 4-7). EPA and DHA contents in breast and thigh samples with increased canola oil from, 0-4% were affected and those differences were 1.66 and 0.14 for

breast and 2.02 and 0.32 is for thigh. In this research T3 with 4% canola oil is best treatment and results show that canola oil could influence meat quality and DHA and EPA contents in broiler meat. (Hulan *et al.*, 1988; López-Ferrer *et al.*, 1999a; González Esquerria and Leeson, 2000; Crespo and Esteve-García, 2001).

ACKNOWLEDGMENT

Financial support for this study (Islamic Azad University, Shabestar Branch) was provided. The authors are also grateful to them valuable support and to oorumieh jahad university for their skilled technical assistance throughout the experimental analyses.

REFERENCES

- Ajuyah, A.O., K.H. Lee, R.T. Hardin and J.S. Sim, 1991. Changes in the yield and in the fatty acid composition of whole carcass and selected meat portions of broiler chickens fed full-fat oil seeds. *Poult. Sci.*, 70: 2304-2314.
- Ajuyah, A.O., R.T. Hardin and J.S. Sim, 1993. Dietary antioxidant and storage affect chemical characteristics of ω -3 fatty acid enriched broiler chicken meats. *J. Food Sci.*, 58: 43-46.
- Bou, R., F. Guardiola, A. Grau, S. Grimpa, A. Manich, A. Barroeta and R. Codony, 2001. Influence of dietary fat source, α -tocopherol and ascorbic acid supplementation on sensory quality of dark chicken meat. *Poult. Sci.*, 80: 1-8.
- Cortinas, L., J. Galobart, A.C. Barroeta, M.S. Castillo and S.K. Jensen, 2001. Influencia del nivel de insaturación dietética sobre el depósito y efecto antioxidante del alfa-tocoferol en muslo de pollo (crudo, cocido y cocido-refrigerado). In: Proceedings of the XXXVIII Symposium Científico de Avicultura. Sección Española de la WPSA. Córdoba, Spain, pp: 141-148.
- Crespo, N. and E. Esteve-García, 2001. Dietary fatty acid profile modifies abdominal fat deposition in broiler chickens. *Poult. Sci.*, 80: 71-78.
- Crespo, N. and E. Esteve-García, 2002a. Dietary polyunsaturated fatty acids decrease fat deposition in separable fat depots but not in the remainder carcass. *Poult. Sci.*, 81: 512-518.
- Crespo, N. and E. Esteve-García, 2002b. Nutrient and fatty acid deposition in broilers fed different dietary fatty acid profiles. *Poult. Sci.*, 81: 1533-1542.
- Fleisch und fett bei stufenweisem austausch von gesättigten durch ungesättigte fettsäuren. *Arch. eflu" Gelk*, 57: 265-274.

- Folch, J., M. Lees and G.H. Sloane-Stanley, 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.*, 226: 497-507.
- Gonza'lez-Esquerria, R. and S. Leeson, 2000. Effects of menhaden oil and flaxseed in broiler diets on sensory quality and lipid composition of poultry meat. *Br. Poult. Sci.*, 41: 481-488.
- Grau, A., F. Guardiola, S. Grimpa, A.C. Barroeta and R. Codony, 2001a. Oxidative stability of dark chicken meat through frozen storage: Influence of dietary fat and α -tocopherol and ascorbic acid supplementation. *Poult. Sci.*, 80: 1630-1642.
- Grau, A., R. Codony, S. Grimpa, M.D. Baucells and F. Guardiola, 2001b. Cholesterol oxidation in frozen dark chicken meat: Influence of dietary fat source and α -tocopherol and ascorbic acid supplementation. *Meat Sci.*, 57: 197-208.
- Hrdinka, C., W. Zollitsch, W. Knaus and F. Lettner, 1996. Effects of dietary fatty acid pattern on melting point and composition of adipose tissues and intramuscular fat of broiler carcasses. *Poult. Sci.*, 75: 208-215.
- Hudspeth, J.P., C.E. Lyon, B.G. Lyon and A.J. Mercuri, 1973. Weight of broiler parts as related to carcass weights and type of cut. *J. Poult. Sci.*, 38: 145-150.
- Hulan, H.W., R.G. Ackman, W.M.N. Ratnayake and F.G. Proudfoot, 1988. Omega-3 fatty acid levels and performance of broiler chickens fed redfish meal or redfish oil. *Can. J. Anim. Sci.*, 68: 533-547.
- Kirchgeßner, M., M. Risitic, M. Kreuzer and F.X. Roth, 1993. Einsatz von fetten mit hohen anteilen an freien fettsäuren in der broilermast. 2. Wachstum sowie qualität von schlachtkörper.
- Klaus, A.M., H. Fuhrmann and H.P. Sallmann, 1995. Peroxidative and antioxidative metabolism of the broiler chicken as influenced by dietary linoleic acid and vitamin E. *Arch. Geflügelk.*, 59: 135-144.
- Krauss, R.M., R.H. Eckel, B. Howard, L. J. Appel and S.R. Daniels, 2001. Revision 2000: A statement for healthcare professionals from the Nutrition Committee of the American Heart Association. *J. Nutr.*, 131: 132-146.
- Lo'pez-Bote, C.J., A.I. Rey, M. Sanz, J.I. Gray and D.J. Buckley, 1997. Dietary vegetable oils and α -tocopherol reduce lipid oxidation in rabbit muscle. *J. Nutr.*, 127: 1176-1182.
- Lo'pez-Ferrer, S., M.D. Baucells, A.C. Barroeta and M.A. Grashorn, 1999a. N-3 enrichment of chickenmeat using fish oil: Alternative substitution with rapeseed and linseed oils. *Poult. Sci.*, 78: 356-365.
- Lo'pez-Ferrer, S., M.D. Baucells, A.C. Barroeta and M.A. Grashorn, 1999b. Influence of vegetable oil sources on quality parameters of broiler meat. *Arch. Geflügelk.*, 63: 29-35.
- Orr, H.L., C. Hunt and C.J. Randall, 1984. Yield of carcass parts, meat, skin and bone of 8 strains of broilers. *Poult. Sci.*, 63: 2197-2200.
- Pan, D.A. and L.H. Storlien, 1993. Dietary lipid profile is a determinant of tissue phospholipid fatty acid composition and rate of weight gain in rats. *J. Nutr.*, 123: 512-519.
- Salamatdoustnobar, R., K. Nazeradl, H. Aghdamshahriyar, A. Ghorbani and P. Fouladi, 2007. The ratio of ω 6: ω 3 fatty acids in broiler meat fed with canola oil and choline chloride supplement. *J. Anim. Vet. Adv.*, 6: 893-898.
- Sanz, M., A. Flores, P. Perez de Ayala and C.J. Lopez-Bote, 1999. Higher lipid accumulation in broilers fed on saturated fats than in those fed on unsaturated fats. *Br. Poult. Sci.*, 40: 95-101.
- Scaife, J.R., J. Moyo, H. Galbraith, W. Michie and V. Campbell, 1994. Effect of different dietary supplemental fats and oils on the tissue fatty acid composition and growth of female broilers. *Br. Poult. Sci.*, 35: 107-118.
- Villaverde, C., M.D. Baucells, L. Cortinas, J. Galobart and A.C. Barroeta, 2003a. Effects of the dietary fat unsaturation level on body fattening in female broiler chickens. In: *Proc. Poult. Sci. Assoc. Ann. Meeting*, Madison, WI, pp: 66.