

Characterisation Phenotypique of Lactic Bacteria Isoles from Believed Milk of Goat of Two Caprine Populations Local Arabia and Kabyle

¹N. Laouabdia Sellami, ²A. Badis, ³D. Guetarni, ⁴R. Ouzrout and ⁵M. Kihal

¹Department of Biology, Faculty of Science, University of Badji Mokhtar,
23000-Annaba, Algeria

²Laboratory of Biochemical Microbiology, Faculty of Science, University of Blida,
9000-Blida, Algeria

³University of Blida 9000-Blida, Algeria

⁴Veterinary Center, University of El Tarf 36000-El Tarf, Algeria

⁵Laboratory of Microbiology, Department of Biology, Faculty of Science,
University Es-senia Oran, Algeria

Abstract: The qualitative distribution of lactic acid bacteria was studied in raw goat's milk of too Algeria's breeds (Kabyle and Arabia). This distribution showed that genus lactobacillus is definitely predominate in kabyle breed (61.48%), follow up, by Lactococcus (22.9%), streptococcus (8.88%), Leuconostoc (5.92%) and by far, the genus Pediococcus (0.74%), while the Leuconostoc and Lactococcus are similarly represented in Arabia (32.64 and 31.02% respectively), follow-up by Lactobacillus (15.27%), streptococcus (14.82%) and pediococcus with fable representation (6.25%). The identification of 240 isolates among morphological, physiological and biochemical criteria have been reattached them with 28 species with the most dominant are Lactobacillus brevis (20 isolates), *Lactobacillus casei* subsp *casei* (20 isolates), *Lactobacillus plantarum* (16 isolates), *Lactococcus lactis* subsp *lactis* (14 isolates) and *pediococcus.acudlactici* (12 isolates).

Key words: Characterisation phenotypique, lactic bacteria, milk of goat, caprine population

INTRODUCTION

The lactic bacteria form a very mixed group composes of hulls and bacilli, whose principal characteristic is the lactic production of acid starting from the fermentation of sugars. Nonpathogenic, these bacteria with positive colouring of gram (Gram-plus) one T an optional anaerobic metabolism and do not produce a catalase. The bacteria colonize varied natural environments such that the surface of the plants and the mucous membranes of the mammals (intestine, stops, vagina and surface of the skin). Their use appeared, since millenia, in the manufacture of food like cheeses, the pork-butcheries, drinks fermented, the leavened bread, sauces, the brines, vegetables fermented, ensilages etc...

They allow, of share their metabolism, to increase the shelf life of origin of the food products and a savour and a different texture confer to them. The insulation of the lactic bacteria starting from goat's milk corresponds on an interesting subject bus little information are available in the literature on the micro-organisms, having

a technological importance, isolated starting from ewe and goat's milk. In Algeria, the caprine breeding comes in second position (14%) after the sheep (26%) (Anonymous, 2000). It is concentrated primarily in the zones of mountains of high plateaus and the arid areas. it is characterized by its adaptation to the climatic conditions of the country and contributes to the formation of the income and the cover of the requirements in, milk and meat of a broad section of the population in the majority of the difficult zones (mountains, high plateaus and arid areas). During the last decade, the annual production of milk is approximately 1 billion liters including goat's milk 13% (Anonymous, 200b). The goat's milk transformation into Raib, Lben and Jben (traditional cheese), generally of varied sensory quality, is done by spontaneous fermentation.

Ea shelf life of origin of the food products and confer a savour and a different texture to them. The insulation of the lactic bacteria starting from goat's milk corresponds on an interesting subject bus little information are available in the literature on the micro-organisms, having

a technological importance, isolated starting from ewe and goat's milk. In Algeria, the caprine breeding comes in second position (14%) after the sheep (26%) (Anonymous, 2000). It is concentrated primarily in the zones of mountains of high plateaus and the arid areas. It is characterized by its adaptation to the climatic conditions of the country and contributes to the formation of the income and the cover of the requirements in, milk and meat of a broad section of the population in the majority of the difficult zones (mountains, high plateaus and arid areas). During the last decade, the annual production of milk is approximately 1 billion liters including goat's milk 13% (Anonymous, 2000b). The goat's milk transformation into Raib, Lben and Jben (traditional cheese), generally of varied sensory quality, is done by spontaneous ferment.

MATERIALS AND METHODS

Site of study and sampling: The taking away of milk came from goats of the population Arabia de Djelfa and the population Kabyle of Tizi Ouzou (large Kabylie). The 19 milk samples were taken on 2 varieties of goats distributed as follows:

- Ten taking away starting from 10 goats of the same Kabyle. Variety
- Last 9 taking away starting from 9 goats of the same Arabia variety.

After washing with soapy water, rinsing with the water chlorinated then dried with a sterile lingette of worse (without trayon) of the goat, the milk collected in sterile bottles (100 mL flacon⁻¹) dice first jets then sampled during the same period due to fodder availability during two years in Mars (2001 for the Kabyle variety and 2002 for the Arabia variety).

Obtaining the isolates: The selective insulation of the lactic bacteria by culture on several mediums was carried out according to the methods described by the International Federation of Milk (Fédération). After dilution of the samples in a solution of Ringer to the ¼ sterile we proceeded to the insulation of the various micro-organisms mentioned in Table 1. The microbiological analyses are made directly starting from the taking away or after enrichment by incubation with 30 and 45°C, until coagulation.

Conservation of the isolates: The short-term conservation of the pure stocks is carried out on tilted solid medium. After growth at the optimal temperature, the cultures are May held with 4°C and the renewal of the stocks is done

Table 1: Mediums used and conditions of incubation for the insulation of the lactic bacteria

Micro-organisms	Mediums of insulation	T°C	Lasted (h)	Incubation
Streptocoques	M17 (Terzaghi and Sandine, 1975)	42-48	72	Aerobiosis
Lactiques	Eliezer (Elliker <i>et al.</i> , 1956)	30	72	Aerobiosis
Lactocoques	Hypersaccharose (Mayeux <i>et al.</i> , 1962)	25	72-144	Aerobiosis
Leuconostocs	M17	30	72	Aerobiosis
Pedicoques	MRS (De Man <i>et al.</i> , 1960)	30	24-36	Anaérobiose
Mésophiles	MRS	45	24-36	Anaérobiose
Lactobacilles Thermophiles				

°C optimal temperature of growth 4

by road repair every 4 weeks. The long-term conservation of the isolates purified is carried out in a medium containing skimmed milk 70% (enriched by 0.05% of extract from yeast and 0.05% of glucose) and glycerol 30% and stored at a temperature of -20°C (Samelis *et al.*, 1994; Herrero *et al.*, 1996).

Determination of the isolates: The identification of the lactic isolates of bacteria at the stage kind was carried out in two stages. The first consists in testing all the isolates by the colouring of Gram, second is based on the morphological study (macroscopic and microscopic) and the type of fermentation. The isolates representative of the dominant flora, were identified at the stage species on the basis of identifying information reported by several authors as follows:

- For the kinds Lactobacillus (Johnson *et al.*, 1980; Kandler and Weiss, 1986; Bottazi, 1988; Falsen *et al.*, 1999; Characteris *et al.*, 2001; Klein, 2001).
- For Streptococcus and Lactococcus (Schleifer *et al.*, 1985; Samelis *et al.*, 1994).
- For Leuconostoc Pediococcus (Devoyod and Poullain, 1988; Lopez and Mayo, 1997; Bissonnette *et al.*, 2000).

The physiological and biochemical criteria are based on the following tests: The test of Scherman and the thermoresistance with 60.5°C during 30 mn (Samelis *et al.*, 1994).

- The growth on M17 medium and MRS, was followed for the temperatures of 5 and 10°C after incubation of 5-7 days and 37, 40 and 45°C after incubation of 24-48 h. According to the temperatures, the isolates of the kinds were tested as follows: Leuconostoc with 5°C, Streptococcus and Lactococcus with 10°C, Leuconostoc with 37°C, Pediococcus, Streptococcus and Lactococcus with 40°C, Streptococcus, Lactococcus, Lactobacillus, Pediococcus and Leuconostoc with 45°C.

- The production of the Arginine Dihydrolase (ADH) on the medium M16 BPC of Thomas (Thomas *et al.*, 1973) and hydrolizes it esculine according to the method of Millière *et al.* (1989).
- The traditional fermentation of the carbohydrates in galleries of tubes on liquid medium MRS and M17 containing the crimson of bromocrésol (0.04 g L⁻¹) like indicator of pH and added with 1% with carbohydrates. The carbohydrates tested are: lactose, glucose, saccharose, galactose, sorbitol, mannose, mélézitose, raffinose, arabinose, xylose, métibiose, fructose and cellobiose.
- The ability to grow on M17 medium and MRS in the presence of NaCl with various concentrations and various values of pH at summer observed during 2-3 days of incubation. The isolates of the kinds were tested as follows: Streptococcus Lactococcus and Lactobacillus with 2% and NaCl 4% and pH 4.5 and 6.5, Leuconostoc with 3 and 6.5% of NaCl and pH 4.5, 4.8 and 6.5 and Pediococcus with 2, 3, 4 and 6.5% of NaCl and pH 4.2, 4.8, 7.0 and 8.0.
- Only the isolates of the kinds Lactococcus, Leuconostoc and Pediococcus were tested for the production of acétoïne in skimmed milk medium according to the technique described by Schmitt *et al.* (1990).
- The production of dextrane starting from saccharose on solid medium MSE of Mayeux *et al.* (1962) was applied for the isolates of Leuconostoc only and the use of citrate to medium Kempler and Mc Kay (1980) pour the isolates Streptococcus and Lactococcus.

RESULTS AND DISCUSSION

Choice of the isolates: From the 19 milk samples, we obtained 919 isolates (48-49 isolates per sample) from which 567 isolates were retained, all in positive Gram, negative catalase and asporulants. Among the 567 remaining isolates, only 240 isolates representative of the dominant flora were identified in a complete way by the phenotypical methods.

Identifications of the isolates: The 567 isolates were attached to 5 groups and were distributed by order of predominance as follows: lactocoques (165 isolates, 29.1%), leuconostocs (149 isolates, 26.28%), lactobacilles (149 isolates, 26.28%), streptocoques (76 isolates, 13.4%) and pedioques (28 isolates, 4.9%) and paid in the Table 2 results of the 240 isolates of stage species are represented in Table 3, the distribution of the 28 species supposed identified is presented according to the kind.

Table 2: Morphological and physiological characters of the supposed kinds of the insulated lactic bacteria

Macro morphology	Micro morphology	Type of fermentation	T°C	Groups (a no. of isolates)
Round or lenticular white colonies	Cocci, Diplocoques and in chain	Homo fermentaire	42-48	Streptocoques (76)
Round or lenticular white colonies	Cocci, Diplocoques and in chain	Homo fermentaire	30	Lactocoques (165)
Very small, Round Transparent colonies	Cocci, ovals, in chain	Hétéro fermentaire	30	Leuconostocs (149)
Colonies smooth Round, white or gray	Cocci into tetrad	Homo fermentaire	30	Pedioques (28)
Small White Colonies in maroon and convex center	Rolled up or Filamentous long Sticks, insulated or in chains	Homo fermentaire	45	Lactobacilles (88)
Round or Lenticular white colonies	Small sticks in chains	Homo fermentaire+ Hétéro fermentaire	30	Lactobacillesb (61)

T°C: Optimal temperature of insulation; Arginine+

Species of lactococcus: On the whole 49 isolates were identified with the lactococcus. Compared to the test of the ADH and the production of acétoïne, they were subdivided in species and under species.

ADH+ and acétoïne+: Ten isolates of *Lc.lactis* subsp.lactis. biovar diacetylactis (all the isolates produced the acétoïne, had the arginine dihydrolase and hydrolized the esculine).

ADH+ and acétoïne-: Fourteen isolates of *Lc.lactis* subsp. lactis (positive growth with 4% NaCl) and 4 isolates of *Lc.lactis* subsp.hordnae (not of growth with NaCl 4%).

ADH- and acétoïne+: Four isolates of *Lc lactis* subsp cremoris (using very few carbohydrates), 13 isolates of *Lc plantarum*, 2 isolates of *Lc gariae* and 2 isolates of *Lc raffinolactis* (these 3 species different only by fermentation from mélézitose and sorbitol).

The species *Lc lactis* subsp. lactis with the biovar, diacetylactis characterized in our isolates is largely used in the dairy products with the other species *Lc lactis* subsp. cremoris, *Lc lactis* subsp. hordinae, *Lc plantarum*, *Lc raffinolactis*, *Lc garvieae* as reported by Garvie. It is important to announce the 4 isolates of *Lactococcus lactis* subsp. cremoris, obtained in our case, also described in Morocco and in the east of Europe (Salama *et al.*, 1995), because these stocks become rare in the industrialized countries.

Species of streptococcus: The 57 isolate identify with the kind *Streptococcus*, 23 were attached to *Streptococcus*

Table 3: Biochemical and physiological criteria of the supposed species of the lactic bacteria isolated from believed milk of goat

Espèce présumées	ADH	ESC	ACT	CTR	DXT	5°C	10°C	37°C	40°C	45°C	PH				PH7	PH	RES	2%	3%	4%	6.5%
											GA-Z	4.2	4.5	4.8							
<i>Lb.acillus helveticus (45)</i>	-	-										+				+					
<i>Lb.plantarum(30)</i>	-	+										+					+				
<i>Lb.brevis (25)</i>	+	V										+					+				
<i>Lb.casei subsp.casei (24)</i>	-	+								V		+					+				
<i>Lb.bulgaricus(26)</i>	-	-										+					+				
<i>Lb.rhamnosus(20)</i>	-	+										+					+				
<i>Lb.casei subsp. alactosus (14)</i>	-	+										+					+				
<i>Lb.lactis (13)</i>	V	+								+		+					+				
<i>Lb.acidophilus (01)</i>	-	+								+		+					+				
<i>Lb.animalis (01)</i>	+	±								+		+					+				
<i>Lb.amylophilus (01)</i>	-	-										±					+		±		+
<i>Streptococcus therphilus(46)</i>	-	-															+	V			
<i>Lc.diacetylactis (18)</i>	+	+	+	+						+							+	V	+		
<i>Lc.lactis subsp.lactis (25)</i>	+	+	-	-						+							+	V	+		
<i>Lc.plantarum (22)</i>	-	+	+	-						V							+		+		
<i>Lc.lactis subsp. hordnae (11)</i>	+	+	-	±													+				
<i>Lc.lactis subsp. cremoris(10)</i>	-	V	++	-													+				
<i>Lc.garviae (02)</i>	+	+	+	-						+							+		+		
<i>Lc.raffinolactis(02)</i>	V	+	-	v													+		+		
<i>Leuconostoc lactis (11)</i>	-	-	-									+					+		+		
<i>Ln.mesenteroides subsp. dextranicum (04)</i>	-	±	+		+					+	+						+		±		
<i>Ln.amelibiosum(07)</i>	-	-	+		+						+						+		+		
<i>W.paramesenteroides(10)</i>	-	-	+		-	+				±	+						±		±		±
<i>Ln. Mesenteroides subsp.cremoris(07)</i>	-	-	+		-	+					+						+				
<i>Ln.pseudomesenteroides(02)</i>	-	±	+		-	+				+	+						+			+	
<i>Pediococcus damnosus (06)</i>	-	-	V														+				
<i>p.acidilactici (12)</i>	+	+	+									V				+	V			+	V
<i>p.parvulus(03)</i>	+	-	-									+	+			+	-			+	+

A number of isolates really tested +: more than 90% of reaction positive, -: less than 10% and less positive creations. ADH: production of the arginine dihydrolase. ESC: hydrolize esculine ACT: production of acetoïne. CTR: Citrate DXT degradation: production of dextrane LMBO: thermorésistance with 63.5°C during 30 min. GAS: production of gas starting from glucose%: Na Cl concentration

thermophilus. These 23 isolates of *St thermophilus* able to grow to heat-resisting pH 6.5 quickly and slowly pH 4.5 et 4.8, not producing an acetoïne and not hydrolizing the esculine, ferment some differently sucres (43% fermented lactose and galactose, 20.6% have ferments the mélibiose, 18% have ferments raffinose and 14% have ferments the mélézitose).

Species of leuconostoc: The 29 isolate of *leuconostoc* are divided into 6 species: 7 of *Ln.lactis*, 5 of *Weissella paramesenteroides* (old *Ln. Paramesenteroides*) 7 stocks of *Ln. amylibiosum*, 7 of *Ln mesenteroides subsp. cremoris*, 4 de *Ln.mesentéroide subsp. dextranicum*, et 2 *Ln. pseudomesenteroide*. Une heterogeneity in the fermentation of sugars was noticed for isolates of 3 species S of *leuconostoc*:

- With a very high degree for *Ln. mesenteroides subsp dextranicum* (galactose = 32% mannose = 50%, raffinose = 56%, xylose = 75%, mélibiose = 45%, cellobiose = 55%).

- Follow-up of *W paramesenteroides* (galactose=50%, glucose = 50%, raffinose=66%, arabinose = 58%, xylose = 76%).
- With a less degree for *Ln. lactis* (mannose=72%, raffinose=38% and mélibiose=66%).

Species of pediococcus: Three species of *Pediococcus* were isolated. They form colonies smooth round grisâtres or blanchâtres on M17 medium. The microscopic observation in a fresh state between blade and plate reveals a form characteristic of cocci into tetrad. Among the 19 identified stocks 12 belong to l'espèce *P.acidilactici*, 4 with the *P.damnossus* species and 3 with the species *P. Parvulus*.

Species of lactobacillus: One hundred and twenty insulators of *Pediococcus* were classified in 11 species normally they are differentiated by their type of fermentation from the carbohydrates but the profile of the fermentation of its isolates was compared with those of stocks of references in the key of identification of

Bergey (Kandler and Weiss, 1986) and of the considerable differences were noted. By way of example: an isolate, identified as *Lb. animalis* differs from the stock of reference by the cellobiose fermentation, from the fructose and the galactose considered typical of the stock of reference. The isolates lactobacillus were subdivided in three groups:

Group of the thermophilous lactobacillus and homofermentaires strict: The isolates this group ferment hexoses exclusively by producing lactate but do not ferment produce lactate exclusively, but do not ferment pentoses. The associated species are:

- 01 isolate of *Lb. acidophilus* which has a typical fermentation as for L stock of reference.
- 12 isolates of *Lb. delbrueckii* of which 7 de *Lb. delbrueckii* subsp. *bulgaricus* (typical Fermentation) and 5 of *Lb. delbrueckii* subsp. *lactis* which ferments four carbohydrates in a different way (50% of the isolates fermented galactose and cellobiose, 44.4% ont ferments the melibiose and 22.2% have ferments the sorbitol).
- Twenty nine isolates of *Lb. helveticus* which ferment three differently carbohydrates (42% saccharoses, 42.8% the sorbitol and 42.9% fructose). Whereas the remainder of the carbohydrates were fermented either by 100% of the isolates or 0% of the isolates.
- 01 isolate of *Lb. animalis* which does not ferment cellobiose, the fructose and galactose, fermentation considered typical for the pilot stock.

Group of the optional lactobacilles mésophiles and homofermentaires: The isolates D this group ferment hexoses exclusively by producing lactate and do not produce a gas starting from glucose. They can ferment pentoses. Three species belonging to this group were isolated:

- 58 isolates of *Lb. casei* including 20 isolates of *Lb. casei* subsp. *casei* (typical fermentation), 10 of *Lb. casei* subsp. *rhamnosus* (typical fermentation) and of 10 of *Lb. casei* subsp. *alactosus* (typical fermentation).
- 30 isolates of *Lb. plantarum* (53.9% fermented saccharose, 53% raffinose and cellobiose and 46.6 fermented xylose). 01 isolate of *Lb. amylophilus* (typical fermentation).

Group of the strict lactobacillus mésophiles or thermophilous and hétérofermentaires: The isolates of group ferment hexoses out of lactate and/or ethanol

and CO₂. One only isolated species was attached to *Lb. brevis* (20 isolates with 40% fermented the arabinose). This species like *Lb. buchneri*, *Lb. fermentum* and *Lb. parabuchneri* were insulated in milk and dairy products, as pays by Dellaglio *et al.* (1994). The species of lactobacilles hétérofermentaires isolated in the meats and incompetents to grow on acetate medium were introduced into the kind carnobacterium (Collins *et al.*, 1987). The search for production of gas starting from gluconate, noncarried out in more of the test of production of gas starting from glucose would undoubtedly have allowed us to classify the lactobacillus more easily insulated according to their type of fermentation (lactobacilles homofermentaires, lactobacilles hétérofermentaires optional and lactobacilles hétérofermentaires obligatory) as described by Demarigny *et al.* (1996). Generally the morphological analyses, physiological and biochemical showed a diversity of kinds and species isolated starting from milk believed from two population caprine local (Kabyle and Arabia) this composition from lactic bacteria is relative and depends on different the criterion used in each study as brought back by Fitzsimmons *et al.* (1999) and Bissonnette *et al.* (2000).

The identification of the isolates by the traditional methods presents much bus difficulties of the significant differences in the profiles of fermentation of sugars were observed. Consequently, the genotypic identification is essential to confirm phenotypical classification.

Distribution of the lactic bacteria

Distribution of the kinds: The distribution of the kinds in the two caprine populations seems to show that the *Lactobacillus* kind is definitely prevalent in the population Kabyle (61.48%); followed of *Lactococcus* (22.9%), *Streptococcus* (8.88%), *Leuconostoc* (5.92%) and by far *Pediococcus* (0.74%), whereas the two kinds *Leuconostoc* and *Lactococcus* are represented in an almost similar way in the population Arabia (32.64% and 31.02, respectively), like *Lactobacillus* (15.27%), *Streptococcus* (14.82%) and *Pediococcus* with a weak representation (6.25%).

The weak rate of the lactobacilles in the Arabia population, compared to the Kabyle population, seems to be with competitiveness "Lactobacilles/ Streptocoques" in the fermentation of the milk which is in favour of the streptocoques ones. This assumption was already advanced by Harrati (1974) which had noted their total absence in Algerian Lben. On the other hand, the prevalence of the lactobacilles recorded in the Kabyle population falls on enrichment by incubation with 30°C and 45°C until coagulation of the milk samples. The same remarks were announced by Tantaoui-Elaraki *et al.* (1983) and Tzantakis and Litopoulou-Tzanetaki (1992)

Table 4: Distribution of the species of BL isolated from believed milk of goat of two caprine populations

Espèce	%	n	A	K
Lb.delbrueckii subsp.bulgaricus	2.92	7	0	7
Lb.helveticus	12.03	29	8	21
Lb.plantarum	6.67	16	9	7
Lb.casei subsp.rhannosus	4.14	10	1	9
Lb.delbrueckii subsp.lactis	2.08	5	5	0
Lb.brevis	8.33	20	6	14
Lb.casei subsp.alactosus	4.17	10	0	10
Lb.casei subsp. Casei	8.33	20	5	15
Lb.acidophilus	0.42	1	1	0
Lb.animalis	0.42	1	1	0
Lb.amylophilus	0.42	1	1	0
Streptococcus thermophilus	9.58	23	11	22
Lc.lactis biovar diacetyllactis	4.17	10	1	9
Lc.lactis subsp.lactis	5.83	14	6	8
Lc.lactis subsp.hordniae	1.67	4	0	4
Lc.lactis subsp.cremoris	1.67	4	1	3
Lc.plantarum	5.42	13	10	3
Lc.garviae	0.83	2	0	2
Lc.raffinolactis	0.83	2	0	2
Leuconostoc lactis	2.92	7	5	2
Ln.mesenteroides subsp.cremoris	1.67	4	0	4
Weissella paramesenteroïdes	2.08	5	3	2
Ln.pseudomesenteroïdes	0.83	2	2	0
Ln.mesenteroides subsp.dextranicum	1.67	4	4	0
Ln.amylibiosum	2.92	7	7	0
Pediococcus damnosus	1.67	14	3	1
P.acidilactici	5	12	12	0
P.parvulus	1.25	3	3	0
Total	99.77	(100)	(43.75)	(56.25)

A: the Arabia population. K: the population Kabyle.%: percentage of each species. N: total of the isolates in the species. *Pourcentage of the isolates in each population

or the prevalence of the lactobacilles is explained by their growth favorable to low pH. The rate high of the leuconostocs in the Arabia population but low in the Kabyle population is not easy to explain. However, for Sawaya *et al.* (1984), a relation exists between the composition in lactic bacteria, the composition of milk and the food. The pediococques ones were not highlighted in the majority of the samples of milk and the number of isolates was always weak, even in the samples of the Arabia population. Resulted similar were brought back by Saidi *et al.* (2002) in the characterization of 206 isolates of BL starting from believed milk of the caprine population of the arid areas, or they showed a clear predominance of lactocoques (76.16%), *Lc.lactis* subsp.lactis. thus, Bekhouche *et al.* (1998) in the insulation of BL starting from cow's milk in the Algerian east or they showed a homogeneous distribution with a great frequency of *Leuconostoc*, follow-up of *Lactobacillus*, *Pediococcus* and *Streptococcus*. characterization of 206 isolates of BL starting from believed milk of the caprine population of the arid areas, or they showed a clear predominance of lactocoques (76.16%), *Lc.lactis* subsp.lactis. thus, Bekhouche *et al.* (1998) in the insulation of BL starting from cow's milk in

the Algerian east or they showed a homogeneous distribution with a great frequency of *Leuconostoc*, follow-up of *Lactobacillus*, *Pediococcus* and *Streptococcus*.

Distribution of the species: The results of the distribution of the lactic bacteria according to the species are reported in Table 4. The number of species shows a low variability, about 19 and 22 at Arabia and Kabyle respectively. The species, *PC acidolactici* (12 isolates), *Ln.amylibiosum* (7 isolates)), *Ln.mesenteroides* subsp *dextranicum* (4 isolates), *PC parvulus* (3 isolates), *Ln.pseudomesenteroides* (2 isolates), *Lb.acidophilus* (01 isolates), *Lb.amylophilus* (01 isolates) and *Lb.animalis* (01 isolates) are found only in the Arabia population. The species *Lc.garviae* (2 isolates), *Lc.raffinolactis* (2 isolats) ne are found that in the population kabyle. This distribution differs from one population to another as follows:

- In the Arabia population, the distribution of the other species is characterized by predominance from the species *Lc.plantarum* (10 isolates), *PC acidolactici* (12 isolates), *St.thermophilus*. (11 isolates) and *Lb.plantarum* (10 isolates).
- In the population Kabyle, the species *Lb.helveticus* (21 isolates), *Lb.casei* subsp.*casei* (15 isolates), found in the other dominant populations.

REFERNECES

Albenzino, M., M.R. Corbo, S.U. Rehman, P.F. Fox, M. de Angelis, A. Corsetti, A. Sevi and M. Gobetti, 2001. Microbiological and biochemical characteristics of canestrato Pugliese cheese made from raw milk, pasteurized milk or by heating the curd in hot whey it. *J. Food. Microbiol.*, 67: 35-48.

Anonyme, 2000a. Statistiques Agricoles, 1990-1999. Série B, superficies et production. ONS (Office National des Statistiques). Alger Algérie.

Anonyme, 2000b. Note de conjoncture sur les performances zootechniques des élevages bovins laitiers en Algérie 1999-2000. Institut Technique des Élevage (ITELV), Alger, Algérie.

Bekhouche, F., Bachir-Cherif and M. Gheribi, 1998. Isolements et identification de bactéries lactiques à partir de lait cru de vaches. XIème Journées Nationales de Microbiologie. Oran. Algérie. 30 novembre et 10 décembre.

Beresford, T.P., N.A. Fitzsimmons, N.L. Brennan and T.M. Cogan, 2001. Recent advances in cheese microbiology. *Int. Dairy J.*, 11: 259-274.

- Bissonnette, F., S. Labrie, H. Deveau, M. Lamoureux and S. Moineau, 2000. Characterization of mesophilic mixed starter cultures used for the manufacture of aged cheddar cheese. *J. Dairy Sci.*, 83: 620-627.
- Bottazi, V., 1988. An introduction to red shaped lactic acid bacteria. *Biochimie*, 70: 303-315.
- Characteris, W.P., P.M. Kelly, L. Morelli and J.K. Collins, 2001. Quality control *Lactobacillus* strains for use With the API 50CH and API ZYM systems at 37°C, *J. Basic. Microbiol.*, 41: 241-251.
- Collins, M.D., J.A. Farrow, B.A. Philips, S. Ferusu and D. Jones, 1987. Classification of *Lactobacillus divergens*, *Lactobacillus piscicola* and some catalase-negative asporogenous, rod-shaped bacteria from poultry in a new genus *Carnobacterium*. *Int. J. Syst. Bacteriol.*, 37: 310-316.
- De Man, J., M. Rogos and M.E. Sharpe, 1960. A medium for a cultivation of lactobacilli. *J. Applied Bacteriol.*, 23: 130-135.
- Deghuchi, Y. and T. Morishita, 1992. Nutritional requirement multiple auxotrophic lactic acid bacteria: Genetic lesion affecting amino acid biosynthetic pathways in *Lactococcus lactis*, *Enterococcus faecium* and *pediococcus acidilactici*. *Biosci. Biotechnol. Biochem.*, 56: 913-918.
- Dellaglio, F., H. De Roissart, S. Torriani, M. Curd and C. Janssens, 1994. In: *Bactéries lactiques aspects fondamentaux et technologiques*, 1: 25-114.
- Demarigny, Y., E. Beuvier, A. Dasen and G. Duboz, 1996. Influence of raw milk micro flora on the characteristics of Swiss-type cheese. 1. Evolution of micro flora during ripening and characterization of facultatively heterofermentative lactobacilli. *Lait*, 76: 371-387.
- Devoyod, J.J. and F. Poullain, 1988. Les leuco nostocs propriétés: Leur rôle en technologie laitière. *Lait*, 68: 249-280.
- Elliker, P.R., A.W. Anderson and G. Hannesson, 1956. An agar culture medium for lactic streptococci and lactobacilli. *J. Dairy. Sci.*, 39: 1611-1612.
- Falsen, E., C. Pascual and B. Sjoden, 1999. Phenotypic and phylogenic characterisation of a nouvel *Lactobacillus* species from human source: description of *lactobacillus inners*. sp. nov. *Int. J. Syst. Bacteriol.*, pp: 49.
- Fitzsimmons, N.A., T.M. Cogan, S. Condon and T. Beresford, 1999. Phenotypic and Genotypic Characterization of non-strarter lactic Acid Bacteria in Mature Cheddar Cheese. *Applied Environ. Microbiol.*, 65: 3418-3426.
- Frey, L., 1993. Les bactériocines des bactéries lactiques. *Bull. Soc. FR Microbiol.*, 4: 245-251.
- Gavie, E.L., 1984. Taxonomy of identification of bacteria important in cheese and fermented dairy products. In: *advanced in microbiology and biochemistry of cheese fermented Milk*, Elsevier Applied Science Publisher, London N.Y., pp: 35-65.
- Gibbs, P.A., 1987. Novel uses for lactic fermentation in food preservation. *J. Applied Bacteriol.*, 5: 51-58.
- Gomes, A.M., F.X. Malcata and F.A. Klaver, 1998. Growth enhancement of *Bifidobacterium lactis* Bo and *Lactobacillus acidophilus* Ki by milk hydrolysate. *J. Dairy Sci.*, 81: 25-281.
- Harrati, E., 1974. Recherche sur Lben et le Klila algériens, These de doctorat de spécialité. Université de Caen. France.
- Herrero, M., B. Mayo, B. Gonzalez and J.E. Suarez, 1996. Evaluation of technologically important traits in lactic acid bacteria isolated from spontaneous fermentation. *J. Applied Bacteriol.*, 81: 565-570.
- Hitchener, B.J., A.F. Egan and P.J. Rogers, 1982. Characteristics of lactic acid bacteria isolated from Vacuum -Packaged beef. *J. Applied Bacteriol.*, 52: 31-37.
- Huang, D.Q., H. Prévost, M. Kihal and C. Diviès, 1994. Instability of plasmid encoding for β . Galactosidase in *Leuconostoc mesenteroides* subsp *mesenteroides*. *J. Basic. Microbiol.*, 34: 23-30.
- Hutking, R., S.M. Halambeck and H.A. Morris, 1986. Use of gelatos-fermenting *Streptococcus Thermophilus* in the manufacture of Swiss, Mozzarella and short-method Cheddar cheese. *J. Dairy. Sci.*, 69: 1-8.
- Jenness, R., 1982. Composition and characteristics of goat milk: Review 1968-1979. *J. Dairy Sci.*, 63: 605.
- Johnson, J.L., C.F. Phelips, C.S. Cummins, J. London and F. Gasser, 1980. Taxonomy of *lactobacillus acidophilus* group. *Int. J. Syst. Bacteriol.*, 30: 53-68.
- Kandler, O. and N. Weiss, 1986. Regular, non-sporforming Gram-positive rods, In: *Sneath, P.H.A., Mair N., Sharpe M.E. et Holt J.G. (EDS.), Bergey's Manual of Systematic Bacteriology. Vol. 2. Williams and Wilkins, Baltimore*, pp: 1208-1234.
- Kempler, G.M. and L.L. Mc Kay, 1980. Improved medium for detection of citrate-fermenting *Streptococcus lactis* subsp *diacetylactis*. *J. Applied Environ. Microbiol.*, 39: 956-957.
- Klein, G., 2001. International committee of Systematic Bacteriology. Subcommittee on the taxonomy of *Bifidobacterium*, *Lactobacillus* and related organisms. Minutes of the meeting. *Int. J. Syst. Evol. Microbiol.*, 51: 259-261.
- Lopez, S. and B. Mayo, 1997. Identification and characterization of homofermentative mesophilic *Lactobacillus* strain isolated from artisan starter-free cheeses. *Lett. Applied Microbiol.*, 25: 233-238.

- Mayeux, J.V., W.W.E. Sandine and P.R. Elliker, 1962. A selective medium for detecting *Leuconostoc* organisms in mixed strain starter cultures. *J. Dairy Sci.*, 45: 655-656.
- Millière, J.B., A.G. Mathot, P. Schmitt and C. Diviès, 189. Phenotypic characterization of *Leuconostoc* species. *J. Applied Bacteriol.*, 67: 529-542.
- Picque, D., B. Perret, E. Latrille and G. Corrieu, 1992. Caractérisation et classification des bactéries lactiques à partir de la mesure de leur cinétique d'acidification, *Lebensm. Wiss. U. Technol.*, 25: 181-186.
- Remeuf, F., V. Cossin, C. Dervin, J. Lenoir and R. Tomassone, 1991. Relations entre les caractéristiques physico-chimiques des laits et leur aptitude fromagère. *Lait*, 71: 397-421.
- Ross, R.P., S. Morgan and D. Hill, 2002. Preservation and fermentation. Past, present and future. *Int. J. Food Microbiol.*, 79: 3-16.
- Saidi, N., B. Guessas, F. Bensalah, A. Badis, M. Hadadji, D. Henni, H. Prevost and M. Kihal, 2002. Caractérisations des bactéries lactiques isolées du lait cru de chèvre des régions arides. N° 01 juin, pp: 1-10.
- Salama, S.M., T. Musafija-Jeknic, E.W. Sandine and J.S. Giovannoni, 1995. An ecological study of lactic acid bacteria. Isolation of new strains of *Lactococcus lactis* subsp *cremoris*. *J. Dairy. Sci.*, 78: 1004-1007.
- Samelis, J., F. Maurogenakis and J. Metaxopoulos, 1994. Characterization of lactic acid bacteria isolated from naturally fermented Greek dry salami. *Int. J. Food. Microbiol.*, 23: 179-196.
- Sawaya, W.N., W.J. Safi and A.F. Shalhat, 1984. Chemical composition and nutritive value of goats milk. *J. Dairy Sci.*, 67: 1655-1659.
- Schleifer, K.H., J. Kraus, C. Dvorak, R. Killper-Blz, M.D. Collins and W. Fisher, 1985. Transfer of *Streptococcus lactis* and related *Streptococci* to the genus *Lactococcus* gen. Nov. System. *Applied Microbiol.*, 6: 183-195.
- Schmitt, P., C. Diviès and C. Merlot, 1990. Utilization of citrate by *leuconostoc mesenteroides* subsp. *cremoris* in continuous culture. *Biotchnol. Lett.*, 12: 127-130.
- Tantaoui-Elaraki, A., M. Berrada, A. El Marrakchi and A. Berramou, 1983. Etude sur le Lben marocain. *Le lait*, 63: 230-245.
- Terzaghi, B.E. and W.E. Sandine, 1975. Improved medium for lactic streptococci and their bacteriophages. *Applied Environ. Microbiol.*, 29: 807-813.
- Thomas, T.D., 1973. Agar medium for differentiation of *streptococcus cremoris* from the other bacteria. *N.Z.J. Dairy. Sci. Technol.*, 8: 620-627.
- Tzanetakis, N. and E. Litopoulou-Tzanetaki, 1992. Changes in numbers and kinds of lactic acid bacteria in Feta and Teleme, tow Greek cheeses from ewes milk. *J. Dairy. Sci.*, 75: 1655-1659.