

Effect of Different Phases of Estrous Cycle on Kisspeptin Expression in the Arcuate Nucleus of the Ewe

¹Jafarzadeh Shirazi Mohammad Reza, ¹Sepehri Hoori and ²Gheisari Hamid Reza

¹School of Biology, University College of Science, University of Tehran, Tehran, Iran

²Department of Food Hygiene, College of Veterinary Medicine,
University of Shiraz, Shiraz, Iran

Abstract: Kisspeptin (KP) is a product of prepro-kisspeptin. KP is a key regulator of GnRH and LH secretion in the ewe. In this study, Effect of different phases of estrous cycle on kisspeptin expression in the arcuate nucleus of the ewe examined. Because of no difference between phases of cycle, we suggested that perhaps KP neurons involved in mediating of both positive and negative feedback effects of steroids to GnRH neurons.

Key words: Kisspeptin, estrous cycle, arcuate nucleus ewe, GnRH

INTRODUCTION

KP is a key regulator in LH and GnRH secretion (Dungan *et al.*, 2006; Smith and Clarke, 2007; Roa and Tena-Sempere, 2007) and causes GnRH and LH releasing in rodent (Gottsch *et al.*, 2004; Thompson *et al.*, 2004; Navarro *et al.*, 2004, 2005) sheep (Messenger *et al.*, 2005) monkey (Plant *et al.*, 2006) and human (Dhillon *et al.*, 2005). Mutation in GPR54 put off puberty beginning in human (De Roux *et al.*, 2003) and mouse (Seminara *et al.*, 2003; Funes *et al.*, 2003). KP has been involved in feedback functions of ovary steroids. In sheep, KP exerts positive feedback actions in medio-basal hypothalamus which result in preovulatory surge of LH (Caraty *et al.*, 1998; Couse and Korach, 1999). Effects of sexual steroids convey to GnRH neurons by different systems of cells. GnRH neurons do not express estrogen receptor- α (ER- α) or progesterone receptor (PR) (Herbison and Pape, 2001) but they express estrogen receptor- β (ER- β) (Dorling *et al.*, 2003). Regarding steroid feedback effects on GnRH/LH, however, ER- β appears to play a slight role in the regulation of reproduction. Strong evidences show that effects of steroid feedback on GnRH/LH results from cells which express ER- α (Goubillon *et al.*, 1999). Possibly, stimulating effects on GnRH surge are influenced by ER- α (Couse and Korach, 1999). Most of ER- α located in arcuate nucleus (ARC). So, other sensitive neurons to steroid, rather than GnRH, should mediate effects of sexual steroids on GnRH secretion. Because KP producing cells in the ARC are abundant in sheep (Franceschini *et al.*, 2006; Pompolo *et al.*, 2006) and

also these cells include ER- α , perhaps KP cells convey positive feedback of steroids to GnRH neurons.

This research aimed to test expression levels of KP neurons in estrous cycle. We proposed that levels of expression of KP in the follicular phase may be more than other phases.

MATERIALS AND METHODS

Animals: Nine Corriedale ewes with similar age and weight were maintained under natural pasture situation. Research was performed during the breeding season. Estrous cycle controlled by using rams in the group. Estrous cycles were synchronized with Cloprostenol. Three groups of animals (n = 3 in each phase) were used, indicating the luteal phase (day 10 of cycle), the follicular phase and estrous phase. Follicular and luteal phases animals were killed 24 h and 10 days after the injection of Cloprostenol, respectively. Estrous animals were killed 1-2 h after the onset of the Standing heat (about 48 h after Cloprostenol injection). Blood samples from the estrous group of ewes were collected prior to the onset of estrous and just before slaughtering of the animals (1-2 h after the onset of standing heat). For the follicular and luteal phase groups, the blood samples were taken just prior to killing the animals. Both ovaries of each animal were also examined for the presence/absence of corpora lutea. Levels of the plasma hormone in these sheep have been explained previously (Estrada *et al.*, 2006). Progesterone levels in luteal phase's ewes were 3.2 ± 0.5 ng mL⁻¹ (MEAN \pm SEM). In other phases, levels of Progesterone

were lower. Plasma LH levels were $<1.0 \text{ ng mL}^{-1}$ in ewes during the luteal phase. These levels in late-follicular and across estrus phase's ewes were 1.4-6.9 and $51.4 \pm 4.2 \text{ ng mL}^{-1}$, respectively.

Tissue collection: Tissue was collected and processed as previously described (Foradori *et al.*, 2006; Pompolo *et al.*, 2006). Briefly, ewes were killed with an overdose of sodium pentobarbital and their heads were perfused with 2 L of heparinized saline (12.5 U mL^{-1}) followed by 2 L of 4% paraformaldehyde plus 15% picric acid in PB and then 1 L of the same fixative containing 20% sucrose. Brains were removed, tissue containing the hypothalamus dissected out, infiltrated with 30% sucrose and frozen coronal sections ($40 \mu\text{m}$ thick) cut and stored at -20°C in cryoprotectant.

Immunohistochemistry: Rabbit's antisera against kisspeptin were provided by Dr. Caraty. Test of Immunohistochemical characteristics of the rabbit polyclonal antibody against kisspeptin-10, in previous experiment has been explained (24), comparing it with a commercially available (Phoenix Pharmaceuticals, Inc) rabbit polyclonal antibody against kisspeptin-10 (human sequence). Three sections/sheep were selected for analysis to represent the rostral, middle and caudal regions of the ARC in each phase of estrous cycle. Sections were washed in 0.05 M PBS, mounted on slides and dried overnight. Kisspeptin-containing cells were labelled using antigen retrieval and Caraty kisspeptin antibody (1: 2000) was used. After incubation of slides with 1: 500 goat's anti-rabbit sera conjugated to Alexa488, they were washed 3 times with PBS and then counterstained with 0.3% Sudan Black B. After further washing, coverslips were applied with fluorescence mounting medium (Dako, Carpinteria, CA). A positive control (no antibody pre-absorption) and a negative control (antibody pre-absorbed with kisspeptin) were used. Images of immunoreactive cell bodies were captured with the proper excitation for Alexa 488. A single viewer then counted the total number of immunopositive cell bodies.

Statistical analysis: Data were analyzed by general linear model of SPSS software (version 11.5). The mean number of cells within areas was compared by 2-way ANOVA (main effects of cycle stages and anatomical area). Duncan's multiple range used as a post hoc test. And $p < 0.05$ was considered as a significance level.

RESULTS

No effects of phase cycle within area of ARC were observed in cell number ($p > 0.05$), except in luteal phase




Fig. 1: Top panel: Fluorescence images of rostral (A) middle (B) and caudal ARC immunostained for kisspeptin. Arrows indicate examples of kisspeptin neurons. Bottom panel: Mean (\pm SEM) number of kiss cells in rostral, middle and caudal of ARC within estrous cycle

that it's cells from middle area were higher than 2 others area ($p < 0.05$). Number of cells in the estrous phase of rostral area also were higher with respect to follicular and luteal phase in this area (Fig 1).

DISCUSSION

This study showed that there is no significant difference between phases of cycle in areas of ARC. In contrast, in previous work (Estrada *et al.*, 2006), up-regulation of KP in the rostral ARC during estrous phase has been reported. Exact details of this experiment whether which kind of antibody used, unknown. We used caraty antibody with high validity that previously demonstrated uncross reactivity of it with other RF-amides (Goodman *et al.*, 2007). Because no difference between phases of cycle, several lines of evidence inferred that perhaps KP neurons involved in mediating of both positive and negative feedback effects of steroids to GnRH neurons. First of all, KP neurons in the ovine contain PR (Smith *et al.*, 2007). We previously showed that KP neurons overlap with Dynorphin neurons in the ARC (Goodman *et al.*, 2007). Dynorphin neurons contain PR (Foradori *et al.*, 2002) and an antagonist to the Dynorphin receptor stimulates episodic LH secretion in luteal phase ewes (Goodman *et al.*, 2004). Because

progesterone has negative feedback effects on GnRH neurons which have no PR, then our data suggest that KP neurons may relay progesterone negative feedback effects to GnRH neurons. Second, KP neurons abundant in the ARC of sheep (Franceschini *et al.*, 2006; Herbison and Pape, 2001) and these neurons contain ER- α (Franceschini *et al.*, 2006), we can suggest that KP neurons in the ARC transmit the positive feedback effect to the GnRH cells which do not own ER- α .

CONCLUSION

KP neurons may be implicated in conveying of positive and negative feedback effects of ovarian steroids to GnRH neurons in preoptic area that have no PR or ER- α . Attempts are ongoing to apply the knowledge of the functions of KP to therapeutic uses including under fertility and over fertility.

REFERENCES

- Blache, D., C.J. Fabre-Nys and G. Venier, 1991. Ventromedial hypothalamus as a target for oestradiol action on proceptivity, receptivity and luteinizing hormone surge of the ewe. *Brain Res.*, 546: 241-249.
- Caraty, A., C. Fabre-Nys, B. Delaleu, A. Locatelli, G. Bruneau, F.J. Karsch and A. Herbison, 1998. Evidence that the mediobasal hypothalamus is the primary site of action of estradiol in inducing the preovulatory gonadotropin-releasing hormone surge in the ewe. *Endocrinol.*, 139: 1752-1760.
- Couse, J.F. and K.S. Korach, 1999. Estrogen receptor null mice: What have we learned and where will they lead us. *Endocr. Rev.*, 20: 358-417.
- De Roux, N., E. Genin, J.C. Carel, F. Matsuda, J.L. Chaussain and E. Milgrom, 2003. Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. *Proc. Natl. Acad. Sci., USA*, 100: 10972-10976.
- Dhillon, W.S., O.B. Chaudhri, M. Patterson, E.L. Thompson, K.G. Urphy, M.K. Badman, B.M. McGowan, V. Amber, S. Patel, M.A. Ghatei and S.R. Bloom, 2005. Kisspeptin-54 stimulates the hypothalamic-pituitary gonadal axis in human males. *J. Clin. Endocrinol. Metab.*, 90: 6609-6615.
- Dorling, A.A., M.G. Todman, K.S. Korach and A.E. Herbison, 2003. Critical role for estrogen receptor alpha in negative feedback regulation of gonadotropin-releasing hormone mRNA expression in the female mouse. *Neuroendocrinology*, 78: 204-209.
- Dungan, H.M., D.K. Clifton and R.A. Steiner, 2006. Minireview: Kisspeptin neurons as central processors in the regulation of gonadotropin-releasing hormone secretion. *Endocrinology*, 147: 1154-1158.
- Estrada, K.M., C.M. Clay, S. Pompolo, J.T. Smith and I.J. Clarke, 2006. Elevated Kiss-1 expression in the arcuate nucleus prior to the cyclic preovulatory gonadotropin-releasing hormone/luteinising hormone surge in the ewe suggests a stimulatory role for kisspeptin in oestrogen-positive feedback. *J. Neuroendocrinol.*, 18: 806-809.
- Foradori, C.D., M. Amstalden, R.L. Goodman and M.N. Lehman, 2006. Colocalisation of DYN and neurokinin B immunoreactivity in the arcuate nucleus and median eminence of the sheep. *J. Neuroendocrinol.*, 18: 534-541.
- Foradori, C.D., L.M. Coolen, M.E. Fitzgerald, D.C. Skinner, R.L. Goodman and M.N. Lehman, 2002. Colocalization of progesterone receptors in the parvicellular DYN neurons of the ovine preoptic area and hypothalamus. *Endocrinology*, 143: 4366-4374.
- Franceschini, I., D. Lomet, M. Cateau, G. Delsol, Y. Tillet and A. Caraty, 2006. Kisspeptin immunoreactive cells of the ovine preoptic area and arcuate nucleus co-express estrogen receptor alpha. *Neurosci. Lett.*, 401: 225-230.
- Funes, S., J.A. Hedrick, G. Vassileva, L. Markowitz, S. Abbondanzo, A. Golovko, S. Yang, F.J. Monsma and E.L. Gustafson, 2003. The KiSS-1 receptor GPR54 is essential for the development of the murine reproductive system. *Biochem. Biophys. Res. Commun.*, 312: 1357-1363.
- Goodman, R.L., L.M. Coolen, G.M. Anderson, S.L. Hardy, M. Valent, J.M. Connors, M.E. Fitzgerald and M.N. Lehman, 2004. Evidence that DYN plays a major role in mediating progesterone negative feedback on gonadotropin-releasing hormone neurons in sheep. *Endocrinology*, 145: 2959-2967.
- Goodman, R.L., M.N. Lehman, J.T. Smith, L.M. Coolen, C.V. De Oliveira, M.R. Jafarzadehshirazi, A. Pereira, J. Iqbal, A. Caraty, P. Ciofi and I.J. Clarke, 2007. Kisspeptin neurons in the arcuate nucleus of the ewe express both dynorphin A and neurokinin B. *Endocrinology*, 148: 5752-5760.
- Gottsche, M.L., M.J. Cunningham, J.T. Smith, S.M. Popa, B.V. Acohido, W.F. Crowley, S. Seminara, D.K. Clifton and R.A. Steiner, 2004. A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology*, 145: 4073-4077.
- Goubillon, M.L., B. Delaleu, Y. Tillet, A. Caraty and A.E. Herbison, 1999. Localization of estrogen-receptive neurones projecting to the GnRH neuron-containing rostral preoptic area of the ewe. *Neuroendocrinol.*, 70: 228-236.

- Herbison, A.E. and J.R. Pape, 2001. New evidence for estrogen receptors in gonadotropin-releasing hormone neurons. *Front Endocrinol.*, 22 (4): 292-308.
- Messenger, S., E.E. Chatzidaki, A.G. Hendrick, D. Zahn, J. Ixon, R.R. Thresher, I. Malinge, D. Lomet, M.B. Carlton, W.H. Colledge, A. Caraty and S.A. Aparicio, 2005. Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *Proc. Natl. Acad. Sci., USA*, 102: 1761-1766.
- Navarro, V.M., J.M. Castellano, R. Fernandez-Fernandez, S. Tovar, J. Roa, A. Mayen, R. Nogueiras, M.J. Vazquez, M.L. Barreiro, P. Magni, E. Aguilar, C. Dieguez, L. Pinilla and M. Tena-Sempere, 2005. Characterization of the potent luteinizing hormone-releasing activity of KiSS-1 peptide, the natural ligand of GPR54. *Endocrinology*, 146: 156-163.
- Navarro, V.M., J.M. Castellano, R. Fernandez-Fernandez, M.L. Barreiro, J. Roa, J.E. Sanchez-Criado, E. Aguilar, C. Dieguez, L. Pinilla and M. Tena-Sempere, 2004. Developmental and hormonally regulated messenger ribonucleic acid expression of KiSS-1 and its putative receptor, GPR54, in rat hypothalamus and potent luteinizing hormone-releasing activity of KiSS-1 peptide. *Endocrinology*, 145: 4565-4574.
- Plant, T.M., S. Ramaswamy and M.J. DiPietro, 2006. Repetitive activation of hypothalamic G protein-coupled receptor 54 with intravenous pulses of kisspeptin in the juvenile monkey (*Macaca mulatta*) elicits a sustained train of gonadotrophin-releasing hormone discharges. *Endocrinology*, 147: 1007-1013.
- Pompolo, S, A. Pereira, K.M. Estada and I.J. Clarke, 2006. Colocalization of kisspeptin and gonadotropin-releasing hormone in the ovine brain. *Endocrinology*, 147: 804-810.
- Roa, J. and M. Tena-Sempere, 2007. Kiss-1 system and reproduction: Comparative aspects and roles in the control of female gonadotropic axis in mammals. *Gen. Comp. Endocrinol.*, 153: 132-140.
- Seminara, S.B., S. Messenger, E.E. Chatzidaki, R.R. Thresher, J.J. Aciermo, J.K. Shagoury, Y. Bo-Abbas, W. Kuohung, K.M. Schwino, A.G. Hendrick, D. Zahn, J. Dixon, U.B. Kaiser, S.A. Slaugenhaupt, J.F. Gusella, S. O'Rahilly, M.B. Carlton, J.W. Crowley, S.A. Aparicio and W.H. Colledge, 2003. The GPR54 gene as a regulator of puberty. *N., Engl., J., Med.*, 349: 1614-1627.
- Smith, J.T. and I.J. Clarke, 2007. Kisspeptin expression in the brain: Catalyst for the initiation of puberty. *Rev. Endocr. Metab. Disord.*, 8: 1-9.
- Smith, J.T., C.M. Clay, A. Caraty and I.J. Clarke, 2007. KiSS-1 messenger ribonucleic acid expression in the hypothalamus of the ewe is regulated by sex steroids and season. *Endocrinology*, 148: 1150-1157.
- Thompson, E.L., M. Patterson, K.G. Murphy, K.L. Smith, W.S. Dhillon, J.F. Todd, M.A. Ghatei and S.R. Bloom, 2004. Central and peripheral administration of kisspeptin-10 stimulates the hypothalamic-pituitary-gonadal axis. *J. Neuro Endocrinol.*, 16: 850-858.