

## Effect of Sodium Bentonite in Broiler Chickens Fed Diets Contaminated with Aflatoxin B<sub>1</sub>

<sup>1</sup>H. Kermanshahi, <sup>2</sup>A.R. Hazegh and <sup>2</sup>N. Afzali

<sup>1</sup>Department of Animal Science, Faculty of Agricultural, Ferdowsi University of Mashhad, Excellence Center, P.O. Box 91775-1163, Mashhad, Iran

<sup>2</sup>Faculty of Agriculture, Birjand University, Iran

---

**Abstract:** In an *in vivo* study, Sodium Bentonite (SB) was evaluated for its ability to reduce the deleterious effects of Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) in broiler chickens. In this study, 288 days old Ross male broiler chickens were used and randomly assigned to 9 treatment groups, with 4 replicates of 8 birds each. Three levels of Aflatoxin B<sub>1</sub>, AFB<sub>1</sub> (0, 500 and 1000 ppb) and 3 levels of SB (0.0, 0.5 and 1.0%) were fed to chickens from 0-42 days of age. Feeding levels of AFB<sub>1</sub> alone significantly decreased ( $p < 0.05$ ) feed intake, body weight and body weight gain, while, SB levels alone or in combination with AFB<sub>1</sub>, had no effect on the mentioned traits. No significant difference was seen for feed conversion ratio during periods of 0-42 days of age. In the diet containing AFB<sub>1</sub>, relative weight of the carcass, thigh and breast were significantly decreased ( $p < 0.05$ ). Adding AFB<sub>1</sub> alone into the diets significantly increased ( $p < 0.05$ ) serum ALT, AST and LDH enzymes and decreased serum GGT enzyme ( $p < 0.05$ ). However, SB alone or in combination with AFB<sub>1</sub> had no effect on the above mentioned enzymes. It is suggested that sodium bentonite at the levels of 0.5-1.0% might be used for reducing the adverse effects of aflatoxins in broiler chickens.

**Key words:** Aflatoxin B<sub>1</sub>, sodium bentonite, performance, blood enzymes, broilers

---

### INTRODUCTION

Mycotoxins are secondary toxic metabolites formed by certain fungi growing on Food and animal feeds usually belonging to the genus of *Aspergillus flavus* and *Aspergillus parasiticus* (Tedesco *et al.*, 2004) and can happen as natural contaminants of poultry feed (Leeson *et al.*, 1995). Contamination of cereals and grains and their related products with mycotoxins causes food and feed-born intoxicants in human and animals (Meltor, 2001). Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) is the most potent mycotoxins causes a wide range of clinical and sub-clinical problems in poultry (Kermanshahi *et al.*, 2007; Hussein and Brasel, 2001). Many different toxic effects for aflatoxins, including reduced performance, hepatic intoxication, adverse effects on carcass as well as egg shell quality, immunosuppression and carcinogenicity have been reported in poultry (Charmley *et al.*, 1995). Increase in relative weights of liver, kidney, heart, proventriculus, gizzard, spleen and pancreas in broilers consuming aflatoxin contaminated diets, have been reported (Kubena *et al.*, 1990). Liver enlargement and discoloration in broilers have been reported by

Kermanshahi *et al.* (2007). Also, it has been demonstrated that aflatoxins are able to alter concentrations of some blood enzymes (Kubena *et al.*, 1990). Negative effects of aflatoxins in poultry production are both dose and time dependent (Leeson *et al.*, 1995). At present, aflatoxins are considered unavoidable contaminants of feed and food. To combat animal aflatoxicosis and detoxification of mycotoxin-contaminated feeds, different physical methods are recommended but since mycotoxins are heat stable and resistance to heat, current agricultural practices and routine feed processing procedures can not reduce their toxicity (Bassapa and Shanta, 1996; Doyle *et al.*, 1982). In animal nutrition, the use of mycotoxin adsorbents as feed additives is one of the most promising and widely used approaches to reduce the deleterious effect of aflatoxicosis in farm animals. It is good method to reduce carry-over of mycotoxins from the contaminated feeds into animal-derived products (Ramos *et al.*, 1996). Phillips *et al.* (1987, 1988) in *in vitro* and *in vivo* studies reported that Hydrated Sodium Calcium Aluminosilicate (HSCAS) have the high capacity to adsorb aflatoxins in poultry HSCAS is a generic description and it does not uniquely define the material

of use. Most of the classified HSCAS are as natural montmorillonites and bentonites (Ramos and Fernandez, 1997). In many reports, different adsorbents including zeolites, activated charcoal and phyllosilicate minerals have been used to decrease the negative effect of aflatoxins (Devegowda *et al.*, 1996; Rosa *et al.*, 2001). There are many adsorbents in the markets and their capacity to adsorb specific mycotoxins are not clear. Although, from nutritional point of view, bentonite type of binders would supply little nutritive value but, it is the most potent mineral substances to reduce the deleterious effect of aflatoxins in poultry (Miazzo *et al.*, 2000, 2005; Rosa *et al.*, 2001). Therefore, the purpose of this research was to evaluate the capacity of locally available bentonites in male broiler chickens fed diets contaminated with aflatoxin B<sub>1</sub>.

### MATERIALS AND METHODS

Total 288 days old Ross male broiler chickens were divided into 9 treatments of 4 replicates each and randomly assigned to each of 36 floor pens of 1×1 m. All birds had free access to feed and water during the experiment (0-42 days). Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) for this experiment was produced using *Aspergillus parasiticus* strain 2999 according to the method described by Shotwell *et al.* (1966) and West *et al.* (1973). More than 80% of the aflatoxin produced by this strain is B<sub>1</sub> type (Rosa *et al.*, 2001). AFB<sub>1</sub> concentration was determined using TLC according to Afzali (1998) method. Relative weights of edible meat, thigh and breast to live body weight measured at 42 days of age by killing 2 birds from each replicate of treatments. Serum concentrations of Aspartate amino Transferase (AST), Alanine amino Transferase (ALT), Gama Glutamyl Transferase (GGT) and Lactate Dehydrogenase (LDH) were determined at 21 and 42 days of age using commercial available kits (Darman Kave Research Laboratory, Isfahan, Iran). Starter, grower and finisher diets (Table 1) were formulated for 0-14, 14-28 and 28-42 days of age, respectively, according to the Ross 308 manual recommendations. Dietary treatments were consisted of 3 levels of AFB<sub>1</sub> in diet (0.0, 500 and 1000 ppb) and 3 levels of a locally prepared sodium bentonite (0.0, 0.5 and 1.0% SB, Ashkaftook mine, South of Khorasan, Iran). The SB composition was SiO<sub>2</sub>, 58.80%; Al<sub>2</sub>O<sub>3</sub>, 18.50%; Fe<sub>2</sub>O<sub>3</sub>, 5.00%; CaO, 2.15%; MgO, 2.10%; Na<sub>2</sub>O, 3.50%; K<sub>2</sub>O, 0.45%; TiO<sub>2</sub>, 0.15%; moisture, 9.35%. In order to reach intended AFB<sub>1</sub> concentrations in diets, an appropriate mixture of *Aspergillus parasiticus* culture on rice was added into the basal diet. AFB<sub>1</sub> contaminated diets were fed to broiler chickens from 0-42 days of age. Feed intake and body weight gain recorded weekly and then performance data expressed as week basis.

Table 1: Composition of the experimental diets

Ingredients (%)	Starter (0-14 days)	Grower (14-28 days)	Finisher (28-42 days)
Yellow com	59.97	60.38	64.65
Soybean meal (44%)	30.00	30.00	28.13
Fish meal	6.20	2.50	0.00
Veg. oil	3.46	3.51	3.30
Dicalcium phosphate	1.40	1.46	1.71
Oyster shell	1.10	1.14	1.25
Vitamin premix <sup>2</sup>	0.25	0.25	0.25
Mineral premix <sup>3</sup>	0.25	0.25	0.25
Salt	0.25	0.25	0.25
DL-Methionine	0.12	0.19	0.16
L-Lysine	0.00	0.07	0.05
Sodium Bentonite (SB) <sup>1</sup>	0.00	0.00	0.00
Aflatoxin B <sub>1</sub> (ppb) (AFB <sub>1</sub> )	0.00	0.00	0.00
Total	100.00	100.00	100.00
<b>Calculated composition</b>			
Metabolic energy (kcal kg <sup>-1</sup> )	3000.00	3000.00	3000.00
Crude protein (%)	22.00	20.00	18.00
Calcium (%)	1.00	0.90	0.90
Available phosphorus (%)	0.50	0.45	0.45
Lysine (%)	1.20	1.09	0.92
Methionine + cystine (%)	0.81	0.81	0.72
Tryptophan (%)	0.22	0.20	0.18
<b>Arrangements of 9 treatments in each period (AFB<sub>1</sub> (ppb))</b>			
SB (%)	0	500	1000
0	0.0 + 0	0.0 + 500	0.0 + 1000
0.5	0.5 + 0	0.5 + 500	0.5 + 1000
1.0	1.0 + 0	1.0 + 500	1.0 + 1000

<sup>1</sup>In each period, 9 treatment diets prepared as arranged in treatment arrangements, <sup>2</sup>Each kg of vitamin premix contained: vitamin A, 3,600,000 IU; vitamin D<sub>3</sub>, 800,000 IU; vitamin E, 7,200 IU; vitamin K<sub>3</sub>, 800 mg; vitamin B<sub>1</sub>, 720 mg; vitamin B<sub>2</sub>, 2,640 mg; vitamin B<sub>3</sub>, 4,000 mg; vitamin B<sub>5</sub>, 12,000 mg; vitamin B<sub>6</sub>, 1,200 mg; vitamin B<sub>9</sub>, 400 mg; vitamin B<sub>12</sub>, 6 mg; vitamin H<sub>2</sub>, 40 mg; choline chloride, 200,000 mg. <sup>3</sup>Each kg of mineral premix contained: Mn, 40,000 mg, Fe, 20,000 mg, Zn, 40,000 mg; Cu, 4,000 mg, Se, 80 mg

**Statistical analysis:** The 9 treatments were allotted in a completely randomized design, with 3×3 factorial arrangements. All data were analyzed using GLM procedure of SAS and treatment means were compared by Touki test (SAS, 1990).

### RESULTS AND DISCUSSION

**Appearance and performance:** In the 1st and 2nd week of experiment, chickens didn't show any sign of toxicity in their appearance. Gradually, at week 3 and later on, chickens fed AFB<sub>1</sub> showed some sign of aggressions, pale color in their combs and skins, fluffy feathers and retarded growth rate. Chickens receiving SB or SB with AFB<sub>1</sub> had normal appearance.

Data of weekly Feed Intake (FI) is presented in Table 2. Treatments had no effect on Feed intake till the end of week 3. However, chickens receiving AFB<sub>1</sub> alone (500 and 1000 ppb) fed lower feed when compared to other treatments. At week 4-6 FI significantly decreased in birds fed AFB<sub>1</sub> contaminated diets (p<0.05). The lowest FI was seen in birds fed 1000 ppb AFB<sub>1</sub> alone. However, the difference between 500 and 1000 ppb AFB<sub>1</sub> was not

Table 2: Effect of treatments on feed intake of broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite

Treatments <sup>1</sup>		Experimental periods (weeks)					
AFB <sub>1</sub> (ppb)	SB (%)	1	2	3	4	5	6
0	0	127.7	273.4	507.6	753.3 <sup>a</sup>	861.8 <sup>a</sup>	1051.1 <sup>a</sup>
500	0	125.1	255.1	500.3	598.7 <sup>b</sup>	716.9 <sup>b</sup>	740.50 <sup>b</sup>
1000	0	124.2	246.7	496.6	589.8 <sup>b</sup>	665.1 <sup>c</sup>	663.90 <sup>c</sup>
0	0.5	129.2	276.3	507.4	751.9 <sup>a</sup>	880.6 <sup>a</sup>	1075.4 <sup>a</sup>
500	0.5	130.7	276.5	521.6	758.9 <sup>a</sup>	871.6 <sup>a</sup>	1056.7 <sup>a</sup>
1000	0.5	128.7	277.5	502.7	752.3 <sup>a</sup>	875.8 <sup>a</sup>	1058.8 <sup>a</sup>
0	1	133.8	278.6	514.4	741.3 <sup>a</sup>	882.3 <sup>a</sup>	1081.3 <sup>a</sup>
500	1	130.6	270.6	515.8	735.4 <sup>a</sup>	885.3 <sup>a</sup>	1060.9 <sup>a</sup>
1000	1	130.7	274.2	519.9	759.8 <sup>a</sup>	872.2 <sup>a</sup>	1068.4 <sup>a</sup>

<sup>1</sup>AFB<sub>1</sub>: Aflatoxin B<sub>1</sub>; SB: Sodium Bentonite; Results for each treatments are obtained from 32 chickens in each treatment with ±SEM = 11.7; <sup>a,b,c</sup>In each column, means with different superscripts are significantly different (p<0.05)

Table 3: Effect of treatments on body weight gain of broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite

Treatments <sup>1</sup>		Experimental periods (weeks)					
AFB <sub>1</sub> (ppb)	SB (%)	1	2	3	4	5	6
0	0	95.0	190.1	368.0 <sup>a</sup>	504.7 <sup>a</sup>	459.2 <sup>a</sup>	520.6 <sup>a</sup>
500	0	91.9	170.0	337.8 <sup>b</sup>	355.8 <sup>b</sup>	365.6 <sup>b</sup>	347.5 <sup>b</sup>
1000	0	89.5	166.3	323.5 <sup>b</sup>	349.6 <sup>b</sup>	327.3 <sup>b</sup>	304.4 <sup>b</sup>
0	0.5	97.9	196.7	368.3 <sup>a</sup>	500.4 <sup>a</sup>	478.4 <sup>a</sup>	521.9 <sup>a</sup>
500	0.5	96.0	194.8	366.1 <sup>a</sup>	498.6 <sup>a</sup>	458.1 <sup>a</sup>	520.0 <sup>a</sup>
1000	0.5	95.2	192.4	366.3 <sup>a</sup>	486.9 <sup>a</sup>	472.3 <sup>a</sup>	510.2 <sup>a</sup>
0	1	97.3	195.6	369.1 <sup>a</sup>	480.9 <sup>a</sup>	475.0 <sup>a</sup>	523.6 <sup>a</sup>
500	1	95.2	185.1	369.4 <sup>a</sup>	476.6 <sup>a</sup>	478.8 <sup>a</sup>	520.8 <sup>a</sup>
1000	1	95.0	188.3	364.4 <sup>a</sup>	468.9 <sup>a</sup>	463.4 <sup>a</sup>	520.5 <sup>a</sup>

<sup>1</sup>AFB<sub>1</sub>: Aflatoxin B<sub>1</sub>; SB: Sodium Bentonite; Results for each treatments are obtained from 32 chickens in each treatment with ±SEM = 14.5; <sup>a,b,c</sup>In each column, means with different superscripts are significantly different (p<0.05)

different only at 4th week. Addition of SB into contaminated diets increased FI similar to control birds. Data indicated that by increasing levels of AFB<sub>1</sub>, FI decreases more and SB addition alleviated this effect. These results are in agreement with those of Ledoux *et al.* (1999), Santin *et al.* (2003), Kermanshahi *et al.* (2007), Rosa *et al.* (2001) and Miazzo *et al.* (2005). Data of weekly Mean Body Weight (MBW) and weekly Body Weight Gain (BWG) of treatments are presented in Table 3 and 4. The results and trends were similar with those of FI. Mean body weight and body weight gain of chickens fed AFB<sub>1</sub> alone started to decrease from 3rd week and continued to end of the experiment. The deleterious effect of AFB<sub>1</sub> was also changed to normal conditions (control) when SB added into their contaminated diets. The results of MBW and BWG in this experiment are also in agreement with those of other studies (Rosa *et al.*, 2001; Desheng *et al.*, 2005; Miazzo *et al.*, 2005; Kermanshahi *et al.*, 2007). Dersjant-Li *et al.* (2003) reviewed the impact of low concentrations of aflatoxins in poultry diets and suggested that the growth reduction due to aflatoxin contaminated diets can be related to reduction in both feed intake and feed efficiency. In this experiment, AFB<sub>1</sub> did not negatively influence feed conversion ratio (Table 5) that

Table 4: Effect of treatments on mean body weight of broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite

Treatments <sup>1</sup>		Experimental periods (weeks)					
AFB <sub>1</sub> (ppb)	SB (%)	1	2	3	4	5	6
0	0	141.8	331.8	699.9 <sup>a</sup>	1204.5 <sup>a</sup>	1663.8 <sup>a</sup>	2184.4 <sup>a</sup>
500	0	139.0	308.9	646.8 <sup>b</sup>	1002.5 <sup>b</sup>	1368.1 <sup>b</sup>	1715.6 <sup>b</sup>
1000	0	135.9	302.2	625.7 <sup>b</sup>	975.30 <sup>b</sup>	1302.7 <sup>b</sup>	1607.0 <sup>b</sup>
0	0.5	144.6	341.3	709.6 <sup>a</sup>	1210.0 <sup>a</sup>	1688.4 <sup>a</sup>	2210.3 <sup>a</sup>
500	0.5	143.7	338.5	704.6 <sup>a</sup>	1203.1 <sup>a</sup>	1661.3 <sup>a</sup>	2181.3 <sup>a</sup>
1000	0.5	142.3	334.8	701.1 <sup>a</sup>	1188.0 <sup>a</sup>	1660.3 <sup>a</sup>	2170.5 <sup>a</sup>
0	1	144.2	339.8	708.9 <sup>a</sup>	1189.8 <sup>a</sup>	1664.8 <sup>a</sup>	2188.4 <sup>a</sup>
500	1	142.8	327.9	697.3 <sup>a</sup>	1173.9 <sup>a</sup>	1652.7 <sup>a</sup>	2173.4 <sup>a</sup>
1000	1	141.5	329.8	694.3 <sup>a</sup>	1163.1 <sup>a</sup>	1626.6 <sup>a</sup>	2147.0 <sup>a</sup>

<sup>1</sup>AFB<sub>1</sub>: Aflatoxin B<sub>1</sub>; SB: Sodium Bentonite; Results for each treatments are obtained from 32 chickens in each treatment with ±SEM = 24.5; <sup>a,b,c</sup>In each column, means with different superscripts are significantly different (p<0.05)

Table 5: Effect of treatments on Feed Conversion Ratio (FCR) of broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite

Treatments <sup>1</sup>		Experimental periods (weeks)					
AFB <sub>1</sub> (ppb)	SB (%)	1	2	3	4	5	6
0	0	1.34	1.44	1.38	1.51	1.94	2.02
500	0	1.36	1.51	1.48	1.69	1.96	2.16
1000	0	1.39	1.49	1.56	1.69	2.03	2.18
0	0.5	1.32	1.41	1.40	1.51	1.86	2.06
500	0.5	1.37	1.42	1.43	1.52	1.91	2.03
1000	0.5	1.35	1.45	1.38	1.55	1.87	2.08
0	1	1.38	1.44	1.39	1.55	1.88	2.07
500	1	1.37	1.47	1.41	1.54	1.87	2.05
1000	1	1.38	1.46	1.44	1.63	1.89	2.07

<sup>1</sup>AFB<sub>1</sub>: Aflatoxin B<sub>1</sub>; SB: Sodium Bentonite; Results for each treatments are obtained from 32 chickens in each treatment with ±SEM = 0.07; In each column, means with no superscripts are not significantly different (p>0.05)

Table 6: Effect of treatments on relative organ weights (g/100 g of body weight) of broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite

Treatments <sup>1</sup>		Relative organ weights (%)		
AFB <sub>1</sub> (ppb)	SB (%)	Carcass weight	Thigh	Breast
0	0	73.460 <sup>a</sup>	22.250 <sup>a</sup>	22.840 <sup>a</sup>
500	0	67.410 <sup>b</sup>	19.890 <sup>b</sup>	19.660 <sup>b</sup>
1000	0	66.540 <sup>b</sup>	18.840 <sup>b</sup>	18.240 <sup>b</sup>
0	0.5	72.120 <sup>a</sup>	22.040 <sup>a</sup>	21.820 <sup>a</sup>
500	0.5	71.920 <sup>a</sup>	22.010 <sup>a</sup>	21.950 <sup>a</sup>
1000	0.5	71.080 <sup>a</sup>	21.970 <sup>a</sup>	21.810 <sup>a</sup>
0	1	71.720 <sup>a</sup>	22.030 <sup>a</sup>	22.200 <sup>a</sup>
500	1	71.500 <sup>a</sup>	21.960 <sup>a</sup>	21.770 <sup>a</sup>
1000	1	71.310 <sup>a</sup>	21.280 <sup>a</sup>	21.540 <sup>a</sup>
±SEM		1.040	0.410	0.600

<sup>1</sup>AFB<sub>1</sub>, aflatoxin B<sub>1</sub>; SB, Sodium Bentonite, <sup>a, b, c</sup>In each column, means with different superscripts are significantly different (p<0.05)

is in agreement with the results of Edrington *et al.* (1997). Feed Conversion Ratio (FCR) is an index of feed intake and body weight gain and usually wouldn't be considered alone.

**Relative organ weights:** The effect of AFB<sub>1</sub> alone or in combination with SB on carcass weight, thigh and breast yield relative to live body weight is presented in Table 6. The data clearly shows the negative effect of AFB<sub>1</sub> on the

**Table 7: Serum ALT, AST, GGT and LDH concentrations (U/L) in broiler chickens fed diets containing different levels of aflatoxin B<sub>1</sub> and sodium bentonite**

Treatments <sup>1</sup>		ALT <sup>2</sup> (day)		AST (day)		GGT (day)		LDH (day)	
AFB <sub>1</sub> (ppb)	SB (%)	21	42	21	42	21	42	21	42
0	0	57.130 <sup>a</sup>	45.000 <sup>a</sup>	139.250 <sup>a</sup>	109.500 <sup>a</sup>	32.55 <sup>a</sup>	35.59 <sup>a</sup>	318.25 <sup>a</sup>	314.00 <sup>a</sup>
500	0	62.630 <sup>ab</sup>	55.000 <sup>bc</sup>	155.630 <sup>bc</sup>	124.130 <sup>b</sup>	24.58 <sup>b</sup>	26.88 <sup>b</sup>	362.88 <sup>ab</sup>	376.38 <sup>b</sup>
1000	0	68.130 <sup>b</sup>	60.880 <sup>b</sup>	163.130 <sup>c</sup>	135.250 <sup>c</sup>	20.63 <sup>b</sup>	21.50 <sup>c</sup>	391.50 <sup>b</sup>	432.75 <sup>c</sup>
0	0.5	57.630 <sup>a</sup>	46.130 <sup>a</sup>	140.500 <sup>a</sup>	110.880 <sup>a</sup>	32.13 <sup>a</sup>	34.38 <sup>a</sup>	319.00 <sup>a</sup>	317.00 <sup>a</sup>
500	0.5	60.380 <sup>a</sup>	49.250 <sup>ac</sup>	143.380 <sup>a</sup>	114.500 <sup>a</sup>	30.88 <sup>a</sup>	33.46 <sup>a</sup>	326.00 <sup>a</sup>	322.00 <sup>a</sup>
1000	0.5	59.630 <sup>a</sup>	50.380 <sup>ac</sup>	146.630 <sup>ab</sup>	118.250 <sup>ab</sup>	31.00 <sup>a</sup>	32.63 <sup>a</sup>	332.75 <sup>a</sup>	329.25 <sup>ab</sup>
0	1	57.880 <sup>a</sup>	48.500 <sup>ac</sup>	142.000 <sup>a</sup>	112.880 <sup>a</sup>	31.25 <sup>a</sup>	33.75 <sup>a</sup>	321.00 <sup>a</sup>	318.00 <sup>a</sup>
500	1	59.130 <sup>a</sup>	48.750 <sup>ac</sup>	143.630 <sup>a</sup>	116.750 <sup>ab</sup>	31.19 <sup>a</sup>	33.25 <sup>a</sup>	325.13 <sup>a</sup>	324.25 <sup>ab</sup>
1000	1	58.880 <sup>a</sup>	49.750 <sup>ac</sup>	147.250 <sup>ab</sup>	115.880 <sup>ab</sup>	31.80 <sup>a</sup>	32.56 <sup>a</sup>	336.88 <sup>a</sup>	326.13 <sup>ab</sup>
±SEM	-	2.450	3.3700	1.480	18.580	-	-	-	-

<sup>1</sup>AFB<sub>1</sub>: Aflatoxin B<sub>1</sub>, SB: Sodium Bentonite, Results for each treatments are obtained from 32 chickens in each treatment, <sup>2</sup>ALT: Alanine Aminotransferase, AST: Aspartate Aminotransferase, GGT: Gamma Glutamyl Trasferase, LDH: Lactate Dehydrogenase, <sup>a, b, c</sup>In each column, means with different superscripts are significantly different (p<0.05)

mentioned organ weights. AFB<sub>1</sub> significantly decreased carcass weight, thigh and breast yield (p<0.05). This negative effect was cleared when SB added as feed supplement into the experimental diets. Kermanshahi *et al.* (2007) showed that AFB<sub>1</sub> significantly increases some of the organ weights. Liver is the first organ affected by AFB<sub>1</sub> (Leeson *et al.*, 1995). Increase in liver relative weight has also been reported by Kubena *et al.* (1993) during aflatoxicosis. This enhancement in liver weight is usually due to fat deposition in the liver (Leeson *et al.*, 1995). Increase in internal organ weights like liver may decrease relative carcass weight. As seen in this study, the lower values of edible parts of the carcass (most valued parts) brings lots of losses to poultry producers and using SB is one of the economical ways to combat with this problem.

**Blood enzymes:** Serum concentrations of ALT, AST, GGT and LDH for different treatment groups are shown in Table 7. AFB<sub>1</sub> alone in the diet at both 21 and 42 days of age caused a significant increase in serum concentrations of ALT and AST (p<0.05).

The results obtained in this study concerning increase activity of ALT and AST when chickens fed AFB<sub>1</sub> are in agreement with those of many studies (Devegowda *et al.*, 1996; Celik *et al.*, 2005; Kermanshahi *et al.*, 2007). Dafalla *et al.* (1987) also reported an increase in serum concentration of AST and ALT due to AFB<sub>1</sub> administration into the diet. Generally, AST and ALT are intracellular enzymes that do not belong to plasma; so, their appearance in serum indicates cell injury (Coles, 1974).

Increase in serum concentrations of AST and ALT in this experiment, might be a result of hepatocytes injury. Using bentonite brought the ALT and AST enzyme activities back to normal conditions that is also showed by Eraslan *et al.* (2006). AFB<sub>1</sub> alone significantly increased GGT activity in both 21 and 42 days of age (p<0.05).

The lowest level of GGT was seen in chickens fed 1000 ppb AFB<sub>1</sub>. Supplementation of contaminated diets with SB corrected the GGT values similar to that of control treatment in both ages. Our results concerning decrease in GGT activity and AFB<sub>1</sub> effect is similar to those of Jansen *et al.* (2006) and Ersalan *et al.* (2006). Eraslan *et al.* (2006) was also shown that using SB alleviates the effect of AFB<sub>1</sub> on GGT and increases its activity. AFB<sub>1</sub> significantly increased activity of LDH in both 21 and 42 days of age (p<0.05).

This effect was more pronounced in 1000 ppb concentration of AFB<sub>1</sub>. Addition of SB into the contaminated diets corrected the increasing effect of AFB<sub>1</sub> so LDH activity gain decreased and reached to control group.

## CONCLUSION

Under the conditions of this study, it was concluded that feeding aflatoxin B<sub>1</sub> at the used levels can impair feed intake, growth performance and efficiency of carcass yield, thigh and breast meat. Using sodium bentonite showed a positive effect to chickens and nearly cleared the negative effect of aflatoxin B<sub>1</sub>. It seems that using 0.5-1% sodium bentonite into the diets is the acceptable range to combat the negative effects of aflatoxin B<sub>1</sub> in broiler chickens.

## ACKNOWLEDGEMENT

The financial support of the Excellence Center of Animal Science, Ferdowsi University of Mashhad, Iran is gratefully acknowledged by the authors.

## REFERENCES

- Afzali, N., 1998. Biotechnological method to counteract Aflatoxicosis in broiler breeders. Ph.D Thesis, University of Agricultural Sciences. [dbase.irandoc.ac.ir/00234/00234048.htm](http://dbase.irandoc.ac.ir/00234/00234048.htm).

- Bassapa, S.C. and T. Shanta, 1996. Methods for detoxification of aflatoxins in food and feeds. A critical appraisal. *J. Food Technol. India*, 33 (2): 95-107. [ps.fass.org/cgi/content/full/87/8/1569](http://ps.fass.org/cgi/content/full/87/8/1569).
- Celik, S., Z. Erdogan, S. Erdogan and R. Bal, 2005. Efficacy of Tribasic Copper Chloride (TBCC) to reduce the harmful effects of aflatoxin in broilers. *Turk. J. Vet. Anim. Sci.*, 29: 909-916. [www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20053126743](http://www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20053126743).
- Charmley, L.L., H.L. Trenholm and D.B. Prelusky, 1995. Mycotoxins. Their Origin, Impact and Importance, Insight into Common Methods of Control and Elimination. In: Lyons, T. and K.A. Jacques (Eds.). *Bio-technology in the feed industry. Proc. Alltech's 11th Ann. Symposium*, pp: 41-63. [www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20063209477](http://www.cababstractsplus.org/abstracts/Abstract.aspx?AcNo=20063209477).
- Coles, E.H., 1974. *Veterinary Clinical Pathology*. 2nd Edn. Saunders, W.B. and Philadelphia. [www.blackwell-synergy.com/doi/pdf/10.1111/j.1532-950X.1977.tb01641.x](http://www.blackwell-synergy.com/doi/pdf/10.1111/j.1532-950X.1977.tb01641.x).
- Dafalla, R., A.I. Yagi and S.E.I. Adam, 1987. Experimental aflatoxicosis in Hybro-type chicks: Sequential changes in growth and serum constituents and histopathological changes. *Vet. Hum. Toxicol.*, 29: 222-226. [www.ncbi.nlm.nih.gov/pubmed/3604040](http://www.ncbi.nlm.nih.gov/pubmed/3604040).
- Dersjant-Li, Y., M.W.A. Verstegen and W.J.J. Gerrits, 2003. The impact of low concentrations of aflatoxin, deoxynivalenol of fumonisin in diets on growing pigs and poultry. *Nutr. Res. Rev.*, 16: 223-239. [journals.cambridge.org/abstract\\_S0954422403000167](http://journals.cambridge.org/abstract_S0954422403000167).
- Desheng, Q., L. Fan, Y. Yanhu and Z. Niya, 2005. Adsorption of aflatoxin B<sub>1</sub> on montmorillonite. *Poult. Sci.*, 84: 959-961. [ps.fass.org/cgi/content/abstract/84/6/959](http://ps.fass.org/cgi/content/abstract/84/6/959).
- Devegowda, G., B.I.R. Aravind and P. Morton, 1996. *Saccharomyces cerviciae* mannan-oligosaccharides to counteract aflatoxins in broilers. *Proc. Aust. Poult. Sci. Symp.*, 8: 103-106. [www.animalscience.com/upload/additionalfiles/WPSAVerona%5C10675.pdf](http://www.animalscience.com/upload/additionalfiles/WPSAVerona%5C10675.pdf).
- Doyle, M.P., R.S. Applebaum, R.E. Bracket and E.H. Marth, 1982. Physical, chemical and biological degradation of mycotoxins in foods and agricultural commodities. *J. Food Prot.*, 45: 964-971. [www.springerlink.com/index/P3198K9341752321.pdf](http://www.springerlink.com/index/P3198K9341752321.pdf).
- Eraslan, G., D. Essiz, M. Akdogan, E. Karaoz, M. Oncu and Z. Ozyildiz, 2006. Efficacy of dietary sodium bentonite against subchronic exposure to dietary aflatoxin in broilers. *Bull. Vet. Inst. Pulawy*, 50: 107-112. [www.piwet.pulawy.pl/doc/biuletyn\\_50-1/21\\_Eraslan.pdf](http://www.piwet.pulawy.pl/doc/biuletyn_50-1/21_Eraslan.pdf).
- Edrington, T.S., L.F. Kubena, R.B. Harvey and G.E. Rottinghaus, 1997. Influence of a superactivated charcoal on the toxic effects of aflatoxin or t-2 toxin in growing broilers. *Poult. Sci.*, 76: 1205-1211. [ps.fass.org/cgi/content/abstract/76/9/1205](http://ps.fass.org/cgi/content/abstract/76/9/1205).
- Hussein, H.S. and J.M. Brasel, 2001. Toxicity, metabolism and impact of mycotoxins on humans and animals. *Toxicol.*, 167: 101-134. [www.ncbi.nlm.nih.gov/pubmed/11567776](http://www.ncbi.nlm.nih.gov/pubmed/11567776).
- Jansen, V.R.C., C.E.J. Van Rensburg, J.B.J. Van Ryssen, N.H. Casey and G.E. Rottinghaus, 2006. *In vitro* and *In vivo* assessment of humic acid as an aflatoxin binder in broiler chickens. *Poult. Sci.*, 85: 1576-1583. [cat.inist.fr/?aModele=afficheN&cpsidt=18039145](http://cat.inist.fr/?aModele=afficheN&cpsidt=18039145).
- Kermanshahi, H., M.R. Akbari, M. Maleki and M. Behgar, 2007. Effect of prolonged low level inclusion of aflatoxin B<sub>1</sub> into diet on performance, Nutrient digestibility, histopathology and blood enzymes of broiler chickens. *J. Anim. Vet. Adv.*, 6 (5): 686-692. [www.medwelljournals.com/new/5/detail.php?id=5&jid=java&theme=5&issueno=56&articulo=932](http://www.medwelljournals.com/new/5/detail.php?id=5&jid=java&theme=5&issueno=56&articulo=932).
- Kubena, L.F., R.B. Harvey, W.E. Huff, D.E. Corrier, T.D. Phillips and G.E. Rottinghaus, 1990. Efficacy of a hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and T-2 toxin. *Poult. Sci.*, 69: 1078-1086. [www.ncbi.nlm.nih.gov/pubmed/2235827](http://www.ncbi.nlm.nih.gov/pubmed/2235827).
- Kubena, L.F., R.B. Harvey, W.E. Huff, M.H. Elissalde, A.G. Yersin, T.D. Phillips and G.E. Rottinghaus, 1993. Efficacy of a hydrated sodium calcium aluminosilicate to reduce the toxicity of aflatoxin and diacetoxyscirpenol. *Poult. Sci.*, 72: 51-59. [www.ncbi.nlm.nih.gov/pubmed/8381229](http://www.ncbi.nlm.nih.gov/pubmed/8381229).
- Ledoux, D.R., G.E. Rottinghaus, A.J. Bermudez and M. Alonso-Debolt, 1999. Efficacy of a hydrated sodium calcium aluminosilicate to ameliorate the toxic effects of aflatoxin in broiler chicks. *Poult. Sci.*, 77: 204-210. [grande.nal.usda.gov/ibids/index.php?mode2=detail&origin=ibids\\_references&throw=419355-7k-Cached](http://grande.nal.usda.gov/ibids/index.php?mode2=detail&origin=ibids_references&throw=419355-7k-Cached).
- Leeson, S., G. Diaz and J.D. Summers, 1995. *Poultry metabolic disorders and mycotoxins*. University books, Guelph, Ontario, Canada, pp: 249-298. [japr.fass.org/cgi/reprint/4/2/216.pdf](http://japr.fass.org/cgi/reprint/4/2/216.pdf).
- Meltor, S., 2001. Mycotoxins in feed: A global challenge. *Feed Mix.*, 9: 26-28. [www.allaboutfeed.net/article-database/download/mycotoxins-in-feed-a-global-challenge-id954.pdf](http://www.allaboutfeed.net/article-database/download/mycotoxins-in-feed-a-global-challenge-id954.pdf).
- Miazzo, R., C.A.R. Rosa, E.C. De Queiroz Carvalho, C. Magnoli, S.M. Chiacchiera, G. Palacio, M. Saenz, A. Kikot, E. Basaldella and A. Dalcero, 2000. Efficacy of synthetic zeolite to reduce the toxicity of aflatoxin in broiler chickens. *Poult. Sci.*, 79: 1-6. [ps.fass.org/cgi/content/abstract/79/1/1](http://ps.fass.org/cgi/content/abstract/79/1/1).

- Miazzo, R., M.F. Peralta, C. Magnoli, M. Salvano, S. Ferrero, S.M. Chia-cchiera, E.C.Q. Carvalho, C.A.R. Rosa and A. Dalcero, 2005. Efficacy of sodium bentonite as a detoxifier of broiler feed contaminated with aflatoxin and fumunisin. *Poult. Sci.*, 84: 1-8. [www.poultryscience.org/ps/abs/05/p0510001.htm](http://www.poultryscience.org/ps/abs/05/p0510001.htm).
- Phillips, T.D., L.F. Kubena, R.B. Harvey, D.R. Taylor and N.D. HaidelBaugh, 1987. Mycotoxin hazards in agriculture: New approach to control. *J. Anim. Vet. Med. Assoc.*, 190: 1617-1618. [tih.sagepub.com/cgi/content/refs/23/2/83](http://tih.sagepub.com/cgi/content/refs/23/2/83).
- Phillips, T.D., L.F. Kubena, R.B. Harvey, D.R. Taylor and N.D. HaidelBaugh, 1988. Hydrated sodium calcium aluminosilicate: A higher affinity sorbent for aflatoxin. *Poult. Sci.*, 67: 243-247. [www.ncbi.nlm.nih.gov/pubmed/2837754](http://www.ncbi.nlm.nih.gov/pubmed/2837754).
- Rosa, C.A.R., R. Miazzo, C. Magnoli, M. Salvano, S.M. Chiacchiera, S. Ferrero, M. Saenz, E.C.Q. Carvalho and A. Dalcero, 2001. Evaluation of the efficacy of bentonite from the South of Argentina to ameliorate the toxic effects of aflatoxin in broilers. *Poult. Sci.*, 80: 139-144. [ps.fass.org/cgi/content/abstract/80/2/139](http://ps.fass.org/cgi/content/abstract/80/2/139).
- Ramos, A.J., J. Fink-Gremmels and E. Hernandez, 1996. Prevention of toxic effects of mycotoxins by means of non-nutritive adsorbent compounds. *J. Feed Prot.*, 59: 631-641. [cat.inist.fr/?aModele=afficheN&cpsidt=3126413](http://cat.inist.fr/?aModele=afficheN&cpsidt=3126413).
- Ramos, A.J. and E. Fernandez, 1997. Prevention of aflatoxicosis in farm animals by means of hydrated sodium calcium aluminosilicate addition to feed stuffs: A review. *Anim. Feed Sci. Technol.*, 65: 197-206. [linkinghub.elsevier.com/retrieve/pii/S0921448805005006](http://linkinghub.elsevier.com/retrieve/pii/S0921448805005006).
- SAS Institute, 1990. SAS/STAT User's Guide. 4th Edn. Version 6.06, SAS institute Inc., Cary, NC. [www.amazon.com/Users-Guide-Version-Fourth-Volumes/dp/1555443761-174k](http://www.amazon.com/Users-Guide-Version-Fourth-Volumes/dp/1555443761-174k).
- Santin, E., A.C. Paulillo, A. Maiorka, L.S.O. Nakaghi, M. Macari, A.V.F. Da Silva and A.C. Alessi., 2003. Evaluation of the efficacy of *Saccharomyces cerevisiae* cell wall to ameliorate the toxic effects of aflatoxin in broilers. *Int. J. Pult. Sci.*, 2 (5): 341-344. [www.pjbs.org/ijps/fin110.pdf](http://www.pjbs.org/ijps/fin110.pdf).
- Shotwell, O.L., C.W. Hesseltine, R.D. Stubblefield and W.G. Sorenson, 1966. Production of aflatoxin on rice. *Applied Microbiol.*, 14: 425-428. [aem.asm.org/cgi/reprint/14/3/425.pdf](http://aem.asm.org/cgi/reprint/14/3/425.pdf).
- Tedesco, D., S. Steidler, S. Galletti, M. Tamani, O. Sanzogni and L. Ravarotto, 2004. Efficacy of silymarine-phospholipid complex in reducing the toxicity of aflatoxin B<sub>1</sub> in broiler chicks. *Poult. Sci.*, 83: 1839-1843. [www.ncbi.nlm.nih.gov/pubmed/15554059](http://www.ncbi.nlm.nih.gov/pubmed/15554059).
- West, S., R.D. Wyatt and P.B. Hamilton, 1973. Improved yield of aflatoxin by incremental increases of temperature. *Applied Microb.*, 25: 1018-1019. [www.pubmedcentral.nih.gov/articlerender.fcgi?artid=380959](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=380959).