

A Hermaphroditic Specimen of Chub Mackerel *Scomber japonicus* in the Dardanelles, Turkey

Ugur Özekinci, Adnan Ayaz, Ugur Altinagaç, Özgür Cengiz and Alkan Öztekin
Department of Fishing and Processing Technologies, Faculty of Fisheries,
Çanakkale Onsekiz Mart University, Çanakkale, Turkey

Abstract: A hermaphroditic individual of the Chub mackerel, *Scomber japonicus*, caught in the Dardanelles is described in this study. This specimen is the first record of hermaphroditic Chub mackerel for the Mediterranean Sea.

Key words: Scomberidae, *Scomber japonicus*, hermaphroditism, chub mackerel, dardanelles, Mediterranean sea

INTRODUCTION

Fish migrate in large schools during certain periods of the year through the Dardanelles (Kocatas and Bilecik, 1992). One of the main migrating fish species through this strait is Chub mackerel, *Scomber japonicus* Houttuyn. It is widespread over the coastal waters of the tropical and subtropical regions of the Pacific, Indian, Atlantic Ocean, the Mediterranean and Black seas. It can undertake long migrations and is vertically distributed from the sea surface up to 300 m (Collette and Nauen, 1983).

MATERIALS AND METHODS

During a study of the hook selectivity, 23 individuals of Chub mackerel were caught in 18 March 2009 in the Dardanelles. Fish were identified based on Collette and Nauen (1983) and scientific names of species were checked with the fish base (Froese and Pauly, 2009). Fish were measured to the nearest cm (total length) and weighed to the nearest g.

RESULTS AND DISCUSSION

Internal examination of fish samples revealed that one individual was hermaphrodite, bearing an ovary and a testis. The tissues of ovary and testis were easily discernible as indicated by clearly divided regions in the same lobe of the hermaphrodite gonad (Fig. 1). The total length and body weight of the hermaphrodite individual were 24.5 cm and 145.09 g, respectively. The length and body weight of the rest of the fish samples ranged between 22.4-27.5 cm and 107.44-206.09 g, respectively. The weight of the hermaphrodite gonad was 1.35 g.




Fig. 1: A hermaphroditic Chub mackerel and their gonads (Ov, ovary and T, testis)

Normal hermaphroditism occurs in species in which the individual is born as a male and changes sex to a female (Protandry) or born as a female and then changes sex to a male (Protogyny) (Atz, 1964). All other forms of sex changes are termed abnormal hermaphrodites or intersexes (Atz, 1964; Devlin and Nagahama, 2002). Abnormal hermaphroditism has been reported for several fish species and both environmental factors and endogenous processes have been suggested to potentially affect sex-determination in fishes (Devlin and Nagahama, 2002).

Abnormal hermaphroditism in scombrids fishes were reported earlier for *Scomber scombrus* Lin. (Stewart, 1891), *Rastrelliger kanagurta* Cuvier (Prabhu and Raja, 1959), *Katsuwonus pelamis* Lin. (Raju, 1960; Uchida, 1961), *Thunnus orientalis* Temminck and Schlegel (Sawada *et al.*, 2002) and *Thunnus thynnus* Lin. (Caprioli *et al.*, 2007). Information on the abnormal

hermaphroditism of Chub mackerel was previously reported only from the Japan sea (Takizaki, 1953; Okiyama and Kawaguchi, 1974).

CONCLUSION

This specimen is the first record of abnormal hermaphroditic Chub mackerel for the Mediterranean and Black sea.

ACKNOWLEDGEMENTS

This study was carried out financial support of TUBITAK TOVAG (Project No. 106O097). The authors would like to thank Mr. Cahit Ceviz for their assistance in the field research. We are grateful to Dr. Umur Önal for his help.

REFERENCES

- Atz, J.W., 1964. Intersexuality in Fishes. In: Armstrong, C.N. and A.J. Marshall (Eds.). Intersexuality in Vertebrates Including Man, Academic Press, London and New York, pp: 145-232.
- Caprioli, R., C. Costa, E. Cataldi, C. Boglione and S. Cataudella, 2007. First record of female germ cells in the testis of a wild Mediterranean Northern bluefin tuna (*Thunnus thynnus* L., 1758). J. Fish Biol., 71:620-622. <http://dx.doi.org/10.1111/j.1095-8649.2007.01509.x>.
- Collette, B.B. and C.E. Nauen, 1983. FAO species catalogue, vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. FAO Fisheries Synopsis No. 125, pp: 2-137. Rome: <ftp://ftp.fao.org/docrep/fao/009/ac478e/ac478e00.pdf>.
- Devlin, R.H. and Y. Nagahama, 2002. Sex determination and sex differentiation in fish: An overview of genetic, physiological and environmental influences. Aquaculture, 208: 191-364. DOI: 10.1016/S0044-8486(02)00057-1.
- Froese, R. and D. Pauly, 2009. Fish base. World Wide Web Electronic Publication. www.fishbase.org.
- Kocatas, A. and N. Bilecik, 1992. Aegean Sea and its living resources. The Republic of Turkey, Ministry of Agriculture and Rural Affairs, Bodrum, No. 7, pp: 88 (in Turkish).
- Okiyama, M. and T. Kawaguchi, 1974. A hermatophroditic specimen of mackerel *Scomber japonicus* from the Japan sea. Japanese J. Ichthyol., 21: 168.
- Prabhu, M.S. and A. Raja, 1959. An instance of hermaphroditism in the Indian mackerel *Rastrelliger canagurta* (Cuvier). Curr. Sci., 28: 73-74. http://www.ias.ac.in/j_archive/curresci/28/2/73-74/viewpage.html.
- Raju, G., 1960. A case of hermaphroditism and some other gonadal abnormalities in the skip jack *Katsuwonus pelamis*. J. Mar. Biol. Assoc. India, 2: 95-102.
- Sawada, Y., M. Seoka, T. Okada, S. Miyashita, O. Murata and H. Kumai, 2002. Hermaphroditism in a captive-raised pacific bluefin tuna. J. Fish Biol., 60 (1): 263-265. DOI: 10.1006/jfbi.2001.1831. www.blackwell-synergy.com/doi/abs/10.1111/j.1095-8649.2002.tb02405.x.
- Stewart, C., 1891. On a hermaphroditic mackerel *Scomber scombrus*. J. Lin. Soc. London (Zoology), 24: 70.
- Takizaki, Y., 1953. On the accidental hermaphroditism in *Scomber japonicus*. Collect. Breed., 15: 63-64.
- Uchida, R.N., 1961. Hermaphroditic skip jack. Pasific Sci., 15: 294-296.