

## Development of Mussel (*M. galloprovincialis* L., 1819) Seed on Different Combined Collectors Used on Raft System, in Sinop, Black Sea

Sedat Karayücel, Meryem Yesim Çelik, Ismihan Karayücel and Gökhan Erik  
Department of Aquaculture, Faculty of Fisheries, Sinop University, Sinop, Turkey

**Abstract:** Raft system was practiced for mediterranean mussel (*Mytilus galloprovincialis*) culture in Sinop Region, Black sea. Four different types of collectors (A type collector was in 14 mm diameter nylon rope which is suspended from the the raft in known mussel settlement time. B type collector was in 14 mm diameter nylon rope which is suspended one month early from the known mussel settlement time. C type collector was in 22 mm diameter and made from old ship rope. D type collector was in 19 mm diameter and made from old anchovy net) was conducted to investigate, effect of collector type on settlement. The results showed that collector types had a significant effect spat settlement. Both shape and surface structure of the collector was found significant ( $p < 0.05$ ). The filamentous and thready surfaces of C and D type collectors were preferred by spats for initial settlement. D type collector had a highest density with  $6.37 \pm 0.60$  ind  $\text{cm}^{-2}$ . At the end of 8 months experimental period, the number of spats on A, B, C and D type collectors were found as  $1218 \pm 7.90$ ,  $1344 \pm 6.80$ ,  $3480 \pm 5.10$  and  $3800 \pm 5.40$  ind  $\text{m}^{-1}$ , respectively.

**Key words:** Raft culture, spat, collector types, settlements, Sinop, Black Sea

### INTRODUCTION

Turkish aquaculture sector can be characterized by a low number of species and production diversity, mainly based on small family-owned farms in inland and big farms in the sea. Total fisheries production is 772.323 ton. Aquaculture production has grown steadily over the years. Aquaculture production was reached 139.873 ton while mussel production was 1.100 ton (0.79%) in 2007 (TUIK, 2008).

Mussel seed is critical for the development of industrial mussel cultivation (Fuentes and Molaes, 1994). Many factors, both physical and biological, can influence mussel settlement in marine systems (Pineda, 2000). Once larvae are competent to settle, the presence of suitable substrata can be indicated by either biotic or abiotic cues, for example surface chemical characteristics, biofilm presence, orientation of the substratum, sunlight, texture and the presence of conspecifics, especially if the species shows gregariousness (Stamps and Krishnan, 1990; Harder *et al.*, 2002). These cues and the biological and physical factors that influence settlement operate at different spatial and temporal scales (Bertness *et al.*, 1996) and the high variability frequently observed in settlement rates can be explained by variation in this wide range of factors (Balch and Scheibling, 2000; Jeffery and Underwood, 2000).

The important factors affecting the spat settlement are composition and textural of materials. Settlement is defined as the point when an individual first takes up permanent residence on the substratum (Connell, 1985), although the mussel settlement process is dynamic and may involve various settlement substrates and transitions to new substrates (Alfaro, 2006). It was emphasized that in point of composition, the following characteristics are surface free energy (Nishida *et al.*, 2003), wettability (Alfred *et al.*, 2005) and associated fouling (Alfaro *et al.*, 2006) and in point of textural properties, the available surface area (Walter and Liebezeit, 2003) and the thickness of filaments (King *et al.*, 1990; Pulfrich, 1996; Lekang *et al.*, 2003; Walter and Liebezeit, 2003) are important factors in larval settlement. The physicochemical characteristics of collector ropes can impact on the strength of seed attachment, thereby modifying the probability of detachment by physical disturbance (Lekang *et al.*, 2003). Collection of natural spat is very important for the production of blue mussels, regardless of the type of production technology used (Lekang *et al.*, 2003).

Longline system was experienced for both spat collection and production in the Black Sea. The results showed that spat could be collected efficiently on 16 mm diameter polypropylene ropes and the collectors should

preferably be installed in January and February and the region was favourable for mussel culture (Karayücel *et al.*, 2002, 2003).

## MATERIALS AND METHODS

Mussel culture in raft system was carried out at depth of 13 m in Sinop (Fig. 1), in the Black Sea, Turkey from June 2005 to May 2006.

**System design:** Raft system was mainly constructed from steel pine wood beams and iron sac. For building raft system, 2 sac float with 3.5 mm thickness, 50×75×300 cm dimensions were used. Three steel bar (0.8×12×400 cm) were attached to the float diagonally while 6 pine wood beams (10×10×400 cm) were attached to steel bars by steel screws.

The raft system was moored from 4 side of the raft by using concreted block and 32 mm polypropylene riser (4:1 scope). Each concreted blocks with 10 m of 22 mm openlink ground chain were connected to riser rope. Two rectangle sacs with 3.5 thickness was used as floats and 4 mm of angle iron welded all over corners of sacs to make the float system strong. Then 2 rectangle float were combined by 2 m galvanised pipe from the bottom of floats.

**Experimental conditions:** Four types of spat collector ropes (A type collector was in 14 mm diameter nylon rope which is suspended from the the raft in known mussel settlement time. B type collector was in 14 mm diameter nylon rope which is suspended one month early form the

known mussel settlement time. C type collector was in 22 mm diameter and made from old ship rope. D type collector was in 19 mm diameter and made from old anchovy net) were used for spat settlement experiment (Fig. 2). Each collector types were represented by 5 collector ropes.

Length of 6 m ropes used as spat collectors and 25 cm of wooden pegs (2 cm Ø pipe) inserted crosswise to collector ropes with 30-50 cm intervals and 3 kg concreted weight tied to end of ropes against wave action and tangle of culture rope.

A, C, D types of collectors were hunged on May 2005 while B type of collectors was hunged on April 2005 from raft system with a distance of 0.5 m (Fig. 3).

**Sampling procedure:** During the sampling period mussel weight and length were measured from each type of collectors from June 2005 to May 2006. To determine spat density, triplicate spat samples (selected one each of 4 type collectors) were taken at monthly intervals from A, B, C and D type collectors. In each sampling date, three from each types collectors were selected and a 30 cm section of the collector were grazed.

The collector ropes were lifted gently to prevent loss of spat and the spat were removed from 30 cm of each rope into 20 L tank filled with sea water. The spat were transferred to the laboratory in a box. All spats were scrubbed for encrusting organism (e.g. barnacles, epifauna and seaweeds). They counted according to owing each type collectors. Then mussel samples were taken by a sub-sampling method for biometric measurements. (Quayle and Newkirk, 1989).



Fig. 1: The map of experiment site in Turkey



Fig. 2: Four types of collector ropes (A type collector was in 14 mm diameter nylon rope which is suspended from the raft in known mussel settlement time. B type collector was in 14 mm diameter nylon rope which is suspended one month early from the known mussel settlement time. C type collector was in 22 mm diameter and made from old ship rope. D type collector was in 19 mm diameter and made from old anchovy net)



Fig. 3: Raft system and spat collectors

**Environmental parameters:** Temperature, salinity, chlorophyll a seston (Total Particulate Matter (TPM)), Particulate Inorganic Matter (PIM) and Particulate Organic Matter (POM), were measured monthly from May 2005-2006. Seawater samples were taken at 3 m depth using a nansen bottle at the experimental site. *In situ* water temperature and salinity was measured using a probe (YSI 6600). In the laboratory, triplicate water samples (3 L) were filtered onto Whatman GF/C filters to determine chlorophyll a ( $\mu\text{g L}^{-1}$ ), seston (TPM) ( $\text{mg L}^{-1}$ ) and POM ( $\text{mg L}^{-1}$ ) concentration according to Stirling (1985).

**Morphometric measurements and statistical analyses:**

Growth was estimated from changes in shell length and live weight. Live weight (total weight of mussel) was obtained by blotting animals with tissue paper and weighing to the nearest 0001 g. Shell length (maximum anterior-posterior axis) was measured to the nearest 0.1 mm with a caliper (Seed, 1969). Monthly Specific Growth Rate (SGR%) were found from following formulate:

$$\text{SGR}\% = \left[ \frac{(\ln L_2 - \ln L_1)}{(T_2 - T_1)} \right] \times 100$$

where,  $L_1$  and  $L_2$  are the mean shell lengths at times  $T_1$  and  $T_2$  (Chatterji *et al.*, 1984).

A correlation matrix was used to determine the relationships between the environmental factors and growth parameters. Statistical analyses were carried out by using MINITAB software 13.

**RESULTS**

**Environmental parameters:** Environmental parameters are shown in Fig 4 and 5. The temperature ranged from 7.5-25.05°C with a mean of 14.9±1.7°C. Salinity ranged from 16.8-17.97‰ with a mean of 17.59±0.10‰ and there




Fig. 4: Monthly distribution of mean temperature (a), salinity (b), chlorophyll a (c) from May 2005-2006




Fig. 5: Monthly distribution of mean particulate organic matter (a), particulate inorganic matter (b) and seston (c) from May 2005-2006

was not a clear seasonal pattern. Chlorophyll a peaked in March ( $16.3 \mu\text{g L}^{-1}$ ) as a result of spring algal bloom and decreased reached its lowest value in January




Fig. 6: Monthly mean changes in spat density on all type of the collectors from June 2005 May 2006 (A type collector: 14 mm diameter nylon rope which is suspended from the the raft in known mussel settlement time. B type collector: 14 mm diameter nylon rope which is suspended one month early form the known mussel settlement time. C type collector: 22 mm diameter and made from old ship rope. D type collector: 19 mm diameter and made from old anchovy net)

( $0.53 \mu\text{g L}^{-1}$ ), with a mean of  $3.07 \pm 1.18 \mu\text{g L}^{-1}$ . Monthly seasonal chlorophyll a concentration was significantly different ( $p < 0.05$ ) and in general higher in spring and lower in winter. POM ranged  $1.58-4 \text{ mg L}^{-1}$  with a mean of  $2.42 \pm 0.17 \text{ mg L}^{-1}$ , while seston ranged from  $5.45-6.92 \text{ mg L}^{-1}$ , with a mean of  $6.21 \pm 0.12 \text{ mg L}^{-1}$ . There was a positive relationship between chlorophyll a and POM ( $p < 0.05$ ). Salinity and temperature did not significantly correlate with chlorophyll a, POM and seston ( $p > 0.05$ ). When chlorophyll a, POM and seston peaked, the temperature was minimum. There was a clear seasonal pattern in the temperature but the other environmental factors did not show clear seasonal pattern.

**Growth:** There was higher settlement on the C type collector than others types in 1st and 2 months.

Young spat have a more rounded shape and brighter shell color than older spat. The monthly spat density on all type of the collectors are depicted in Fig. 6.

Mean monthly SGR% on the A, B, C, D type of the collectors was  $20.02 \pm 4.59$ ,  $19.73 \pm 4.38$ ,  $19.61 \pm 4.16$  and  $19.10 \pm 3.41$ , respectively. At the end of 11 months, mean monthly shell length and live weight were  $44.00 \pm 0.65 \text{ mm}$  and  $11.60 \pm 0.31 \text{ g}$  on the A type;  $45.00 \pm 0.68 \text{ mm}$  and  $11.0 \pm 0.26 \text{ g}$  on B type;  $44.40 \pm 0.78 \text{ mm}$  and  $11.8 \pm 0.45 \text{ g}$  on C type;  $44.90 \pm 0.60 \text{ mm}$  and  $11.43 \pm 0.35 \text{ g}$  on D type, respectively (Fig. 7). Length frequencies of spat on all types of collectors for October (A), January (B) and May 2006 (C) are shown in Fig. 8.

The observation showed that barnacles settled in July-August, continued to developed throughout the




Fig. 7: Monthly mean shell length and live weight of spat from July 2005 to May 2006 for A, B, C, D type of collectors




Fig. 8: Length frequencies of spat on all types of collector in October (a), January (b) and May 2006 (c)

summer and autumn and the few larvae of sea snail was observed in September and October on all type of the collectors (A, B, C, D type). The amount of barnacles decreased with depth.

There was a positive relationship between shell length increment and temperature however POM positively correlated with live weight increment ( $p < 0.05$ ).

## DISCUSSION

In marine systems, filamentous substrates such as hydroids and algae facilitate or enhance larval mussel settlement (Bayne, 1964; Standing, 1976; Dean and Hurd, 1980). Present study also showed that there was a significant effect of collector types on spat settlement.

The most of spat settlement were occurred from July-October and new recruitment (seed) were observed rest of the year. But the number of larvae decreased during winter.

As the mussels had been grown under identical conditions, the characteristics of the materials in the collectors seem to be of importance for collection, growth and attachment of the mussels to the material (Lekang *et al.*, 2003). Ropes with filamentous loops record the highest density values followed by ropes with non-filamentous loops (Filgueira *et al.*, 2007). According to Mason and Drinkwater (1981), Pulfrich (1996), Folino-Rorem *et al.* (2006), blue mussel spats prefer thready and filamentous surfaces to settle. Examples of such surfaces are substrates covered with hydroids, filamentous algae or ropes with a thready surface (Alfaro *et al.*, 2006; Filgueira *et al.*, 2007; Peteiro *et al.*, 2007) and could explain the greater settlement densities observed on filamentous structures in rope (Lekang *et al.*, 2003). One of the main factors influencing settlement density on collector ropes is the available surface area (Walter and Liebezeit, 2003). In the present study, mussel attachment to the collectors varied according to collectors types. Greater settlement density was recorded on C (22 mm diameter) and D (19 mm diameter) types with filamentous ropes while in the early time the highest settlement density on C type collectors. Higher mussel settlement densities are obtained on C with greater available surface area, although a filamentous structure enhances the amount of settled individuals. Dense surface structure provides several attachment points for byssal threads. However C type collector had fringed filament, the reason of spat settlement on fringed filament, could be fallen in the long run. So the highest density on C type did not continue to the end of the study. As compared with SGR%, low density was resulted in better growth but not significant. The density of mussels on the collector gives an indication of the competition for area and nutrients. Lekang *et al.* (2003) were declared similar results.

The seasonal pattern of spat development is related to primary production. The large natural river supply of phosphorus and nitrogen, essential nutrients for marine plants and algae, has always made the Black Sea very fertile (Bakan and Büyükgüngör, 2000). So the relatively

high nutrient levels and hence high phytoplankton content of coastal waters of the Black sea support great good growth and mussel farming. Growth rates were highest in autumn and spring and low in winter but resumed from April. Similar results were declared in many studies (Page and Hubbard, 1987; Mallet and Carver, 1989; Lauzon-Guay *et al.*, 2005) and attributed to high water temperature and food availability in the spring and summer. Several studies showed that when chlorophyll a was high (between March and July), SGR% was found highest, too (Chatterji, 1984; Small and Van Stroken, 1990; Karayücel and Karayücel, 1999, 2000; Filgueira *et al.*, 2007).

In the present study, mean shell length was better than those obtained in Scotland's west coast sea lochs because of high food rate and temperature (Karayücel and Karayücel, 2001). The difference on growth and survival rates is influenced to exposure air, environmental conditions, salinity and food availability (Dickie *et al.*, 1984).

### CONCLUSION

The experiment showed clearly that there was a significant effect of collector types on spat settlement due to it's material and structure. Growth was affected by environmental factors. There was not significant effect of collector types on growth.

### REFERENCES

- Alfaro, A.C., 2006. Population dynamics of the green-lipped mussel, *Perna canaliculus*, at various spatial and temporal scales in northern New Zealand. J. Exp. Mar. Biol. Ecol., 334: 294-315.
- Alfaro, A.C., B.R. Copp, D.R. Appleton, S. Kelly and A.G. Jeffs, 2006. Chemical cues promote settlement in larvae of the green-lipped mussel *Perna canaliculus*. Aquac. Int., 14: 405-412.
- Alfred, N., L.K. Ista, M.E. Callow, J.A. Callow, G.P. Lopez and A.S. Clare, 2005. Mussel (*Mytilus edulis*) byssus deposition in response to variations in surface wettability. J. R. Soc. Interface, 3 (6): 37-43.
- Bakan, G. and H. Büyükgüngör, 2000. The Black Sea. Mar. Poll. Bull., 41: 24-43.
- Balch, T. and R.E. Scheibling, 2000. Temporal and spatial variability in settlement and recruitment of echinoderms in kelp beds and barrens in Nova Scotia. Mar. Ecol. Prog. Ser., 205: 139-154.
- Bayne, B.L., 1964. Primary and secondary settlement in *Mytilus edulis* L (Mollusca). J. Anim. Ecol., 33: 513-523.
- Bertness, M.D., S.D. Gaines and R.A. Wahle, 1996. Wind-driven settlement patterns in the acorn barnacle *Semibalanus balanoides*. Mar. Ecol. Prog. Ser., 137: 103-110.
- Chatterji, A., Z.A. Ansari, B.S. Ingole and A.H. Parulekar, 1984. Growth of the Green Mussel, *Perna viridis* L., in A Sea Water Circulating System. Aquaculture, 40: 47-50.
- Connell, J.H., 1985. The consequences of variation in initial settlement vs. post-settlement mortality in rocky intertidal communities. J. Exp. Mar. Biol. Ecol., 93: 11-45.
- Dean, T.A. and L.E. Hurd, 1980. Development in an estuarine fouling community: The influence of early colonists on later arrivals. Oecologia, 46: 295-301.
- Dickie, L.M., P.R. Boudreau and K.R. Freeman, 1984. Influence of Stock and Site on Growth and Mortality in the Blue Mussel (*Mytilus edulis*). Can. J. Fish. Aquat. Sci., 41: 134-140.
- Filgueira, R., L.G. Peteiro, U. Labarta and M.J. Fernández-Reiriz, 2007. Assessment of spat collector ropes in Galician mussel farming. Aquacult. Eng., 37: 195-201.
- Fuentes, J. and J. Molares, 1994. Settlement of the mussel *Mytilus galloprovincialis* on collectors suspended from rafts in the Ria de Arousa (NW of Spain): Annual pattern and spatial variability. Aquaculture, 122: 55-62.
- Folino-Rorem, N., J. Stoeckel, E. Thom and L. Page, 2006. Effects of artificial filamentous substrate on zebra mussel (*Dreissena polymorpha*) settlement. Biol. Invasions, 8: 89-96.
- Harder, T., C. Lam and P.Y. Qian, 2002. Induction of larval settlement in the polychaete *Hydroides elegans* by marine biofilms: An investigation of monospecific diatom films as settlement cues. Mar. Ecol. Prog. Ser., 229: 105-112.
- Jeffery, C.J. and A.J. Underwood, 2000. Consistent spatial patterns of arrival of larvae of the honeycomb barnacle *Chamaesipho tasmanica* Foster and Anderson in new South Wales. J. Exp. Mar. Biol. Ecol., 252: 109-127.
- Karayücel, S. and I. Karayücel, 1999. Growth, production and biomass in raft cultivated blue mussels (*Mytilus edulis* L.) in 2 Scottish sea lochs. Isr. J. Aquacult./Bamidgeh, 51 (1): 65-73.
- Karayücel, S. and I. Karayücel, 2000. Influence of environmental factors on condition index and biochemical composition in index and biochemical composition in *Mytilus Edulis* L. in cultivated-raft system, in 2 Scottish Lochs. Turk. J. Mar. Sci., 3 (3): 149-166.

- Karayücel, S. and I. Karayücel, 2001. Spat collection, growth and associated problems in mussel (*Mytilus edulis*, L.) in 2 Scottish sea lochs. *Turk. J. Mar. Sci.*, 7: 195-205.
- Karayücel, S., M. Erdem, O. Uyan, S. Saygun and I. Karayücel, 2002. Spat settlement and growth on long-line culture system of the mussel, *Mytilus galloprovincialis*, in the Southern Black Sea. *Isr. J. Aquacult. Bamidgeh*, 54 (4): 163-172.
- Karayücel, S., I. Karayücel, M. Erdem, S. Saygun and O. Uyan, 2003. Growth and production in long-line cultivated mediterranean mussel (*Mytilus galloprovincialis*) in Sinop, Black Sea. *Isr. J. Aquacult. Bamidgeh*, 55 (3): 169-178.
- King, P.A., D. McGrath and W. Britton, 1990. The use of artificial substrates in monitoring mussel (*Mytilus edulis* L.) settlement on an exposed rocky shore in the west of Ireland. *J. Mar. Biol. Ass. U.K.*, 70: 371-380.
- Lauzon-Guay, J., M. Dionne, M.A. Barbeau and D.J. Hamilton, 2005. Effects of seed size and density on growth, tissue-to-shell ratio and survival of cultivated mussels (*Mytilus edulis*) in Prince Edward Island, Canada. *Aquaculture*, 250: 652-665.
- Lekang, O.I., T.K. Stevik and A.M. Bomo, 2003. Evaluation of different combined collectors used in longlines for blue mussel farming. *Aquac. Eng.*, 27: 89-104.
- Mallet, A.L. and C.E.A. Carver, 1989. Growth, mortality and secondary production in natural population of the blue mussel (*Mytilus edulis* L.), *Can. J. Aquat. Sci.*, 46: 1154-1159.
- Mason, J. and J. Drinkwater, 1981. Experiment on Suspended Cultivation on Mussels in Scotland. *Scott. Fish. Inf., Pamphlet. 4. DAFS*, pp: 16.
- Nishida, A., K. Ohkawa, I. Ueda and H. Yamamoto, 2003. Green mussel *Perna viridis* L.: Attachment behaviour and preparation of antifouling surfaces. *Biomol. Eng.*, 20: 381-387.
- Quayle, D.B. and G.F. Newkirk, 1989. Farming Bivalve Molluscs: Methods for Study and Development. In: Sandifer, P.A. (Ed.), *Advances in World Aquaculture*. World Aqua. Soc. IDRC, 1: 294.
- Page, H.M. and D.M. Hubbard, 1987. Temporal Spatiat Patterns of Growth in Mussels *Mytilus edulis* on an Offshore Platform: Relationships to Water Temperature and Food Availability. *J. Exp. Mar. Biol. Ecol.*, 111: 159-179.
- Peteiro, L.G., R. Filgueira, U. Labarta and M.J. Fernández-Reiriz, 2007. Effect of submerged time of collector ropes on the settlement capacity of *Mytilus galloprovincialis* L. *Aquacult. Res.*, 38: 1679-1681.
- Pineda, J., 2000. Linking larval settlement to larval transport: Assumptions, potentials and pitfalls. *Oceanogr. East. Pac.*, 1: 84-105.
- Pulfrich, A., 1996. Attachment and settlement of post-larval mussel (*Mytilus edulis* L.) in the Schleswig-Holstein Wadden Sea. *J. Sea Res.*, 36: 239-250.
- Seed, R., 1969. The ecology of the *Mytilus edulis* L. (Lamellibranchiata) on exposed rocky shores. 2. Growth and mortality. *Ecologia*, 3: 317-350.
- Small, A.C. and M.R. Van Stralen, 1990. Average annual growth and condition of mussels as a function of food source. *Hydrobiologia*, 195: 179-188.
- Standing, J.D., 1976. Fouling Community Structure: Effect of the Hydroid *Obelia Dichotoma* on Larval Recruitment. In: Mackie, G.O. (Ed.). *Coelenterate Ecology and Behaviour*, Plenum Press, New York, pp: 155-164.
- Stamps, J.A. and V.V. Krishnan, 1990. The effect of settlement tactics on territory sizes. *Am. Nat.*, 135: 527-546.
- Stirling, H.P., 1985. *Chemical and Biological Methods of Water Analyses for Aquaculturist*. Institute of Aquaculture, University of Stirling, Scotland, UK, pp: 119.
- TUIK, 2008. Turkish Statistic Department, Fishery Statistic, 2008.11.18. [www.tuik.gov.tr](http://www.tuik.gov.tr).
- Walter, U. and G. Liebezeit, 2003. Efficiency of blue mussel (*Mytilus edulis*) spat collectors in highly dynamic tidal environments of the Lower Saxonian coast (southern North Sea). *Biomol. Eng.*, 20: 407-411.