

Comparative Determination of Serobiochemical Constituents In-Door and Free Grazing Camels

Salah A. AL-Shami

Department of Veterinary Public Health and Animal Husbandry,
College of Veterinary Medicine and Animal Resources, King Faisal University,
P.O. Box 1069, AL-Ahsa 31982, Saudi Arabia

Abstract: Serum concentrations of metabolites and minerals were compared in indoor and free grazing camels. Glucose, triglycerides, cholesterol, immunoreactive insulin and blood urea nitrogen concentrations were significantly higher in indoor than in free grazing camels. Serum Mg and K were significantly higher in free grazing compared to indoor camels. Non-esterified fatty acids concentrations were higher in serum free grazing suggesting that supplementary feeding should be considered in these animals.

Key words: Camel, biochemical constituents, indoor, free-grazing, comparative concentrations

INTRODUCTION

The camel is an important animal in Arabian culture and for various reasons, camels are now transformed from the position of subsistence to one of production (FAO, 1995). The camel in this part of the world was in the past mostly selected for survival to feed the nomads. Camels do not systematically graze on one spot but usually only takes one or 2 bites before moving on, therefore do not destroy pasture (Kanoses, 1977). Due to settlement of many Bedouins, draught or shrinking of grazing land the methods of camel keeping are now fast changing (Mokhtar and El-Hisanonein, 1998) and camel farms are now growing in many parts of Saudi Arabia.

Investigations determining normal values of blood constituents in camels are limited and the way they are affected by nutritional status and other factors seem to be limited (Hassan *et al.*, 1968; Eldirdiri *et al.*, 1987; Abdalla *et al.*, 1988).

Comparison of blood values under different managerial systems seems to be essential as these values may reflect the well-being of the animal and could be used as diagnostic tools in health and disease of animals.

The objective of this study, was to compare some blood constituents in grazing and in-door camels.

MATERIALS AND METHODS

Animals: A total of 105 blood samples were obtained from 2-3 years old male and females, which were naturally grazing in Al-Ahsa area (area extending from Kheras to

Table 1: Raw materials and composition percentages of concentrate mixture given to camels

Raw materials (%)	
Barley	35-45
Soya bean meal	2-4
Wheat bran	15-20
Maize	20-25
Salt	0.5-1
Dried dates	2-3
Dicalcium phosphate	0.5-1
Premix	0.5-0.8
Composition (%)	
Moisture (max)	13
Crude Protein (min)	12
Digestible Protein (min)	9
Total digestible nutrients (min)	72
Fat (min)	5-6
Calcium (min)	1
Total phosphorous (min)	0.5
Ash (max)	9

Salwa, 40000 km²). The predominant pasture plants were *Haloxylon salicornium*, *Cetium elegans*, *Alhagi maurum*, *Anabasis stefera* and *Panicum turgidum* (Gaili *et al.*, 2000). A further 110 samples were obtained from indoor camels kept in 6 farms owned by one person. All animals were group-fed and were given concentrate mixture (Table 1), Rhodes grass (*Chloris gayana*) for roughage and water *ad libitum*.

Blood samples were centrifuged at 3000 rpm and serum was separated and stored at -30°C until analysis.

Laboratory analysis: Serum concentrations of metabolites were determined using the Vet Test 8008 autoanalyser (Sanofi animal health Ltd., England) using specific kits. Serum concentrations of minerals were determined using Shimadzu AA6800 Atomic Absorption Spectrophotometer.

Statistical analysis: Data were analyzed by one way ANOVA, using GLM procedure of SAS (Goodnight *et al.*, 1986). Student t-test was used to detect significant differences at $p < 0.05$.

RESULTS AND DISCUSSION

Serum concentrations of metabolites in camels are given in Table 2. Serum concentration of glucose, cholesterol, immunoreactive insulin and blood urea nitrogen were significantly ($p < 0.05$) higher in indoor camels compared to free grazing camels. Non-esterified fatty acids concentrations were significantly ($p < 0.05$) higher in free grazing camels than in indoor camels. Concentrations of protein and albumin were similar in both groups. Serum minerals in camels are given in Table 3. Magnesium and potassium concentrations were higher in free grazing camels than in indoor camels. Concentration of Na, Ca and P were similar in the 2 groups.

Higher levels of glucose, triglycerides, cholesterol, immunoreactive insulin and blood urea nitrogen in indoor over that of grazing camels indicate that nutritional factor can influence normal values of blood constituents of metabolites (Mokhtar and El-Hisanonein, 1998). Most of these biochemical values obtained were at the upper physiological limit reported elsewhere (Mehrotra and Gupta, 1989; AL-Busadah and Osman, 2001; Al-Suwaigh and AL-Suhaimi, 2005; Mohammed *et al.*, 2007). Rich source of sugars like dried dates may explain the significant increase of serum glucose concentrations in indoor compared to free grazing camels. High intake of grain containing carbohydrate in indoor camels would be expected to increase propionate, the most potent stimulus for production of insulin reflected in the increase of immuno-reactive insulin in this group compared to grazing group (Rabelo *et al.*, 2001; Shingu *et al.*, 2001; Sako *et al.*, 2007). As a result of greater amount of concentrate diet supplementation in indoor camels triglycerides, cholesterol and blood urea nitrogen concentrations were also expected to increase (Sako *et al.*, 2007). The higher values for serum K and Mg in free grazing camels compared to indoor camels could be due to the reason that the free grazing camels graze on plants rich in these minerals.

Serum concentration of Ca, Na and P obtained in this study were similar to those obtained by Haroun *et al.* (2000) and Mohammed *et al.* (2007). The lower non-esterified fatty acids values in the supplemented indoor camels may indicate that animals in this group were able to draw most of the required energy from diet and there was little need for catabolizing endogenous source. On the contrary, the higher non-esterified fatty acids

Table 2: Mean (\pm SD) serum concentration of metabolites indoor and free grazing camels

Parameter	Indoor camels	Free grazing camels
Glucose (mg dL ⁻¹)	130.20 \pm 5.11	105 \pm 4.2*
Total protein (mg dL ⁻¹)	6.85 \pm 0.11	6.62 \pm 0.12
Albumin (mg dL ⁻¹)	3.31 \pm 0.12	3.21 \pm 0.11
Cholesterol (mg dL ⁻¹)	82.1 \pm 2.1	60.5 \pm 1.6*
Triglyceride (mg dL ⁻¹)	95.4 \pm 1.6	60.2 \pm 0.95*
Creatinin (mmol L ⁻¹)	0.42 \pm 1.6	0.36 \pm 0.05
Blood urea nitrogen (mmol L ⁻¹)	5.6 \pm 0.21	3.1 \pm 0.11*
Immunoreactive Insulin (Pmol L ⁻¹)	460 \pm 11	220 \pm 6*
Non-esterified fatty acids (μ mol L ⁻¹)	0.32 \pm 0.03	0.52 \pm 0.06*

* $p < 0.05$

Table 3: Mean (\pm SD) serum concentration of some minerals in indoor and free grazing camels

Parameter	Indoor camels	Free grazing camels
Na (mmol L ⁻¹)	161 \pm 12	146 \pm 11
K (mmol L ⁻¹)	4.2 \pm 0.21	6.8 \pm 0.31*
Ca (mg dL ⁻¹)	9.9 \pm 0.52	10.2 \pm 0.49
P (mg dL ⁻¹)	0.36 \pm 0.02	0.37 \pm 0.02
Mg (mg dL ⁻¹)	0.123 \pm 0.011	0.311 \pm 0.012*

* $p < 0.05$

concentrations seen in free grazing camels would suggest that supplementary feeding should be considered during times when pasture conditions are inadequate.

CONCLUSION

This study concluded that although some of nutrients were available in natural pasture, but supplementary feeding should also be considered.

ACKNOWLEDGEMENT

The author thanks the Deanship of Scientific Research of King Faisal University for financial support.

REFERENCES

- Abdalla, O.M., I.A. Wasfi and F.A. Gadir, 1988. The Arabian race camel normal parameters. Hemogram, Enzymes and Minerals. *Comp. Biochem. Physiol.*, 90A: 237-239.
- Al-Busadah, K.A. and T.E.A. Osman, 2001. Some biochemical and haematological indices in different breeds of camels in Saudi Arabia. *J. Camel Practice Res.*, 7: 149-152.
- AL-Suwaigh, B.R. and E.A. AL-Suhaimi, 2005. Comparative study of some behavioural constituents of plasma in male camels and goats. *J. Camel Practice and Res.*, 21: 141-143.
- Eldirdiri, N.I., H.B. Suliman and A.M. Shommein, 1987. Normal serum activities of some diagnostic enzymes in dromedary camel in Sudan. *Vet. Res. Comm.*, 11: 201-203.
- FAO, 1995. Quarterly Bulletin of statistics. Food and Agriculture Organization, UN Rome, 8: 31-36.

- Gaili, E.S.E., M.M. AL-Eknaah and M.H. Gadek, 2000. Comparative milking performance of three types of Saudi camels (*Camelus dromedarius*). *J. Camel Practice and Res.*, 7: 73-76.
- Goodnight, J.H., J.P. Sal and R.F. Sarle, 1986. In SAS User's Guide. Statistics. SAS Institute Inc., Box 8000 CARY, North Carolina, USA.
- Haroun, E.M., M. Magzoub, O.M. Mohmoud, A.A. AL-Qarawi, A.M. AL-Hawas and O.H. Omer, 2000. Some clinico-pathological aspect of experimental *Trypanosoma evansi* infection in Najdi camels. *J. Camel Practice Res.*, 7: 101-106.
- Hassan, Y.M., H. Hoeller and I.M. Hassan, 1968. Observation on the blood constituents of Camels in the Sudan. *Sud. J. Vet. Sci. Anim. Husb.*, 9: 464-473.
- Kanoses, K.H., 1977. The camel as milk and meat animal. *World Anim. Rev.*, 22: 39-44.
- Mohammed, A.K., A.K.B. Sackey, L.B. Tekdeh and J.O. Gefu, 2007. Serum biochemical values of healthy adult one humped camel (*Camelus dromedarius*) introduced into a sub-humid climate in Shika-Zaria, Nigeria. *J. Camel Practice Res.*, 14 (2): 191-194.
- Mehrotra, V. and M.L. Gupta, 1989. Seasonal variations in certain blood constituents in camel. *Indian J. Anim. Sci.*, 59: 1559-1561.
- Mokhtar, M. and R. EL-Hisanonein, 1998. Proceeding of the International Symposium Constraints and Possibilities of Ruminant Production. Cairo, Egypt.
- Rabelo, E., S.J. Bertics, J. Mackovic and R.R. Grummer, 2001. Strategies for increasing energy density of dry cow diets. *J. Dairy Sci.*, 70: 1543-1549.
- Sako, T., S. Urabe, A. Kusaba, N. Kimura, I. Yoshimura, H. Tazaki, S. Imai, K. Ono and T. Arai, 2007. Comparison of plasma metabolite concentrations and lactate dehydrogenase activity in dogs, cats, horses, cattle and sheep. *Vet. Res. Commun.*, 31: 413-417.
- Shingu, H., K. Hodate, S. Kushibiki, Y. Ueda, A. Watanabe, M. Shimoda and M. Matsumoto, 2001. Profiles of growth hormones and insulin secretion and glucose responses to insulin in growing Japanese Black heifer (beef type): Comparison with Holstein heifer (dairy type). *Comparative Biochem. Physiol.*, 130C: 259-270.