

Distribution of Multidrug Resistant Bacteria in Inanimate Objects Within Stalls in Hidalgo, Mexico

¹H.B. Barrios-Garcia, ²M.A. Flores-Valdez, ³V.M. Martinez-Juarez,
³R. Acosta-Salinas and ⁴J. Acosta-Dibarrat

¹Facultad de Medicina Veterinaria y Zootecnia, Universidad Autonoma de Tamaulipas, Mexico

²Centro de Investigacion y Asistencia en Tecnologia y Diseno del Estado de Jalisco, Mexico

³Medicina Veterinaria y Zootecnia, Instituto de Ciencias Agropecuarias,
Universidad Autonoma del Estado de Hidalgo, Mexico

⁴Centro de Investigacion y Estudios Avanzados en Salud Animal,
Universidad Autonoma del Estado de Mexico, Mexico

Abstract: The emergence of multidrug resistant bacteria is a threat to veterinary and human health. The aim of this study was to determine what would be, if any, the contribution of inanimate objects present in stalls in Hidalgo, Mexico, as a possible reservoir for these microorganisms. About 72 strains are isolated from different locations within stalls, where 54.2% were gram positives and 45.8% were gram negatives and found that 38 out of 72 strains were resistant to at least one antibiotic (52.77%) and 13 out of 38 (34.2%) were resistant to more than 2 antibiotics. The present study underscores the need of carefully assessing what are the correct measures for using and disposing of antibiotics in the veterinary setting.

Key words: Multidrug resistance, antibiotics, stalls, inanimate objects, emergence, veterinary

INTRODUCTION

In recent years, there has been a worldwide increase of drug resistant bacteria; a routine approach is to evaluate antibiotic resistance of clinical isolates. However, considering the frequently reported appearance of pathogens that are able to resist a number of current antibiotics (Aarestrup *et al.*, 1998; Wegener *et al.*, 1997), we thought worth studying organisms isolated from environmental samples, aiming to ascertain whether or not they could be implicated in the rise and maintenance of bacterial drug resistance, possibly by its capacity to horizontally transfer it to acceptor organisms.

Antibiotics presence in the environment due to their production by competing bacteria (D'Costa *et al.*, 2006; Wright, 2007) or as a side product of improper use by humans as well as the presence of naturally resistant saprophytic microorganisms could play a key role as sources for drug resistance transmission to other cells. From the veterinary point of view, it is necessary to consider that in general, antibiotic resistance might be the consequence of a production-oriented approach, whereby

treatment is required to stop a spreading illness or from the perspective of improving profits, such as the use of antibiotics as promoters of growth in some species (Aarestrup *et al.*, 2000; Bager *et al.*, 1999). In this study, we would like to draw attention to the possible role for inanimate objects as reservoir for a persistent-like population which could infect animals or humans.

MATERIALS AND METHODS

We performed a sampling process on material from inert areas in 4 stalls located in Tulancingo Valley, Hidalgo, Mexico, a moderate climate zone with rains in summer and a dry winter; area handling and climate conditions allow maintaining bacterial viability (Makinson and Swan, 2006).

A total of 40 samples are obtained taken from utensils located in handling corrals, including water troughs, troughs, as well as from materials used for routine work like buckets water and milk tanks. We used the 2 most abundant isolated colonies of each sample for further analysis giving a total of 72 specimens.

For the isolation and identification of bacterial species, we followed the protocols recommended in the Bergey's Manual of Systematic Bacteriology. Samples were inoculated onto Blood and MacConkey agar (BBL®) and incubated at 37°C for 24 h; species identification was carried out using biochemical tests.

For antibiotic susceptibility tests, used a diffusion disk assay (Bauer *et al.*, 1966) where each bacterial strain was grown overnight and a suspension was made using 0.85% NaCl, adjusted to a 0.5 reading of Mc Farland's nephelometer (Murray and Zeiting, 1983) and inoculated on Mueller-Hinton agar (BBL, Sparks, MD), supplemented with 5% defibrinated sheep blood. The panel of antibiotics we used is shown in Table 1. The results were interpreted according to the National Committee for Clinical Laboratory Standards (NCCLS, 2002).

RESULTS AND DISCUSSION

We isolated the following strains: *Escherichia coli*, *Citrobacter* sp. *Serratia marcescens*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Staphylococcus aureus*

faecalis, *Enterococcus* sp., *Bacillus subtilis*. A summary of the results is shown in Table 2. Out of the 72 isolated strains, 54.2% were gram positives and 45.8% were gram negatives.

From the first group, *Enterococcus* sp. (12.5%) and *Staphylococcus aureus* coagulase negative (11.11%) were the most commonly isolated species. In the gram negatives group, we frequently found *E. coli* (18%) followed by *P. aeruginosa* (15.2%).

Interestingly, 38 out of 72 strains were resistant to at least one antibiotic (52.77%) and 13 out of 38 (34.2%) were resistant to >2 antibiotics (Table 1). Species showing Multidrug Resistance (MDR) include *S. aureus* coagulase negative (5/8), *E. coli* (3/5) and *P. aeruginosa* (4/8).

It is relevant to note that several places in this area routinely use antibiotics for treatment of their ill animals or as growth enhancers, thus potentially providing a means for bacteria to develop drug resistance and then these organisms can be harbored in inanimate areas among the objects of daily work (Kramer *et al.*, 2006).

Table 1: Detection of multi-resistant gram-positive and gram-negative bacteria

Bacteria	Antibiotic concentration											Total Isolations									
	Amikacin		Ampicillin		Sulfamethoxazol/ trimethoprim		Nalidixic acid		Norfloxacin		Cefotaxime		Erythromycin		Penicilline		Tetracycline		Gentamicine		
	30 µg	10 µg	23.75/1.25 µg	30 µg	10 mg	30 µg	15 µg	10 UI	30 µg	15 µg	10 UI		30 µg	10 µg	10 µg	30 µg	10 µg	10 µg	10 µg	10 µg	10 µg
Gram negative																					
<i>Escherichia coli</i>	0	1	0	3	0	1	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	11
<i>Citrobacter</i> sp.	0	0	0	2	0	0	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	4
<i>Serratia marcescens</i>	0	2	0	0	0	0	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	5
<i>Pseudomonas aeruginosa</i>	3	1	0	0	2	2	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	13
<i>Proteus vulgaris</i>	1	2	1	0	0	0	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	N/D	6
Gram positive																					
<i>Staphylococcus aureus</i> coagulase negative	N/D	N/D	N/D	N/D	N/D	0	3	2	1	0	8										
<i>Micrococcus</i> sp.	N/D	N/D	N/D	N/D	N/D	0	1	1	1	0	6										
<i>Enterococcus faecalis</i>	N/D	N/D	N/D	N/D	N/D	0	0	2	0	1	4										
<i>Enterococcus</i> sp.	N/D	N/D	N/D	N/D	N/D	1	1	3	0	1	9										
<i>Bacillus subtilis</i>	N/D	N/D	N/D	N/D	N/D	0	0	0	0	0	6										
Total	4	6	1	5	2	4	5	7	2	2	38/72										

N/D = Not Determined

Table 2: Origin and occurrence of bacteria isolated from different environmental sites

Subjects	<i>Staphylococcus aureus</i> coagulase negative										Total
	<i>Escherichia coli</i>	<i>Citrobacter</i> sp.	<i>Serratia marcescens</i>	<i>Pseudomonas aeruginosa</i>	<i>Proteus vulgaris</i>	<i>Micrococcus</i> sp.	<i>Enterococcus faecalis</i>	<i>Enterococcus</i> sp.	<i>Bacillus subtilis</i>		
Water troughs	2	0	2	3	0	2	2	3	2	16	
Troughs	3	1	0	2	1	1	0	2	1	13	
Buckets	3	1	1	3	1	1	1	0	1	14	
Water tanks	0	1	2	3	1	0	1	1	0	10	
Milk tanks	0	0	0	1	0	1	0	1	0	5	
Handling corral	3	1	0	1	3	1	0	2	2	14	
Total	11	4	5	13	6	6	4	9	6	72	

CONCLUSION

It is important to conduct a study that assess the distribution and permanency of bacteria as well as the resistance to the current therapeutically relevant agents and use it as an indicator of drug resistance transmission in areas where antibiotics are used as a preventive measure. In this manner, safety measures can be taken, to avoid transferring to other animals a variety of microorganisms to which directly or indirectly humans could have helped in becoming antibiotic resistant, remaining unnoticed in inert areas, as a potential source for widespread tolerance or resistance to multiple drugs.

REFERENCES

- Aarestrup, F.M., A.M. Seyfarth, H.D. Emborg, F. Bager, K. Pedersen and S.E. Jorsal, 2000. Antibiotic use in food-animal production in Denmark. *APUA Newslett.*, 18: 1-3.
- Aarestrup, F.M., F. Bager, N.E. Jensen, M. Madsen, A. Meyling and H.C. Wegener, 1998. Resistance to antimicrobial agents used for animal therapy in pathogenic-, zoonotic- and indicator bacteria isolated from different food animals in Denmark: A baseline study for the Danish Integrated Antimicrobial Resistance Monitoring Programme (DANMAP). *Apmis*, 106: 745-770.
- Bager, F., F.M. Aarestrup, N.E. Jensen, M. Madsen, A. Meyling and H.C. Wegener, 1999. Design of a system for monitoring antimicrobial resistance in pathogenic, zoonotic and indicator bacteria from food animals. *Acta Vet. Scand. Suppl.*, 92: 77-86.
- Bauer, A.W., W.M. Kirby, J.C. Sherris and M. Turck, 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.*, 45: 493-496.
- D'Costa, V.M., K.M. McGrann, D.W. Hughes and G.D. Wright, 2006. Sampling the antibiotic resistome. *Science*, 311: 374-377.
- Kramer, A., I. Schwebke and G. Kampf, 2006. How long do nosocomial pathogens persist on inanimate surfaces: A systematic review. *BMC Infect. Dis.*, 6: 130-130.
- Makinson, C. and J. Swan, 2006. The effect of humidity on the survival of MRSA on Hard Surfaces. *Indoor Built Environ.*, 15: 85-91.
- Murray, R.P. and R.J. Zeiting, 1983. Evaluation of mueller-hinton agar for disk diffusion susceptibility test. *J. Clin. Microbiol.*, 18: 1269-1271.
- NCCLS, 2002. Performance Standards for Antimicrobial Susceptibility Testing; Twelfth Informational Supplement, M100-S12 (M2). National Committee for Clinical Laboratory Standards, Wayne, PA.
- Wegener, H.C., F. Bager and F.M. Aarestrup, 1997. Surveillance of antimicrobial resistance in humans, food stuffs and livestock in Denmark. *Euro Surveill.*, 2: 17-19.
- Wright, G.D., 2007. The antibiotic resistome: The nexus of chemical and genetic diversity. *Nat. Rev. Microbiol.*, 5: 175-186.