

Plasma Thyroid Hormone Concentrations and pH Values of Some GI-Tract Segments of Broilers Fed on Different Dietary Citric Acid and Microbial Phytase Levels

Rouhollah Nourmohammadi, Seyed Mohammad Hosseini, Hamed Saraee,
Alireza Arab and Hamed Arefinia
Department of Animal Science, College of Agriculture, University of Birjand,
P.O. Box 331/97175, Birjand, Iran

Abstract: An experiment was conducted to study the effect of microbial phytase supplementation and citric acid on thyroid activity, relative weight of lymphoid organs and pH values of some GI-tract segments in broiler chickens fed corn-soybean meal based diets. The data was analyzed using a Randomized Complete Block Design (RCBD) with factorial arrangement of 3×3 with three levels of citric acids (0, 3 and 6%) and three levels of phytase (0, 500 and 1000 IU kg⁻¹). There were three replicates (with ten chicks in each replicate) for each treatment. A total of 270 Ross 308 broilers chicks were used. The results indicated that addition of citric acid to diets caused significant decrease in pH values of crop, gizzard, duodenum, jejunum and ileum (p<0.05) and caused significant increase (p<0.01) in plasma Triiodothyronine (T₃) concentration, T₃:T₄ ratio and relative weight of bursa and thymus but had no significant effect on Thyroxin (T₄) concentrations. Microbial phytase significantly increased relative weight of thymus (p<0.01) but had no significant effects on thyroid gland activity, relative weight of bursa and values of pH in different parts of the GI-tract.

Key words: Broilers, acidify, phytase, pH value, thyroid hormones, Iran

INTRODUCTION

Phytate and phytate-bound Phosphorus (P) are present in all poultry diets and the partial availability of phytate-P has long been recognized (Lowe *et al.*, 1939). Possibly, Warden and Schaible (1962) were the first to show that exogenous phytase enhances phytate-P utilization and bone mineralization in broiler chickens. Nevertheless, three decades elapsed before an *Aspergillus niger* derived phytase feed enzyme with the capacity to liberate phytate bound P to reduce P excretion the was commercially introduced in 1991. Then, it was considered that the use of Microbial Phytases (MP) would be confined to areas where financial penalties were imposed on excessive P levels for excess from intensive pig and poultry units (Chesson, 1993). In contrast, the inclusion of phytase feed enzymes in monogastric diets has been far more widely accepted and now exceeds that of NSP degrading enzymes (Bedford, 2003). Phytase feed enzymes have more general application as their substrate is consistently present in pig and poultry diets and their dietary inclusion economically effect bioavailability of P and reduces the P load in the environment. Also, prohibition of animal origin

protein meals accelerate P acceptance as phytase feed enzymes in certain countries. Phytase is naturally found in a number of seeds including cereals, legume, by-products, other feedstuffs and microbial sources (Viveros *et al.*, 2000). Supplementation of diets with MP has known to increase availability of phytate P and Zn in chicks (Sebastian *et al.*, 1996; Ravindran *et al.*, 2000). Phytase increases availability and retention of Ca, improves absorption and retention of Mg, Cu and Fe (Sebastian *et al.*, 1996).

Previous researches have shown that poultry digestive tract acidity is not desirable to be completely hydrolyzed or established phytate by phytase (Brenes *et al.*, 2003). Citric Acid (CA) may change the intestinal pH and improve phytase enzyme activity because the phytase efficiency is correlated in both acidity and concentration of other free cations (Nourmohammadi *et al.*, 2010a). Therefore, it might be used with properties of chelated organic acids to enable intensification of phytase efficiency. Kemme *et al.* (1998) reported that phytase efficiency is related to plant or microbial sources in hydrolyzed phytate and gut pH term and durability time. Therefore, between CA and MP may have been synergistic effect. This study was carried out

to investigate the effect of supplementing diet with both MP and CA interaction on plasma hormones concentration, relative weight of lymphoid organs and pH values of some gastrointestinal tract segments.

MATERIALS AND METHODS

Diets, birds and experimental design: This experiment took place at Poultry Research Station and nutrition laboratory in University of Birjand, Iran. About 270 day old male chicks (Ross-308) were obtained from a commercial hatchery, weighed on arrival and randomly assigned to 27 pens of 10 birds each. The experiment was carried out using a Randomized Complete Block Design (RCBD) with factorial arrangement of 3×3, three levels of CA (0, 30 and 6%) and 3 levels of phytase (0, 500 and 1000 IU kg⁻¹ enzyme). There were 9 experimental diets, 3 replicates with 10 chicks in each replicate. Feed and water were provided *ad libitum* and a continuous lighting schedule were used throughout the experimental period. A basal diet (without phytase enzyme or CA) was formulated with corn-soybean meal from 7-21 and 22-42 day periods according to National Research Council recommendations. Diets were provided in the mash form. Broiler chicks were fed the following diets with equal energy and protein levels: T₁) basal diet, T₂) basal diet+500 IU kg⁻¹ MP, T₃) basal diet+1000 IU kg⁻¹ MP, T₄) basal diet+3% CA, T₅) basal diet+3% CA+500 MPIU kg⁻¹, T₆) basal diet+3% CA+1000 IU kg⁻¹ MP, T₇) basal diet+6% CA, T₈) basal diet+6% CA+500 IU kg⁻¹ MP, T₉) basal diet+6% CA+1000 IU kg⁻¹ MP. CA was supplied as monohydrate with 99.5% purity and phytase (Natuphos® 500, BASF Corp., Mt. Live, Nj) diet source also had 10,000 active phytase unit per gram.

Collection of samples and analysis: At the end of the experimental period (42 days of age), two birds per replicate (6 birds per treatment) were selected randomly and killed by cervical dislocation. Blood samples (approx. 10 mL) were collected in heparinized vacutainer tubes for measuring plasma hormones concentration (triiodothyronine and thyroxin). Immediately after collection, tubes were placed in an ice bath and transferred to the laboratory. Plasma was harvested subsequently after centrifuging the whole blood samples at 3000 rpm for 15 min. The heparinized plasma samples were stored at -20°C in Eppendorf tubes and analyzed subsequently. The triiodothyronine (T₃) and thyroxin (T₄) concentrations in the plasma samples were determined by Radioimmunoassay (RIA) using the procedure described by Darras *et al.* (1992).

At the end of the experimental period, two birds per replicate (6 birds per treatment) were taken randomly, weighed and killed by cervical dislocation then scalded

and defeathered. Thymus (all lobes of both sides) and bursa were removed and their relative percentages of live body weights were calculated.

pH values for different segments of the Gastrointestinal tract (GI-tract) were measured immediately by using a digital pH meter. About 10 g of contents from crop, gizzard, duodenum, jejunum and ileum were collected aseptically in 90 mL sterilized physiological saline (1:10 dilution) and their pHs were determined (Al-Natour and Alshawabkeh, 2005).

Statistical analysis: The data were subjected to an Analysis of Variance (ANOVA) through fitting General Linear Model (GLM) using SAS® (SAS, 2006) software and the corresponding means were compared by Tukey-Kramer test. The statistical model was as follows:

$$Y_{ijkl} = \mu + CA_i + MP_j + (CA \times MP)_{ij} + B_k + e_{ijkl}$$

Where:

- Y_{ijkl} = The individual observation
- μ = The experimental mean
- CA_i = The Citric Acid effect
- MP_j = The Microbial Phytase effect
- B_k = The block effect
- e_{ijkl} = The error term with mean 0 and variance σ²_e

RESULTS AND DISCUSSION

Results of the main effects of CA and MP on plasma hormones and relative weight of lymphoid organs are shown in Table 1. The results indicated that CA caused significant increase in T₃, T₃:T₄ ratio, bursa and thymus (p<0.01) but had no significant effect on plasma T₄ concentration. Also, MP caused an increase in relative weight of thymus (p<0.01) although, MP main effect was not significant for T₃, T₄, T₃:T₄ ratio and relative weight of bursa. Moreover, the present findings showed that there was significant effect between T₃, T₃:T₄ ratio and relative weight of bursa and thymus (p<0.05) (Table 2). It is well known that bursa and thymus are considered as parts of the immunity system (Sturkie, 1986) and this system is responsible for producing cells that protect the birds from the invaded microorganism. From Table 1, it is clearly observed that supplemental MP and CA, significantly increased the relative weight of both primary lymphoid organs (bursa and thymus). Increasing the weight of thymus may be due to the impact of MP on the functional activities of the immune system responses that led to increase in the number of lymphocytes in the primary lymphoid organs. These results may imply that broilers fed acidifiers diets obtained higher immune response and disease resistance. In this respect, Katanbaf *et al.* (1989)

Table 1: The main effects of CA and MP on plasma thyroid hormones and relative weight of lymphoid organs (g kg⁻¹ of live weight)

| Main effects | Thyroid gland activity | | | Lymphoid organs | |
|--------------------------------|---------------------------------------|---------------------------------------|--------------------------------|-------------------|-------------------|
| | T ₃ (ng mL ⁻¹) | T ₄ (ng mL ⁻¹) | T ₃ :T ₄ | Bursa | Thymus |
| MP (IU kg⁻¹) | | | | | |
| 0 | 1.702 | 16.133 | 0.105 | 0.27 | 0.33 ^b |
| 500 | 1.704 | 16.251 | 0.105 | 0.28 | 0.35 ^a |
| 1000 | 1.701 | 16.245 | 0.105 | 0.28 | 0.35 ^a |
| CA (%) | | | | | |
| 0 | 1.571 ^c | 16.234 | 0.097 ^c | 0.25 ^c | 0.28 ^c |
| 3 | 1.718 ^b | 16.141 | 0.106 ^b | 0.27 ^b | 0.34 ^b |
| 6 | 1.818 ^a | 16.255 | 0.112 ^a | 0.31 ^a | 0.41 ^a |
| SEM | 0.0020 | 0.0643 | 0.0005 | 0.003 | 0.005 |
| Probabilities | | | | | |
| MP | NS | NS | NS | NS | 0.01 |
| CA | 0.01 | NS | 0.01 | 0.01 | 0.01 |
| MP×CA | NS | NS | NS | NS | NS |

Mean values within a column with no common superscript differ significantly from each other (p<0.05). T₃ = Triiodothyronine, T₄ = Thyroxine, MP = Microbial Phytase, CA = Citric Acid, SEM = Standard Error of Mean, NS = Not Significant

Table 2: Interaction effects between CA and MP on plasma thyroid hormones and relative weight of lymphoid organs (g kg⁻¹ of live weight)

| Treatments | | Thyroid gland activity | | | Lymphoid organs | |
|------------|---------------------------|---------------------------------------|---------------------------------------|--------------------------------|--------------------|---------------------|
| CA (%) | MP (IU kg ⁻¹) | T ₃ (ng mL ⁻¹) | T ₄ (ng mL ⁻¹) | T ₃ :T ₄ | Bursa | Thymus |
| 0 | 0 | 1.5710 ^c | 16.228 | 0.0970 ^c | 0.240 ^b | 0.270 ^c |
| 0 | 500 | 1.5750 ^c | 16.247 | 0.0970 ^c | 0.260 ^b | 0.280 ^c |
| 0 | 1000 | 1.5690 ^c | 16.228 | 0.0970 ^c | 0.260 ^b | 0.290 ^{bc} |
| 3 | 0 | 1.7190 ^b | 15.918 | 0.1080 ^{ab} | 0.270 ^b | 0.330 ^d |
| 3 | 500 | 1.7180 ^b | 16.251 | 0.1060 ^b | 0.270 ^b | 0.350 ^{bc} |
| 3 | 1000 | 1.7170 ^b | 16.254 | 0.1060 ^b | 0.270 ^b | 0.350 ^{bc} |
| 6 | 0 | 1.8170 ^a | 16.254 | 0.1120 ^a | 0.310 ^a | 0.380 ^{ab} |
| 6 | 500 | 1.8210 ^a | 16.256 | 0.1120 ^a | 0.320 ^a | 0.420 ^a |
| 6 | 1000 | 1.8170 ^a | 16.255 | 0.1120 ^a | 0.300 ^a | 0.420 ^a |
| SEM | | 0.0034 | 0.1113 | 0.0005 | 0.006 | 0.009 |

Mean values within a column with no common superscript differ significantly from each other (p<0.05). T₃ = Triiodothyronine, T₄ = Thyroxine, MP = Microbial Phytase, CA = Citric Acid, SEM = Standard Error of Mean, NS = Not Significant

Table 3: The main effects of CA and MP on pH values of some gastrointestinal tract segments in broiler chicks

| Main effects | Crop | Gizzard | Duodenum | Jejunum | Ileum |
|--------------------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| MP (IU kg⁻¹) | | | | | |
| 0 | 5.000 | 3.160 | 5.770 | 6.450 | 7.200 |
| 500 | 5.020 | 3.160 | 5.770 | 6.450 | 7.190 |
| 1000 | 5.030 | 3.170 | 5.770 | 6.450 | 7.190 |
| CA (%) | | | | | |
| 0 | 5.170 ^a | 3.210 ^a | 5.800 ^a | 6.630 ^a | 7.220 ^a |
| 3 | 5.000 ^b | 3.190 ^a | 5.790 ^a | 6.490 ^b | 7.210 ^a |
| 6 | 4.890 ^c | 3.090 ^b | 5.710 ^b | 6.230 ^c | 7.160 ^b |
| SEM | 0.012 | 0.006 | 0.003 | 0.007 | 0.004 |
| Probabilities | | | | | |
| MP | NS | NS | NS | NS | NS |
| CA | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| MP×CA | NS | NS | NS | NS | NS |

Mean values within a column with no common superscript differ significantly from each other (p<0.05). SEM = Standard Error of Mean, MP = Microbial Phytase, CA = Citric Acid, NS = Not Significant

reported that increases in the relative lymphoid organs weights are considered as an indication of the immunological advances. The fact that thyroid hormones has a major role in regulating oxidative metabolism of birds has been established (Sturkie, 1986). Triiodothyronine (T₃), the metabolic activity of thyroid hormone, plays an active role in energy metabolism and metabolic rate. Any pronounced alteration in thyroid function (hyperthyroidism or hypothyroidism) is reflected

in an altered metabolic rate. The results pointed out superior metabolic and growth rate due to the addition of acidifiers into broiler chickens diet. The hyperthyroidism and peripheral conversion of T₄ to T₃ signified better. The concentration of thyroid hormones circulating in chicken blood plasma was found to be around 1.2 µL/100 mL (Davison, 1976) showing daily variations due to an extremely short half-life and showing T₃ to T₄ ratio to be 60:40, in favour of T₄. Current study data indicated that CA significantly decreased pH value of GI-tract segments (p<0.01) but MP had no significant effect on pH values (Table 3). Also, the results showed that there was significant effect between treatments for pH values of GI-tract parts (p<0.05) (Table 4). Similar results with the present study were found by Abdel-Fattah *et al.* (2008). In contrast, the other studies in which ascorbic acid and citric acid (Brown and Southern, 1985) were added to broiler diets, indicated that experimental treatments of intestinal content pH levels were not different compared to control that did not support the results. Decreasing pH in GI-tract had a beneficial effect in the inhibition of intestinal bacteria competition with the host for available nutrients and that possibility of reducing bacterial toxicity e.g., ammonia and amines thus improving weight gain of

Table 4: Interaction effect between CA and MP on pH values of some gastrointestinal tract segments in broiler chicks

| Treatments | | | | | | |
|------------|---------------------------|---------------------|--------------------|--------------------|--------------------|---------------------|
| CA (%) | MP (IU kg ⁻¹) | Crop | Gizzard | Duodenum | Jejunum | Ileum |
| 0 | 0 | 5.170 ^a | 3.210 ^a | 5.810 ^a | 6.640 ^a | 7.220 ^a |
| 0 | 500 | 5.170 ^a | 3.200 ^a | 5.800 ^a | 6.630 ^a | 7.220 ^a |
| 0 | 1000 | 5.170 ^a | 3.210 ^a | 5.800 ^a | 6.630 ^a | 7.220 ^a |
| 3 | 0 | 4.990 ^b | 3.200 ^a | 5.800 ^a | 6.480 ^b | 7.210 ^a |
| 3 | 500 | 5.000 ^b | 3.200 ^a | 5.790 ^a | 6.500 ^b | 7.200 ^{ab} |
| 3 | 1000 | 5.000 ^b | 3.180 ^a | 5.790 ^a | 6.490 ^b | 7.210 ^a |
| 6 | 0 | 4.850 ^c | 3.080 ^b | 5.720 ^b | 6.230 ^c | 7.170 ^{bc} |
| 6 | 500 | 4.900 ^{bc} | 3.090 ^b | 5.720 ^b | 6.230 ^c | 7.160 ^c |
| 6 | 1000 | 4.920 ^{bc} | 3.110 ^b | 5.710 ^b | 6.230 ^c | 7.150 ^c |
| SEM | | 0.022 | 0.010 | 0.004 | 0.012 | 0.007 |

Mean values within a column with no common superscript differ significantly from each other (p<0.05) SEM = Standard Error of Mean, MP = Microbial Phytase, CA = Citric Acid, NS = Not Significant

the host animals. Furthermore, the growth inhibition of potential pathogen bacteria e.g., *E. coli* and *Salmonella* in the feed and GI-tract is beneficial in respect to animal health (Thompson and Hinton, 1997). Organic acids are not antibiotics but if used correctly along with proper nutritional, managerial and biosecurity measures, they can be a powerful tools in maintaining the GI-tract state health of poultry thus improving their zootechnical performances. If applied correctly, organic acids work in poultry, not only as a growth promoter but also as a meaningful tool for controlling both pathogenic and non-pathogenic bacteria (Wolfenden *et al.*, 2007). Moreover, feeding organic acids are believed to have several beneficial effects such as improving feed conversion ratio, growth performance, enhancing mineral absorption and speeding recovery from fatigue (Nourmohammadi *et al.*, 2010b). The antibacterial activity of organic acids is related to the reduction of pH as well as their ability to dissociate that is determined by the pKa-value of the respective acid and pH of the surrounding environment. The antibacterial activity increases with decreasing pH value. Several investigations have shown a strong bactericidal effect of organic acid without significantly decreasing the pH value in the GI-tract. Generally lactic acid bacteria are able to grow at relatively low pH which means that they are more resistant to organic acids than other bacterial species, e.g., *E. coli* (Russell and Diez-Gonzalez, 1997). In poultry, pathogenic bacteria e.g., *Salmonella* enters the GI-tract via crop. The crop environment with respect to microbial composition and pH seems to be very important in relation to the resistance to pathogens.

CONCLUSION

In this study high amounts of Lactobacilli and low pH in the crop have shown to decrease the occurrence of *Salmonella* in the crop (Hinton *et al.*, 2000). Also the

antibacterial effect of dietary organic acids in chickens is believed to take place mainly in the upper part of the digestive tract (crop and gizzard). Therefore, following combination addition of formic and propionic acid (Bio-Add) in high concentrations could only be affected by crop and gizzard (Thompson and Hinton, 1997).

ACKNOWLEDGEMENTS

The researchers would like to acknowledge the contribution of Dan va Oloofeh Shargh Co. for the donation of feed and Broiler Breeder of South Khorasan Complex Productive Co. for the donation of day old chicks. The study was supported by University of Birjand.

REFERENCES

Abdel-Fattah, S.A., M.H. El-Sanhoury, N.M. El-Mednay and F. Abdel-Azeem, 2008. Thyroid activity, some blood constituents, organs morphology and performance of broiler chicks fed supplemental organic acids. *Int. J. Poult. Sci.*, 7: 215-222.

Al-Natour, M.Q. and K.M. Alshwabkeh, 2005. Using varying levels of formic acid to limit growth of *Salmonella gallinerum* in contaminated broiler feed. *Asian Aust. J. Anim. Sci.*, 18: 390-395.

Bedford, M.R., 2003. New enzyme technologies for poultry feeds. *Br. Poult. Sci.*, 44: 14-16.

Brenes, A., A. Viveros, I. Arija, C. Centeno, M. Pizarro and C. Braro, 2003. The effect of citric acid and microbial phytase on mineral utilization in broiler chicks. *Anim. Feed Sci. Technol.*, 110: 201-219.

Brown, D.R. and L.L. Southern, 1985. Effect of citric and ascorbic acids on performance and intestinal pH of chicks. *Poult. Sci.*, 64: 1399-1401.

Chesson, A., 1993. Feed enzymes. *Anim. Feed Sci. Technol.*, 45: 65-69.

Darras, V.M., T.J. Visser, L.R Berghman and E.R Kuhn, 1992. Ontogeny of type I and type III deiodinase activities in embryonic and posthatch chicks: Relationship with changes in plasma triiodothyronine and growth hormone levels. *Comp. Biochem. Physiol. Part A: Physiol.*, 103: 131-136.

Davison, T.F., 1976. Circulating thyroid hormones in the chicken before and after hatching. *General Comp. Endocrinol.*, 29: 21-27.

Hinton Jr, A., R.J. Buhr and K.D. Ingram, 2000. Reduction of *Salmonella* in the crop of broiler chickens subjected to feed withdrawal. *Poult. Sci.*, 79: 1566-1570.

- Katanbaf, M.N., E.A. Dunnington and P.B. Siegel, 1989. Restricted feeding in early and late-feathering chickens. 2. Reproductive responses. *Poult. Sci.*, 68: 352-358.
- Kemme, P.A., A.W. Jongbloed, Z. Mroz and A.C. Beynen, 1998. Diurnal variation in degradation of phytic acid by plant phytase in the pig stomach. *Livest. Prod. Sci.*, 54: 33-44.
- Lowe, J.T., H. Steenbock and C.H. Keiger, 1939. Cereals and rickets. IX. The availability of phytin-P to the chick. *Poult. Sci.*, 18: 40-44.
- Nourmohammadi, R., S.M. Hosseini and H. Farhangfar, 2010a. Influence of citric acid and microbial phytase on growth performance and carcass characteristics of broiler chickens. *Am. J. Anim. Vet. Sci.*, 5: 282-288.
- Nourmohammadi, R., S.M. Hosseini and H. Farhangfar, 2010b. Effect of dietary acidification on some blood parameters and weekly performance of broiler chickens. *J. Anim. Vet. Adv.*, 9: 3092-3097.
- Ravindran, V., S. Cabahug, G. Ravindran, P.H. Selle and W.L. Bryden, 2000. Response of broiler chickens to microbial phytase supplementation as influenced by dietary phytic acid and non-phytate phosphorus levels. II. Effects on apparent metabolizable energy, nutrient digestibility and nutrient retention. *Br. Poult. Sci.*, 41: 193-200.
- Russell, J.B. and F. Diez-Gonzalez, 1997. The effects of fermentation acids on bacterial growth. *Adv. Microb. Physiol.*, 39: 205-234.
- SAS, 2006. User's Guide. Release 9.1 Edn., SAS Institute Inc., Cary NC., USA.
- Sebastian, S., S.P. Touchburn, E.R. Chavez and P.C. Lague, 1996. The effects of supplemental microbial phytase on the performance and utilization of dietary calcium, phosphorus, copper and zinc in broiler chickens fed corn-soybean diets. *Poult. Sci.*, 75: 729-736.
- Sturkie, P.D., 1986. *Avian Physiology*. 4th Edn., Springer-Verlag, New York.
- Thompson, J.L. and M. Hinton, 1997. Antibacterial activity of formic and propionic acids in the diet of hens on *Salmonella* in the crop. *Br. Poult. Sci.*, 38: 59-65.
- Viveros, A., C. Centeno, A. Brenes, R. Canales and A. Lozano, 2000. Phytase and acid phosphatase activities in plant feedstuffs. *J. Agric. Food Chem.*, 48: 4009-4013.
- Warden, W.K. and P.J. Schaible, 1962. Preliminary investigations concerning utilization of phytin phosphorus by the chick. *Poult. Sci.*, 41: 1692-1692.
- Wolfenden, A.D., J.L. Vicente, J.P. Higgins, R.L. Andreatti-Filho, S.E. Higgins, B.M. Hargis and G. Tellez, 2007. Effect of organic acids and probiotics on *Salmonella enteritidis* infection in broiler chickens. *Int. J. Poult. Sci.*, 6: 403-405.