

RAPD-PCR Analysis of Water Vole, *Arvicola amphibius* (Linnaeus, 1758) (Mammalia: Rodentia) Distributed in Turkey

Sevilay Ustunbas, Reyhan Colak, Gul Olgun Karacan and Ercument Colak
Department of Biology, Faculty of Science, Ankara University, 06100 Ankara, Turkey

Abstract: Water vole, *Arvicola amphibius* is a rodent distributed widely in Palearctic region. Three subspecies of *A. amphibius* are distributed in Turkey. *A. amphibius* lives in Turkey located between Europe and Asia. In Turkey, there is no any study on this species at the level of genetic structure. The aim of the present study was to survey genetic structure based on DNA markers and to contribute to the taxonomic status, population genetics of *A. amphibius*, distributed in Thrace and Anatolia. A total of 38 specimens were collected from nine locations. In order to explore the extent of genetic variation in *A. amphibius* populations, a Randomly Amplified Polymorphic (RAPD) DNA marker system was used. The estimates of Nei's standard genetic identity and standard genetic distance were calculated to show the genetic relationships between populations studied. UPGMA dendrogram constructed with genetic distance data was clustered in 2 groups. The 1st group contains Thrace populations and the 2nd one including Anatolian populations was divided into 3 subgroups. Consequently, RAPD-PCR marker system confirmed the validity of *A.a. cernjavskii* and *A.a. persicus*.

Key words: Water vole, *Arvicola amphibius*, evolution, RAPD-PCR, populations, Turkey

INTRODUCTION

Water voles were once a familiar waterside animal often known locally as a water rat. Water voles are almost wholly vegetarian, feeding on a wide range of plants. They need luxurious bank side vegetation, particularly grasses and sedges to provide food and cover from predators. Although, water voles are widely distributed in palearctic region, they are one of the most rapid and serious declines of any mammal in recent years. This decline is attributed to habitat loss such as through river management and drainage. On this account beside morphological (Kratochvil, 1983; Nikolaeva, 1982; Krystufek and Tvrtkovic, 1984; Ventura, 1991) and karyological studies (Raicu *et al.*, 1971; Kuliev *et al.*, 1978; Zima and Kral, 1984) there are several researches on metapopulation level (Stewart *et al.*, 1999; Berthier *et al.*, 2004, 2005, 2006; Aars *et al.*, 2006).

Also mitochondrial genes were used for phylogeny researches (Martin *et al.*, 2000; Pfunder *et al.*, 2004; Piertney *et al.*, 2005). Mursaloglu reported three subspecies of *Arvicola amphibius* in Turkey: *A. amphibius hintoni* from South Eastern Turkey, *A. amphibius persicus* from Anatolia and *A. amphibius cernjavskii* from Turkish Thrace. Morphological, karyological and biometric characters of *A. amphibius* were studied in Turkey (Ozkurt *et al.*, 1999;

Gozcelioglu *et al.*, 2006). Although, blood proteins and allozyme profiles were investigated, these studies contained limited areas (Iyigun and Colak, 2004). In recent years to reveal intraspecific genetic differentiation and define the origin of the species, mtDNA and microsatellites have been usually used in Europe. But in Turkey, this species has not been studied on genetic level and this cause deficiency of data while considering *A. amphibius* population with other geographic forms in literatures. In this study, RAPD markers were elected for why this marker can provide an efficient assay for polymorphisms which should allow rapid identification and isolation of chromosome-specific DNA fragments (Williams *et al.*, 1990).

MATERIALS AND METHODS

Sampling localities: We examined 38 individuals of *Arvicola amphibius* from nine localities of Turkey. As outgroups, we used Macedonian mouse *Mus macedonicus* from Kirklareli and Konya (N = 4), *Microtus levis* from Konya (N = 2) and *Microtus guentheri* from Kirklareli (N = 2). The sampling localities and sizes were as follows: Kirklareli (N = 6), Denizli (N = 4), Usak (N = 4), Afyon (Lake Eber) (N = 4), Konya (Lake Beysehir) (N = 4), Eskisehir (N = 4), Bolu (N = 4), Ankara (N = 4), Kirsehir (N = 4) (Fig. 1).



Fig. 1: Collecting localities of specimens (Numbers refer to localities were presented in materials and methods; ●: *Arvicola*, ■: *Microtus*, ▲: *Mus*)

Table 1: Sequences and polymorphism percentage of all primers

Primers	Sequences	P (%)
OPA-02	5'-TGC CGA GCT G-3'	4.1
OPA-03	5'-AGT CAG CCA C-3'	5.4
OPA-04	5'-AAT CGG GCT G-3'	9.5
OPA-07	5'-GAA ACG GGT G-3'	8.2
OPA-08	5'-GTG ACG TAG G-3'	7.5
OPA-16	5'-AGC CAG CGA A-3'	7.5
OPB-15	5'-GGA GGG TGT T-3'	5.4
OPB-16	5'-TTT GCC CGG A-3'	5.4
OPB-18	5'-CCA CAG CAG T-3'	4.1
OPB-19	5'-ACC CCC GAA G-3'	5.4
OPB-20	5'-GGA CCC TTA C-3'	4.1
OPD-08	5'-GTG TGC CCC A-3'	4.1
OPD-09	5'-CTC TGG AGA C-3'	5.4
OPD-10	5'-GGT CTA CAC C-3'	6.1
OPD-11	5'-AGC GCC ATT G-3'	5.4
OPD-12	5'-CAC CGT ATC C-3'	6.1
OPD-14	5'-CTT CCC CAA G-3'	5.4

Isolation procedure and amplification conditions: DNA was isolated from kidney tissue according to the CTAB method of Doyle and Doyle (1990). DNA was quantified using a spectrophotometer (Agilent, 2100 Bioanalyser NanoDrop ND-1000 spectrophotometer).

The PCR was run in 25 μ L of a reaction mixture containing 1 μ L of the DNA samples (200 ng μ L⁻¹); 2.5 μ L of buffer (750 mM Tris-HCl pH: 8.8, 200 mM (NH₄)₂ SO₄; fermentas); 0.3 μ L of Taq DNA polymerase (100 unit fermentas); 4 μ L of deoxynucleotide triphosphate mix (200 μ M of each nucleotide); 1.5 μ L of 2 mM MgCl₂; 1 μ L of 1 pmol primers (Thermo electron). The PCR steps were as follows: 95°C for 1 min, 45 cycles of 94°C for 1 min, 36°C for 2 min, 72°C for 2 and 15 min. Pre-screening of 23 random decamer primers revealed that 17 primers could be useful for further study and data collection (Table 1).

Agarose gel electrophoresis: The amplification products were separated on 1.7% agarose gels in 1X TAE (Tris, acetic acid, EDTA) buffer at 100 V for 4 h and visualized

by staining with ethidium bromide. A 100 base pair ladder was used as a size standard marker (DNA ladder plus, fermentas).

Analyzing of amplified products: All visible bands on gels were considered as RAPD loci and all loci were scored as presence (1) and absence (0) of the bands. We used POPGENE Version 1.31 (Yeh *et al.*, 1997) software package to compute the intrapopulation and interpopulation variations. By this software, the percentage of polymorphic loci (P), observed number of alleles (N_a), effective number of alleles (Kimura and Crow 1964) (N_e) and Nei (1972)'s gene diversity (H) and Shannon's Information index (Lewontin, 1972) (I) were computed to display intrapopulation variations. The estimated parameters of interpopulation differentiation included total gene diversity (H_T), intrasample gene diversity (H_S), interpopulation gene diversity (D_{ST}), coefficient of gene fixation (G_{ST}) and coefficient of gene flow (the number of migrants per generation) Nm. Genetic distance matrix (Nei, 1972) was used to draw UPGMA tree by TFPGA software Version 1.3 (Miller, 1997) and MEGA software Version 4.0 (Tamura *et al.*, 2007).

RESULTS

In this study, 17 of 23 RAPD primers were choice to analyze *Arvicola* specimens. While these 17 RAPD primers constituted 147 bands for all individuals, only 95 bands were observed in *Arvicola* specimens. Four primers (OPA-2₆₅₀, OPB-16₇₀₀, OPB-19₂₀₀₀ and OPD-10₅₅₀) were diagnostic between Anatolian and Thrace populations (Fig. 2).

Inference of genetic variation and differentiation analysis: Genetic distance matrix that was computed according to Nei (1972) showed that while the closest populations were Eskisehir and Bolu (D = 0.057), the most distant populations were Kirklareli ve Bolu (D = 0.187) (Table 2).

The mean observed Number of alleles (N_a) was 1.561. When all populations were considered, the mean Ne value was 1.266. Nei's genetic diversity or Heterozygosity (H) was the lowest in Denizli (0.051) and the highest in Afyon (0.094). For all populations, the genetic diversity was calculated as 0.160.

The high G_{ST} value of 0.496 indicated that genetic differentiation among the studied populations was substantial. The total gene diversity (H_T) was 0.146 in *Arvicola amphibius* populations but 50.3% of this was within population variation (H_S = 0.0736). UPGMA tree was constructed using TFPGA and MEGA software




Fig. 2: Amplification products of Primer OPA-02. The arrows show diagnostic RAPD loci in Kirklareli specimens. M: marker (100 bp DNA ladder, fermentas), C: negative control. 1-4: Kirsehir, 5-8: Ankara, 9-14: Kirklareli, 15-18: Konya, 19-22: Afyon, 23-26: Denizli, 27-30: Usak, 31-34: Eskisehir, 35-38: Bolu (1-38: *A. amphibius*), 39-40: Kirklareli (*Microtus guentheri*), 41-42: Konya (*Microtus levis*), 43-44: Konya (*Mus macedonicus*), 45-46: Kirklareli (*Mus macedonicus*)




Fig. 3: Genetic similarity of *A. amphibius* from different localities based on RAPD data and generated using the UPGMA method, respectively within the MEGA software program. At the branching nodes, the BP values exceeding (50% are presented)

Table 2: Pairwise dissimilarity matrix based on Nei (1972) between *Arvicola amphibius* populations

	1	2	3	4	5	6	7	8	9
Kirsehir	-								
Ankara	0.0681	-							
Kirklareli	0.1338	0.1683	-						
Konya	0.1070	0.1084	0.1488	-					
Afyon	0.0847	0.1067	0.1389	0.0885	-				
Denizli	0.1122	0.1295	0.1626	0.0906	0.1042	-			
Usak	0.0623	0.0785	0.1670	0.0812	0.0969	0.1033	-		
Eskisehir	0.0776	0.0794	0.1516	0.0799	0.0773	0.0992	0.0665	-	
Bolu	0.0957	0.0744	0.1879	0.1301	0.1126	0.1141	0.0959	0.0571	-

packages. All specimens analyzed were divided into two major groups while the 1st group contained only Kirklareli

populations called Thrace, the 2nd group contained three sub-groups as Kirsehir-Usak-Ankara-Eskisehir-Bolu, Konya-Afyon and Denizli called Anatolian group (Fig. 3).

DISCUSSION

Geographical variations in subspecific level of *A. amphibius* were investigated in Turkey. Ozkurt *et al.* (1999) and Gozcelioglu *et al.* (2006) separated *A.a. cernjavskii* in Thrace from *A.a. persicus* in Anatolia based on karyotype analysis. In this study, RAPD analyses of Anatolian and Thrace populations supported the existence of these two subspecies. In addition Iyigun and Colak (2004) proved high genetic diversity in Kirsehir populations of Turkey due to absence of the bottleneck based on their esterase and SDS-PAGE studies. Contrarily RAPD loci did not show obvious disparity in heterozigosity of Kirsehir populations, possibly owing to the difference of the markers between two studies.

Afyon population and Konya population were formed from Lake Eber and Lake Beysehir, respectively. These two populations clustered together in UPGMA dendrogram in consequence of the lakes have very similar water vole habitats. Similarly, populations from rivers and brooks were clustered in same group (Kirsehir, Usak, Ankara, Eskisehir, Bolu). This habitat similarity reduces the differentiation between populations. Although, Denizli specimens were collected from riverside too, genetic drift and fluctuation in population density may cause evolution of the populations differently therefore, Denizli population might differ from Anatolian group in this way. According to the RAPD data while there is gene flow between *A. amphibius* populations in Anatolia, presence of subpopulations might be depended on river and lake habitats. In additional, Bosphorus and Marmara sea seem to interrupt gene flow between Thrace and Anatolia populations. This barrier effect may cause differentiation of *A.a. cernjavskii* and *A.a. persicus*.

RAPD-PCR is widely used in rodents (Atopkin *et al.*, 2007; Spiridonova *et al.*, 2008; Dokuchaev *et al.*, 2008; Olgun *et al.*, 2009). RAPD assay may in some instances detect single base changes in genomic DNA. Most single nucleotide changes in a primer sequence caused a complete change in the pattern of amplified DNA segments (Williams *et al.*, 1990).

CONCLUSION

This study was the first molecular study of *Arvicola* in Turkey on DNA level. As a result, there is a significant genetic differentiation between Thrace and Anatolian

populations. In order to discover evolution and population dynamics of *A. amphibius* much more molecular technique should be used to reach definitive conclusion in DNA level.

ACKNOWLEDGEMENT

Specimens used in this study were collected between 2008-2010 funded by the Scientific Research Office of Ankara University (BAPRO No: 2008-07-05-003).

REFERENCES

- Aars, J., J.F. Dallas, S.B. Piertney, F. Marshall, J.L. Gow, S. Telfer and X. Lambin, 2006. Widespread gene flow and high genetic variability in populations of water voles *Arvicola terrestris* in patchy habitats. *Mol. Ecol.*, 15: 1455-1466.
- Atopkin, D.M., A.S. Bogdanov and G.N. Chelomina, 2007. Genetic variation and differentiation in striped field mouse *Apodemus agrarius* inferred from RAPD-PCR analysis. *Genetika*, 43: 804-817.
- Berthier, K., M. Galan, A. Weber, A. Loiseau and J.F. Cosson, 2004. A multiplex panel of dinucleotide microsatellite markers for the water vole, *Arvicola terrestris*. *Mol. Ecol. Notes*, 4: 620-622.
- Berthier, K., M. Galan, J.C. Foltete, N. Charbonnel and J.F. Cosson, 2005. Genetic structure of the cyclic fossorial water vole (*Arvicola terrestris*): Landscape and demographic influences. *Mol. Ecol.*, 14: 2861-2871.
- Berthier, K., N. Charbonnel, M. Galan, Y. Chaval and J.F. Cosson, 2006. Migration and recovery of the genetic diversity during the increasing density phase in cyclic vole populations. *Mol. Ecol.*, 15: 2665-2676.
- Dokuchaev, N.E., A.G. Lapinskii and L.L. Solovenchuk, 2008. Genetic diversity of the striped field mouse (*Apodemus agrarius* Pallas, 1771) in the Russian Far East as assessed by RAPD-PCR. *Biol. Bull.*, 35: 368-373.
- Doyle, J.J. and J.L. Doyle, 1990. Isolation of plant DNA from fresh tissue. *Focus*, 12: 13-15.
- Gozeioglu, B., E. Colak and R. Colak, 2006. Karyotype of *Arvicola terrestris* (Mammalia: Rodentia) in Turkish Thrace. *Pak. J. Biol. Sci.*, 9: 2387-2388.
- Iyigun, C. and R. Colak, 2004. An electrophoretic study on esterase and blood serum proteins of the water vole *Arvicola terrestris* (L., 1758) (Mammalia: Rodentia), in Kirsehir province. *Turk. J. Biol.*, 28: 47-53.
- Kimura, M. and J.F. Crow, 1964. The number of alleles that can be maintained in a finite population. *Genetics*, 49: 725-738.
- Kratochvil, J., 1983. Variability of some criteria in *Arvicola terrestris* (Arvicolidae, Rodentia). The effect of altitude on some taxonomical criteria of the Asian population group of *Arvicola terrestris*. *Acta Scientiarum Bohemoslovaca* (Brno), 17: 1-40.
- Krystufek, B. and N. Tvrtkovic, 1984. Redescription of *Arvicola terrestris illyrica* (Barret-Hamilton, 1899)-Rodentia, Mammalia. *Biosystematika*, 10: 91-97.
- Kuliev, G.N., G.K. Kuliev and S.I. Radjabli, 1978. Karyotypical differences between different populations of *Arvicola terrestris* Rodentia Cricetidae. *Zool. Z.*, 57: 1409-1411.
- Lewontin, R.C., 1972. The Apportionment of Human Diversity. In: *Evolutionary Biology*, Dobzhansky, T., M.K. Hecht and W.C. Steere (Eds.). Appleton Century Crofts, New York, pp: 381-398.
- Martin, Y., G. Gerlach, C. Schlotterer and A. Meyer, 2000. Molecular phylogeny of European muroid rodents based on complete cytochrome b sequences. *Mol. Phylogenet. Evol.*, 16: 37-47.
- Miller, M.P., 1997. Tools for Population Genetic Analysis (TFPGA). Version 1.3, A Windows Program for the Analysis of Allozyme and Molecular Population Genetic Data. <http://www.ccg.unam.mx/~vinuesa/tlem09/docs/TFPGADOC.PDF>.
- Nei, M., 1972. Genetic distance between populations. *Am. Naturalist*, 106: 283-292.
- Nikolaeva, A.I., 1982. Adaptive variation in the masticatory surface of molar teeth in *Arvicola terrestris*. *Zoologicheskii Zhurnal*, 61: 1565-1575.
- Olgun, G., R. Colak, I. Kandemir, E. Colak and N. Yigit, 2009. Genetic variation in rocky mouse, *Apodemus mystacinus* (Danford and Alston 1877) (Mammalia: Rodentia) in Turkey. *Acta. Zool. Bulg.*, 61: 123-129.
- Ozkurt, S., E. Colak, N. Yigit, M. Sozen and R. Verimli, 1999. Contributions to the karyology and morphology of *Arvicola terrestris* (Lin., 1758) (Mammalia: Rodentia) in Central Anatolia. *Turk. J. Zool.*, 23: 253-257.
- Piertney, S.B., W.A. Stewart, X. Lambin, S. Telfer, J. Aars and J.F. Dallas, 2005. Phylogeographic structure and postglacial evolutionary history of water voles (*Arvicola terrestris*) in the United Kingdom. *Mol. Ecol.*, 14: 1435-1444.
- Pfunder, M., O. Holzgang and J.E. Frey, 2004. Development of microarray-based diagnostics of voles and shrews for use in biodiversity monitoring studies, and evaluation of mitochondrial cytochrome oxidase I vs. cytochrome b as genetic markers. *Mol. Ecol.*, 13: 1277-1286.
- Raicu, P., D. Duma, M. Kirillova and A. Tuta, 1971. Chromosomal polymorphism in the water vole (*Arvicola terrestris* L.). *Rev. Roum. Biol. Zool.*, 16: 293-296.

- Spiridonova, L.N., K.V. Korobitsyna, L.V. Yakimenko and A.S. Bogdanov, 2008. Genetic differentiation of subspecies of the house mouse *Mus musculus* and their taxonomic relationships inferred from RAPD-PCR data. *Russian J. Genet.*, 44: 732-739.
- Stewart, W.A., J.F. Dallas, S.B. Piertney, F. Marshall, X. Lambin and S. Telfer, 1999. Metapopulation genetic structure in the water vole, *Arvicola terrestris*, in NE Scotland. *Biol. J. Linn. Soc.*, 68: 159-171.
- Tamura, K., J. Dudley, M. Nei and S. Kumar, 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24: 1596-1599.
- Ventura, J., 1991. Morphological characteristics of the molars of *Arvicola terrestris* (Rodentia, Arvicolidae) in its southwestern distribution area. *Zoologischer Anzeiger*, 226: 64-70.
- Williams, J.G.K., A.R. Kubelik, K.J. Livak, J.A. Rafalski and S.V. Tingey, 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acids Res.*, 18: 6531-6535.
- Yeh C.F., R. Yang and T. Boyle, 1997. POPGENE Version 1.31. Windows-Bsed Software for Population Genetics Analysis.
- Zima, J. and B. Kral, 1984. Karyotypes of european mammals II. *Acta Sci. Nat. Brno.*, 18: 1-62.