

The Identification of Cherry Valley Ducks CD8+Lymphocytes by Using *Homo sapiens* CD8 Antibodies

^{1,2}J.Y. Bai, ^{1,2}W. Li, ¹S. Li, ¹W.P. Zhang, ¹C.Q. Gu, ¹X.Y. Hu and ¹G.F. Cheng

¹College of Veterinary Medicine, Huazhong Agricultural University, Lion Mountain Street No. 1,
South Lake District, Wuhan, 430070 Hubei Province, People's Republic of China

²Xinyang Agricultural College, Xinyang Agricultural College Road No. 1, Xinyang,
464000 Henan Province, People's Republic of China

Abstract: To study the cross-application of the Antibodies (Abs) of *Homo sapiens* and duck origins which provides the fundamentals for potential application of related Abs, we analyzed their differences by molecular biology techniques and preliminary application in peripheral blood. The similarities of deduced aa sequences of *CD8 α* gene reached to about 32% between cherry valley duck and homo sapiens whereas the extracellular regions reached to about 30%. The result showed the obvious differences in protein hydrophilicity, antigenicity and the possibilities of *CD8 α* extracellular region. This study successfully expressed the extracellular region of cherry valley ducks *CD8 α* (ERCVC $CD8$) sequence and prepared for corresponding Abs. The percentage of control group (group 1) *CD8*+lymphocytes was relatively stable during different phases of the post-infected with DPV when using rabbit anti-the recombinant protein of extracellular region of cherry valley ducks *CD8 α* (rCVER $CD8$) serum but the percentage of *CD8*+lymphocytes was unstable when using anti-*Homo sapiens* *CD8 α* mAb. When rabbit anti-rCVER $CD8$ serum was used, the percentage of infection group (group 2) *CD8*+lymphocytes showed regularity however that of *CD8*+lymphocytes using rabbit anti-*Homo sapiens* *CD8 α* Abs was irregular. This study indicates that the identification of cherry valley ducks *CD8*+lymphocytes using anti-*Homo sapiens* *CD8 α* Abs has its limitations.

Key words: Cherry valley duck, *CD8 α* , *Homo sapiens*, blood, region, China

INTRODUCTION

The *CD8* molecule is transmembrane glycoprotein of T lymphocytes surface (Gillooly *et al.*, 2001) and simultaneously it also mainly exists in Cytotoxic T cell (CTL), suppression T cell and NK cell's surface (Fu *et al.*, 2007). *CD8 α* linked Major Histocompatibility Complex (MHC) molecule immunoglobulin superfamily (Ig-SF) is helpful for T Cell Receptor (TCR) to identify Ag-MHC complex. Both intracellular region p56lck of *CD8 α* chain and tyrosine kinase have the adjustment function of *CD3/TCR* complex. *CD8 α* chain has played an important role in transducing signal of T cell proliferation and differentiation (Konno *et al.*, 2002; David and George, 2004). *CD8 α* chain carboxyl terminal does not contain p56lck region currently, there is an idea that *CD8 α* chain is only an assistant action during biologic activity of the *CD8 α* chain and *CD8* molecule is constructed with homodimer (*CD8 $\alpha\alpha$*) or heterodimer (*CD8 $\alpha\beta$*) (Luhtala, 1998). The *CD8*+CTL are essential to protect against viruses, intracellular bacteria infection and tumor cells (Zhang *et al.*, 2009) and it is an integral component of

cellular immunity and factors that influenced *CD8*+lymphocytes cell function which may be critical to an effective immune response in HIV infection (Karen and Copeland, 2001). This *CD8*+Cell Noncytotoxic Anti-HIV Response (CNAR) was mediated by a soluble *CD8*+Cell Antiviral Factor (CAF). CNAR/CAF inhibited HIV-1 replication by blocking viral RNA transcription (Bonneau *et al.*, 2008). The immunity of avian fell far behind that of mammals (Tregaskes *et al.*, 1995). As the chicken cellular immune response related to molecules MHC⁻ (BF), β 2m, TCR, the *CD8 α/β* genes were cloned (Luhtala *et al.*, 1997). Firstly, the reports concerning the duck Ancy-MHC⁻, β 2m cDNA and genomic structure (Xia *et al.*, 2004) at present, the ratio of *CD4*+ and *CD8*+lymphocytes in peripheral blood has been used to evaluate the immune effect of virus vaccine in many countries (Kothlow *et al.*, 2005). In China, a number of studies on the hypothesis found that duck T cells shared a common antigen with human *CD3* epsilon chain (Higgins and Chung, 1986; Bertram *et al.*, 1996; Shawky *et al.*, 2000). Previously it was hard to see the published demonstrations about *CD8 α* . In this study,

Corresponding Author: G.F. Cheng, College of Veterinary Medicine, Huazhong Agricultural University,
Lion Mountain Street No. 1, South Lake District, Wuhan, 430070 Hubei Province,
People's Republic of China

through the analyses of genes coding of CD8 α and the use of different Abs on immunohistochemistry (SABC), the differences between cherry valley ducks CD8 α and *Homo sapiens* CD8 α Abs were comparatively analyzed and during the analytic phase, we observed the changes of CD8+lymphocytes. Therefore, this study will provide us a fundamental basis for potential application of related Abs that can be used as a research tool to observe the changes of the body's cellular immune function in the immunosuppressive diseases study.

MATERIALS AND METHODS

Annotation of CD8 α genes: The *M. domestica* whole genome sequence data released from National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov/>). The MonDom version 2.0 (MonDom2.0) ensemble assembly was analyzed by Blast search (using default search parameters) using certain Chinese domestic duck breeds CD8 α cDNA sequences (GenBank accession Nos.NM_001768, AY519197, AF378373, FJ527828 and FJ527912) as query.

Animal tissues and cDNA preparation: Cherry valley duck spleen tissues from a 30 days old female animal were supplied by the Chinese Academy of Agricultural Sciences, WuHan, China. Total RNA was isolated from Cherry valley duck spleen tissue using TR Trizol reagent (Sigma Company, USA) following the manufacturer's instructions. For RT-PCR, cDNA was synthesized using SuperScript 2 reverse transcription system (TaKaRa, DaLian, China) according to the manufacturer's protocol. The cDNA was stored at -20°C until use.

Sequence data analysis: The Cherry valley duck CD8 α Open-Reading Frame (ORF) genes were aligned with that

of other known avian and *Homo sapiens* using ClustalW(BioManager) and the BLAST program at the NCBI database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) using the default parameters. Putative signal peptide cleavage sites for the amino acid (aa) sequences were predicted using the SignalP program (www.cbs.dtu.dk/services/signalP) and transmembrane domain were predicted using interproscan (www.ebi.ac.uk/interproscan). Analyses of hydrophilicity plot, antigenicity index and surface probability plot of CD8 α extracellular region deduced aa sequences were performed by DNASTar software.

Construction of bacterial expression vector: The extracellular region of cherry valley ducks CD8 α (CVERCD8) sequence was amplified by PCR from the Cherry valley duck cDNA using synthetic oligonucleotide (F: 5-CGC GAA TTC AGA AGA TCA CAG TGA CGG CCA AGT T-3) as the forward primer and synthetic oligonucleotide (R: 5-CGC CTC GAG TCA ACA GAA GAA ATT CAG CTC TTT C-3') as the reverse primer. EcoRI and XhoI sites were incorporated respectively into the forward and reverse primers. PCR reactions that were performed with the BD advantage polymerase mix using a touchdown program were as follows: initial denaturation at 94°C for 2 min followed by 35 cycles of denaturation at 94°C for 30 sec; an annealing temperature commencing at 63°C and extension at 72°C for 30 sec. A final elongation step was carried out at 72°C for 5 min. The PCR products were purified by DNA Fragment Purification Kit (TOYOBO, Japan) and cloned into the pMD18-T vector (TaKaRa, Dalian, China). The CVERCD8 sequence was subsequently released by EcoRI/XhoI digestion and cloned into the EcoRI and XhoI sites of PEGX-KG vector (Novagen) in frame with the gene encoding GST (Fig. 1a).




Fig. 1: a) Shows schematic representation of extracellular region of Cherry Valley duck CD8 α and strategy for construction of the expression plasmid (PEGX-KG-CVERCD8); b) shows PCR product of CVERCD8 detected by 1% agarose gel electrophoresis. Lane M represents DNA marker (DL2000). Lane 1 represents PCR product of CVERCD8; c) shows double digestion analysis of recombinant plasmid (PEGX-KG-CVERCD8) detected by 1% agarose gel electrophoresis. Lane M1 represents DNA Marker (DL2000). Lane 1 represents results of double digestion of recombinant plasmid (PEGX-KG-CVERCD8). Lane M2 represents DNA Marker (DL15000)

The recombinant plasmid (PEGX-KG-CVERCD8) was confirmed by colony PCR, restriction enzyme digestion and sequencing (TaKaRa).

Expression and purification of GST-rCVERCD8 protein:

The confirmed construction described above was used to chemically transform *Escherichia coli* BL21 (DE3) for expression of the CVERCD8 protein. For production of GST-CVERCD8 protein, 100 μ L of fresh stationary-phase culture was inoculated into 10 mL of Luria Broth (LB) supplemented with 50 μ g mL⁻¹ ampicillin (Sigma). To optimize expression, the bacterial culture was bred at 37°C until the optical density at 595 nm was 0.5 and during the process, the protein expression was induced by the addition of 0.8 mM Isopropyl-d-Thiogalactopyranoside (IPTG). The culture was shaken at 210 rpm at 37°C for 4 h in a 100 mL Erlenmeyer flask. After induction, cells were lysed in 2×sample buffer (0.1 M Tris-HCl, pH 6.8, 4% SDS, 0.2% bromophenol blue, 20% glycerol and 0.1 M DTT) and analyzed by SDS-PAGE. The recombinant GST-tagged proteins were purified by nickel affinity chromatography according to the manufacturer's protocol (TaKaRa) and analyzed by SDS-PAGE.

Western blotting: The recombinant protein of extracellular region of cherry valley ducks CD8 α (rCVERCD8) was analyzed by Western blotting analysis with mAb of anti-GST (Boster WuHan, China). *E. coli* expressed GST-tagged rCVERCD8 were harvested by centrifugation. Western blotting technique was described. Specific bands were detected using an Enhanced Chemiluminescence (ECL system) according to the manufacturer's instructions (Amersham). The other reagent was offered by WuHan Boster Company in china.

Generation of polyclonal anti-rCVERCD8 in rabbits: For the preparation of polyclonal Abs, male rabbits were immunized first with 0.5 mg of *E. coli* expressed GST-tagged CVERCD8 emulsified in Freund's complete adjuvant. Inoculations were subcutaneous injections on the shaven back. Freund's incomplete adjuvant and 1mg of purified fusion protein were used for subsequent boosts. Three booster injections were given each at 1 week interval after primary injection. About 18 days after the last boost, blood was collected from an ear vessel. Then sera were collected and stored at -80°C. This experiment employed saturated ammonium sulfate and affinity chromatography (Sephadex gel) to purify serum and the methods of anti-rCVERCD8 serum purification were in accordance with the manufacturer's instructions (boster, WuHan, China). The bandscan 5.0 software were utilized to measure Abs purity. Measuring for titer of anti-rCVERCD8 polyclonal serum using ELISA was performed as Sonja K described (Kothlow *et al.*, 2005).

Preliminary application of antibody: Twelve 35 days old nonimmune ducklings were divided randomly to two groups control group (group 1) and infection group (group 2). The six ducklings of group 1 were intramuscularly inoculated with allantoic liquid containing DPV isolates (China Agricultural University presented). Ducklings in group 2 were inoculated with 0.9% NaCl as negative control. The six living ducklings peripheral blood of group 1 was collected on everyday during 6 days of post-infection and then smeared to be well-distributed on the slide at once. All blood smears were stained with anti-homo sapiens CD8 α mAb (Boster, WuHan, China) and rabbit anti-CVERCD8 serum using an indirect immunoperoxidase technique as described (Grodioa *et al.*, 2009). At second-stage reagent, Vectastain¹ SABC kit (Boster, WuHan, China) was used. Ducklings peripheral blood of group 2 was executed the same test procedure.

Statistical analysis: Test of percentage of CD8 positive lymphocytes concerning the all lymphocytes by Nikon image analysis software which computed means of group 1 on everyday during 6 days of post-infection by different Ads for statistical analysis. All the means of experimental treatments were analyzed by ANOVA with the GLM procedure of SAS. Ducklings peripheral blood of group 2 which collected on everyday during 6 days of post-infection executed the same analysis procedure.

RESULTS AND DISCUSSION

Characterization of cherry valley duck CD8 α ORF gene:

In this study, the Cherry valley duck CD8 α Open Reading Frame (ORF) sequence (GenBank accession no.FJ527828) comprised 714 nucleotides (Fig. 2). The ORF translated into mature protein of 227 amino acid residues (Fig. 3). The mature protein included four domains, signal peptide (1-23AA), extracellular (24-182AA), transmembrane (183-205AA) and cytoplasmic domain (206-237AA). Alignment of the Cherry valley duck CD8 α amino acid sequences with other known avians and *Homo sapiens* sequences is shown in Fig. 2. Overall, the cherry valley duck CD8 α deduced amino acid sequence shared about 91% identity with other known duck breeds, whilst identity with *Homo sapiens* CD8 α sequences was much lower (32%). For the structural domains, signal peptide, 75 and 94%, the transmembrane, 60 and 65% and cytoplasmic, 63 and 71% revealed, respectively high identity between both homo sapiens and chicken CD8 α and the cherry valley duck. The Cherry valley duck of extracellular region showed relatively low identity with other known ducks (88%), gallus (53%) and *Homo sapiens* (30%). The previously published demonstrations indicated that chicken CD8 α gene shows polymorphism,




Fig. 2: Differences of hydrophilicity plot, antigenicity index and surface probability plot of deduced aa sequences of extracellular region of CD8α among; *Homo sapiens* a) and Pekin b) Cherry valley duck c)




Fig. 3: Alignment of the aa sequence of cherry valley duck CD8α with other animal showed the level of sequence identity (Id%) with Cherry valley duck CD8α. The alignment was generated using ClustalW (BioManager). Dashes indicated that the gaps were introduced into the sequence to optimise the alignment. The deduced aa sequences of extracellular region of cherry valley duck CD8a ORF were predicted using interproscan which included signal peptide, extracellular, transmembrane and cytoplasmic four domains. GenBank accession numbers are as follows: Cherry valley duck CD8α FJ527828, Sheldrack CD8a FJ527912, Pekin CD8a AF378373, *Homo sapiens* CD8a NM_001768 and *Gallus gallus* CD8α AY519197

compared with CD8β but as yet there are only a few reports about *Anas platyrhynchos* CD8α sequence analyses (Koskinen *et al.*, 1999; Hu *et al.*, 2006). In this study, sequence analyses of gene scoding for differential species CD8α revealed that the deduced aa sequence had

relatively high homology (91%) compared with that of Cherry valley duck's (Fig. 3) but extracellular aa homology was significantly lower than other domains. The result was in line with Tang's study on chicken (Tang *et al.*, 2007). More significantly, the extracellular domain of




Fig. 4: a) Shows identification of recombinant proteins (rCVERCD8) by Western-blot. Lane M represents Proterm marker. Lane 1 represents identification of GST by Western-blot; Lane 2 represents identification of recombinant protein (rCVERCD8) by Western-blot; b) shows SDS-PAGE analysis of expression of PEGX-KG and PEGX-KG-CVERCD8 in the absence (1, 2) or presence (3-7) of IPTG in *E. coli*. Lane 3 represents SDS-PAGE analysis of PEGX-KG induced by IPTG for 4 h. Lanes 4-7 represent SDS-PAGE analysis of PEGX-PEGX-KG-CVERCD8 induced, respectively by IPTG for 1, 2, 3, 4 h; Lane M represents Proterm marker; c) shows SDS-PAGE analysis of purification of rabbit anti-rCVERCD8 serum. Lane M represents Proterm marker. Lane 1 represents SDS-PAGE analysis of purification of rabbit anti-rCVERCD8 serum

Cherry valley duck had relatively higher genetic polymorphism compared with Pekin and Sheldrack. Li W's studies (Li *et al.*, 2009) indicated the same results. Therefore, by DNASTar software, the differences among hydrophilicity plot, antigenicity index and surface probability plot of deduced aa sequences, CD8 α among homo sapiens, Cherry valley duck and Pekin were obvious (Fig. 2).

Construction and expression of bacterial expression vector: According to the result of sequence analysis of coding CD8 α gene, the researchers employed molecular biology techniques in prokaryotic expression of CVERCD8 sequence as immunogen of preparation for Abs in this experiment. The differentiation antigen of Cherry Valley duck in Peripheral Blood Lymphocytes (PBL) could be purified but the large variety of differentiation antigen molecules made the purification of CD8 α chains from PBL more difficult. Consequently we employed molecular biology techniques. PCR with primers F/R designed for the amplification CVERCD8 sequence resulted in 476 bp DNA fragment by agarose gel electrophoresis (Fig. 1b). The size of the fragment was similar with the expected size of CVERCD8 gene. In order to ensure correct reading frame in recombinant plasmid, the colony PCR of recombinant plasmid identification by agarose gel electrophoresis indicated the size of DNA fragment was about 470 bp. The result of double digestion identification of recombinant plasmid using EcoRI and XhoI showed about 470 bp band and 5000 bp band by agarose gel electrophoresis (Fig. 1c). Insertion of CVERCD8 gene DNA fragment in the correct reading frame was confirmed by sequencing. Expression of the recombinant plasmid (PEGX-KG-CVERCD8) in the host cell *E. coli* BL21 (DE3) was induced by IPTG. The molecular weight of *E. coli* expressed GST-tagged rCVERCD8 was about 43 kDa

(Fig. 4b). The induction with IPTG for 4h was selected in order to produce the largest amount of recombinant protein. Solubility analysis showed that the recombination protein was mainly expressed as inclusion body. Both the *E. coli*-expressed GST-tagged rCVERCD8 and GST proteins were analyzed by Western blotting analysis and the identity of the protein band was confirmed (Fig. 4a). The size of the proteins was similar to the expected size (GST-tagged rCVERCD8, 43 kDa; GST, 26 kDa). The above experiments were intended to ensure the accuracy of protein expression. Simultaneously these test results were consistent in the expected, so these data showed CVERCD8 sequence was successfully expressed.

In the study, the purpose of prokaryotic expression of CVERCD8 sequence was preparing for anti-rCVERCD8 polyclonal serum. The serum was prepared as described in methods. The results are as follows: the purified polyclonal Abs purity of saturated ammonium sulfate was about 56%. After by dextran gel column chromatography, the purity reached to about 80%. The titer of the Abs is 1:12800. There were two bands of heavy chain and light chain with purified serum by SDS-PAGE analysis (Fig. 4c). In order to ensure purity, this experiment further employed saturated ammonium sulfate and affinity chromatography (sephadex gel).

Preliminary application of Abs: All blood smears were detected by immunohistochemistry (SABC) and the recognition site of rabbit anti-rCVERCD8 serum on blood smear was the lymphocytes membrane (Fig. 5). DPV (duck plague virus) showed the affinity with a variety of epithelial cells and lymphocytes. The bursal and thymic atrophy observed in DPV infected ducklings indicated that the viruses were capable of altering defined subpopulations of lymphocytes (Shawky, 2000). For the purpose of comparing the differences between Abs of




Fig. 5: Observation of immunohistochemistry with rabbit anti-rCVERCD8 serum in blood film 400×c




Fig. 6: The Line graph of percentage of CD8 positive lymphocyte in the post-infection with DPV; a) represents the percentage of CD8 positive lymphocytes using rabbit anti-rCVERCD8 serum in group 1; b) represents the percentage of CD8 positive lymphocytes using anti-*Homo sapiens* CD8a mAb in group 1; c) represents the percentage of CD8 positive lymphocytes using rabbit anti-rCVERCD8 Abs in group 2; d) represents the percentage of CD8 positive lymphocytes using anti-*Homo sapiens* CD8a mAb in group 2

anti-*homo sapiens* CD8 α and anti-rCVERCD8, we observed the changes of peripheral blood CD8+ lymphocytes in group 1 and group 2 by immunohistochemistry (SABC). Using rabbit anti-rCVERCD8 serum the percentage of group 1 CD8+ lymphocytes fluctuated between about 31.5 and 29.8%; Using anti-*Homo sapiens* CD8 α mAb, the percentage of group 1 CD8+lymphocytes fluctuated between about 34.2 and 28.7%. Using rabbit anti-rCVERCD8 serum, the percentage of group 2 CD8+lymphocytes showed downward trend during 4 days of post-infection, the percentage started to indicate upward trend after 4 days of post-infection. Using anti-*homo sapiens* CD8 α mAb, the percentage of group 2 CD8+lymphocytes did not show regularity (Fig. 6). The result showed the percentage

of CD8+lymphocytes using anti-*Homo sapiens* CD8 α Abs was obviously instable when compared with that of using anti-rCVERCD8 serum in group 1. In group 2, the percentage of CD8+lymphocytes with anti-rCVERCD8 mAb showed significant regularity. Maybe the reason is that the number of CD8+lymphocytes in peripheral blood dramatically changed as time of the post-infected DPV. The rate of CD8+lymphocytes using anti-*Homo sapiens* CD8 α Abs was also unstable, the same as group 1. Although, immunohistochemistry was a Semi-quantitative method, the data of this experiment in synergy accord with the result of analyses of genes coding of CD8 α .

CONCLUSION

This study showed that identification of cherry valley duck CD8+lymphocytes with Abs of anti-*Homo sapiens* CD8 α has its limitations. In some immunosuppressive diseases, the use of origin Abs can be more effective on observing changes of lymphocytes subsets.

ACKNOWLEDGEMENTS

This research was supported by the grants from the Natural Science Foundation of China (30972161) and Science and Technology Research Foundation of Hubei Province (No. 2006AA202A05). The researchers would like to thank Zhang Wei (Wuhan Institute of Virology, Chinese Academy of Sciences) for her help and guide during the experiment.

REFERENCES

- Bertram, E.M., R.G. Wilkinson, B.A. Lee, A.R. Jilbert and I. Kotlarski, 1996. Identification of duck T lymphocytes using an anti-homo sapiens T cells (CD3) antiserum. *Vet. Immunol. Immunop.*, 51: 353-363.
- Bonneau, K.R., S. Ng, H. Foster, K.B. Choi and B. Berkhout *et al.*, 2008. Derivation of infectious HIV-1 molecular clones with LTR mutations: Sensitivity to the CD8+ cell noncytotoxic anti-HIV response. *Virology*, 373: 30-38.
- David, K.C. and F.G. George, 2004. CD8: Adhesion molecule, co-receptor and immunomodulator. *Mol. Cell. Biol.*, 1: 81-88.
- Fu, S.J., Z.D. Cao and J.L. Peng, 2007. Effect of feeding and fasting on excess post-exercise oxygen consumption in juvenile southern catfish (*Silurus meridionalis* Chen). *Comp. Biochem. Physiol.*, 146: 435-439.

- Gillooly, J.F., J.H. Brown, G.B. West, V.M. Savage and E.L. Charnov, 2001. Effects of size and temperature on metabolic rate. *Science*, 293: 2248-2251.
- Grodioa, J.L., E.L. Bucklesa and K.A. Schat, 2009. Production of house finch (*Carpodacus mexicanus*) IgA specific anti-sera and its application in immunohistochemistry and in ELISA for detection of *Mycoplasma gallisepticum*-specific IgA. *Vet. Immunol. Immunop.*, 132: 288-294.
- Higgins, D.A. and S.H. Chung, 1986. Duck lymphocytes. I. Purification and preliminary observations on surface markers. *J. Immunol. Methods*, 86: 231-238.
- Hu, Q.H., X.A. Jiao, Z.M. Pan, J.L. Huang and Y.L. Yin, 2006. Cloning and sequencing of CD4 and CD8 α cDNA of Chinese local chicken breeds. *Vet. Sci. China*, 37: 216-221.
- Karen, F.T. and Copeland, 2001. Non-lytic control of homo sapiens immunodeficiency virus replication and transcription by CD8-T cells. *Clin. Appl. Immunol. Res.*, 2: 1-16.
- Konno, A., K. Okada, K. Mizuno, M. Nishida and S. Nagaoki *et al.*, 2002. CD8 $\alpha\alpha$ memory effector T cells descend directly from clonally expanded CD8 $\alpha\beta$ High TCR $\alpha\beta$ T cells *in vivo*. *Immunobiology*, 100: 4090-4097.
- Koskinen, R., U. Lamminmaki, C.A. Tregaskes, J. Salomonsen, J.R. Yong and O. Vainio, 1999. Cloning and modeling of the first nonmammalian CD4. *J. Immunol.*, 162: 4115-4121.
- Kothlow, S., N.K. Mannes, B. Schaerer, D.E. Rebeski, B. Kaspers and U. Schultz, 2005. Characterization of duck leucocytes by monoclonal antibodies. *Dev. Comp. Immunol.*, 29: 733-748.
- Li, W., J.Y. Bai, Y.Y. Xu, C.Q. Gu and W.P. Zhang *et al.*, 2009. Cloning and sequence analyses of genes coding for CD3 ϵ , CD4, CD8 α of sheldrake and Cherry Valley Duck. *China Agric. Sci.*, 42: 4428-4434.
- Luhtala, M., 1998. Chicken CD4, CD8- α and CD8- β T cell co-receptor molecules. *Poult. Sci.*, 77: 1858-1873.
- Luhtala, M., C.A. Tregaskes, J.R. Young and O. Vainil, 1997. Polymorphism of chicken CD8- α , but not CD8- β . *Immunogenetics*, 46: 396-401.
- Shawky, S., 2000. Target cells for duck enteritis virus in lymphoid organs. *Avian Pathol.*, 29: 609-616.
- Shawky, S., T. Sandhu and H.L. Shivaprasad, 2000. Pathogenicity of a low virulence duck virus enteritis isolate with immunosuppressive ability. *Avian. Dis.*, 44: 590-599.
- Tang, X.S., L. Wang and D.W. Liao, 2007. Cloning, expression and production of polyclonal antibodies in extracellular regions of chicken CD8 α/β . *J. Chin. Agric. Univ.*, 12: 529-533.
- Tregaskes, C.A., F.K. Kong, E. Paramithiotis, C.L. Chen, M.J. Ratcliffe, T.F. Davison and J.R. Young, 1995. Identification and analysis of the expression of CD8 $\alpha\beta$ and CD8 $\alpha\alpha$ isoforms in chickens reveals a major TCR- γ Δ CD8 $\alpha\beta$ subset of intestinal intestinal lymphocytes. *J. Immunol.*, 154: 4485-4494.
- Xia, C., C.Y. Lin, G.X. Xu, T.J. Hu and T.Y. Yang, 2004. cDNA cloning and genomic structure of the duck (*Anas platyrhynchos*) MHC class I gene. *Immunogenetics*, 56: 304-309.
- Zhang, S., H.J. Zhang and J.D. Zhao, 2009. The role of CD4 T cell help for CD8 CTL activation. *Biochem. Biophys. Res. Commun.*, 384: 405-408.