

## The Efficacy of Antioxidative Therapy in Hepatic Fibrosis Induced Experimentally by Bile Duct Ligation in Rats

Abdulmonem Al-Hayani

Department of Anatomy, Faculty of Medicine, King Abdul Aziz University, Jeddah, Saudi Arabia

**Abstract:** Liver fibrosis which is a common result of chronic hepatic injury of diverse origins might be related to the occurrence of oxidative stress and accumulation of free radicals. It is characterized essentially by increased secretion and accumulation of different components Extracellular Matrix proteins (ECM). To investigate the beneficial effect of combined supplementation of vitamin E and selenium on hepatic fibrosis induced experimentally in bile duct-ligated rats. Forty albino rats were randomly assigned to 4 groups (10 rats in each); control group, sham-operated group Bile Duct-Ligated (BDL) group where the bile ducts of rats were ligated and BDL group co-treated with vitamin E and selenium where the rats underwent bile duct ligation and were fed with chow supplemented with vitamin E (250 mg kg<sup>-1</sup>) and selenium (0.2 mg kg<sup>-1</sup> diet) starting from three days before the operation. At the end of 3rd week, the rats were sacrificed where blood was taken for biochemical estimations of serum enzymes and bilirubin. Fresh liver tissue was taken for determination of hydroxyproline content. Also, other liver samples of 5 mm<sup>3</sup> were immediately fixed and processed for immunohistochemical demonstration of collagen type I and IV, fibronectin and laminin using the indirect immunoperoxidase method. Bile duct ligation resulted in a marked elevation in the levels of serum enzymes, bilirubin and hydroxyproline together with extensive bile duct proliferation. Strong staining was found for collagens I and IV, fibronectin and laminin in enlarged portal spaces around the newly formed bile ductules. Co-treatment with vitamin E and selenium resulted in a shift in the serum enzymes, bilirubin and hydroxyproline towards the normal values. Also, a remarkable decrease of bile duct proliferation and in the intensity of staining for different ECM components in the portal spaces was observed. Antioxidants vitamins E and selenium combination attenuate the development of hepatic fibrosis in bile duct-ligated rats. Long-term, prospective studies in humans with chronic cholestatic liver diseases may be helpful to evaluate the beneficial effects of these elements.

**Key words:** Liver fibrosis, chronic, hepatic injury, supplement, selenium, Saudi Arabia

---

### INTRODUCTION

Hepatic fibrosis which is one feature of liver cirrhosis is a highly integrated cellular response to tissue injury (Friedman, 2000). It is essentially characterized by secretion and accumulation of Extracellular Matrix proteins (ECM) (Schuppan *et al.*, 2001). The ECM of the liver comprises three sets of protein macromolecules, collagens, non-collagenous glycoproteins and glycosaminoglycans (Rojkind and Ponce-Noyola, 1982). Collagens form the major matrix component and include at least four genetically distinct subtypes which have been identified immunochemically and immunohistochemically (Rojkind *et al.*, 1979; Grimaud *et al.*, 1980; Voss *et al.*, 1980). The distribution of various matrix components varies in normal and pathological livers. Some elements, such as laminin, collagen type IV and fibronectin are concentrated in the basement membranes whereas others such as collagens types I, pro III and III are present in intercellular spaces (Martin and Timpl, 1987).

It is generally accepted that once it is established, fibrosis is irreversible (Ramachandran and Iredale, 2009). A few observations however, in man and studies with animals have led to the hypothesis that experimental hepatic fibrosis is a reversible process (Williams *et al.*, 1969; Yeong *et al.*, 1982). Animal models have been proposed for studying the diminishing of hepatic fibrosis. These include Schistosomiasis mansoni in mice (Perez-Tamayo, 1979; Mehlhorn *et al.*, 1982; Morcos *et al.*, 1985) and chemicals, e.g., CCl<sub>4</sub> and ethionine (Perez-Tamayo, 1979). Duration of the treatment and the difficulty of identifying the causes of fibrosis, however had made these models matter in debate.

Extrahepatic cholestasis is another model to investigate the formation of hepatic fibrosis. Prolonged obstruction of bile flow or hepatic diseases that lead to anatomic destruction of the biliary tree results in morphologic and biochemical changes and the development of secondary biliary cirrhosis (Popper and Schaffner, 1985). Changes induced by experimental bile

duct ligation in the rat have been partially analyzed (Carpino *et al.*, 1981; Kountouras *et al.*, 1984). They include an extensive proliferation of bile ducts in enlarged portal spaces with slight inflammation and necrosis and the formation of periportal fibrosis in <2 weeks after obstruction of the biliary tree (Aronson *et al.*, 1988). The possibility that cholestasis-induced fibrosis is a preventable phenomenon has not been explored.

Studies have shown that oxidative stress and accumulation of free radicals might contribute to the development of hepatic fibrosis in biliary obstruction (Poli *et al.*, 1987; Poli, 2000; Parola and Robino, 2001). Cholestasis was found to reduce antioxidative capacities in liver mitochondria in bile duct-ligated rats (Krahenbuhl *et al.*, 1995; Huang *et al.*, 2003). Accumulation of bile acids and inflammatory cells in the liver tissue may cause increased production of free radicals in biliary obstruction (Sokol *et al.*, 1993; Poli and Parola, 1997). Bile acids especially, enhance reactive oxygen species released by polymorpho-nuclear leukocytes (Dahm *et al.*, 1988; Singh *et al.*, 1992).

Many studies have shown that antioxidative nutrients such as vitamin E and selenium can have protective effect against chronic liver damage and cirrhosis. Both antioxidants were found to decrease significantly hepatocellular damage and elevation in the plasma enzymatic activities caused by hepatocellular damage (Geetha *et al.*, 1990; Lee and Clemens, 1992; Nagamatsu and Hasegawa, 1993; Naziroglu *et al.*, 1999). In addition, some previous studies had confirmed that dietary supplementation with vitamin E and selenium can protect hepatocytes and prevent liver fibrosis induced by CCl<sub>4</sub> in rat model (Lee *et al.*, 2001; Svegliati-Baroni *et al.*, 2001; Nieto *et al.*, 2002; Zhan *et al.*, 2003). However, the mechanism of the effects is still not clear (Li *et al.*, 2003). Therefore, in the present study, it was aimed to investigate the beneficial effect of combined supplementation of proper doses of vitamin E and selenium on liver fibrosis induced experimentally in bile duct-ligated rats.

## MATERIALS AND METHODS

**Animals and experimental design:** Forty adult male Sprague-Dawley albino rats (weighing 220-250 g at the beginning of the experiment) were used in this study. They were obtained from the Animal House of King Fahd Medical Research Center. At the beginning of the experiment the animals were kept in separate metallic cages under standard temperature (24±2°C), humidity (55±5%) and lighting (12 h: 12 h Light: Dark) conditions.

Food formed of Purina rat chow and drinking water was supplied *ad libitum*. This study was approved and registered by the Committee of Animal Investigations in Department of Anatomy, Faculty of Medicine, King Abdul-Aziz University during all the steps of the study, the animals were cared in King Fahd Center for Medical Research. The rats were randomly assigned to 4 groups (10 rats in each).

**Control group:** Rats were left without any treatment or operation.

**Bile Duct-Ligated (BDL) group:** The bile ducts of rats were ligated and transected.

**Sham-operated group:** The operation was performed to the rats in the same way as the previous group but without bile duct ligation or transection.

**BDL group co-treated with vitamin E and selenium:** The rats underwent bile duct ligation and transection and were fed with chow supplemented with vitamin E (250 mg kg<sup>-1</sup>) and selenium (0.2 mg kg<sup>-1</sup> diet) starting from 3 days before the operation.

**Operative procedure:** Rats were anesthetized with ketamin and diazepam. Laparotomy was performed under antiseptic conditions. A mid-line incision in the abdomen was made from the xiphosternum to the pubis, exposing the muscle layers and the linea alba which was then incised over a length corresponding to the skin incision. The edge of the liver was then raised and the duodenum pulled down to expose the common bile duct which pursues an almost straight course of about 3 cm from the hilum of the liver to its opening into the duodenum. There was no gall bladder and the duct was embedded for the greater part of its length in the pancreas which opens into it by numerous small ducts. A blunt needle was passed under the part of the duct selected and the duct was double ligated with 5.0 polipropilene sutures and divided in between. The peritoneum and the muscle layers as well as the skin wound were closed with cotton stitches (Kountouras *et al.*, 1984).

**Biochemical assay:** At the end of the study period (3 weeks), the rats were decapitated and blood was withdrawn from all groups of rats by puncturing retroorbital plexus. The blood samples were allowed to coagulate at room temperature. Serum was separated by centrifugation at 3,000 rpm at room temperature for 20 min and subjected to biochemical estimations of serum

transaminases, alkaline phosphatase, lactate dehydrogenase and bilirubin. Determination of serum total and direct bilirubin was done by using COBAS INTEGRA® 400 plus analyzer.

**Determination of hydroxyproline content in the liver:** At the end of the study period, fresh liver tissue (10 mg) was taken and homogenized in 1 mL of 10 N HCl with a homogenizer. This homogenate was used directly for hydroxyproline analysis. Hydroxyprolines attached to strong acid cations were exchanged with resins. Then the resins were washed with distilled water to remove substances which may have caused interference. Peptide bonds in the resins were broken at 100°C for 16 h.

The hydroxyprolines were then oxidized with a pirol derivate. Following incubation at 60°C for 25 min, the samples were kept at room temperature for 30 min to stabilize the color and the optical density of the developed color was read at 560 nm with a Shimadzu UV 1201 (Shimadzu Corp., Japan) spectrophotometer. The results were expressed as mg g<sup>-1</sup> tissue (Stegemann and Stalder, 1967).

**Fixation and immunohistochemical procedure:** At the end of the study period, fresh liver samples of 5 mm<sup>3</sup> were cut and immediately immersed in a 4% paraformaldehyde solution buffered with 0.1 M sodium cacodylate, pH 7.4 at 4°C for 4 h. Then, the samples were extensively washed in 0.1 M phosphate buffered saline at 4°C for 18 h followed by soaking in 10% glycerol in phosphate buffered saline for 1 h.

Then, the fragments were frozen in liquid nitrogen-cooled isopentane and stored at -70°C. Extracellular matrix components were localized in situ using the indirect immunoperoxidase technique (Clement *et al.*, 1985). Briefly, 5 µm cryostat sections were prepared from different frozen liver fragments and incubated in 0.05% saponin in phosphate buffered saline with 10% fetal calf serum for 1 h before a further another 1 h incubation with the specific primary antibodies.

After three phosphate buffered saline washes, sections were incubated with sheep anti-rabbit immunoglobulin labelled with peroxidase enzyme. The peroxidase activity was revealed by using 3, 3'-diaminobenzidine/H<sub>2</sub>O<sub>2</sub> solution for 20 min. Then, the sections were mounted on glass and examined by Seize light microscope (the positive reaction was visualized as a brown staining). Control liver sections were first incubated with normal immunoglobulins or directly with the medium employed to reveal peroxidase activity.

**Source of antibodies:** The specific primary antibodies against collagen type I and IV, laminin and fibronectin

were bought from Sigma Agency. These antibodies were raised in New Zealand white rabbits and purified by affinity chromatography.

**Statistical analysis:** Data are expressed as mean±Standard Deviation (SD). For all variables, the statistical differences between treated groups and the control group were analyzed statistically by student t-test, p<0.05 was considered statistically significant.

## RESULTS AND DISCUSSION

**Biochemical findings:** Table 1 and 2 showed the activities of various liver enzymes, total and conjugated bilirubin measurements in control, bile duct ligated and bile duct ligated co-treated with vitamin E and selenium groups. In bile duct ligated group there was significant and dramatic increase in serum enzyme levels, compared to control, while vitamin E and selenium co-treated groups showed a decrease in all enzyme levels towards the normal levels. Also, there was significant rise in total and conjugated bilirubin measurements which returned back towards the normal values in the bile duct ligated rats co-treated with vitamin E and selenium.

**Liver collagen content:** As shown in Table 2, hepatic injury induced by bile duct ligation caused a significant rise in the hydroxyproline content whereas vitamin E and selenium co-treatment resulted in a decrease towards the normal values.

Table 1: Showing the mean values of different liver enzymes in different groups

Group (n=10)	Aspartate transaminase (IU L <sup>-1</sup> )	Alanine transaminase (IU L <sup>-1</sup> )	Alkaline phosphatase (IU L <sup>-1</sup> )	Lactate dehydrogenase (IU L <sup>-1</sup> )
Control	67±3.3	48±4.4	175±7.8	150±6.7
Sham operated	64±4.2	54±4.8	179±8.1	149±6.6
Bile duct ligated	157±13.4**	128±11.8**	424±26.8**	331±15.9**
Bile duct ligated co-treated with vit E and selenium	82±5.1*	71±5.3*	198±9.3*	178±8.1

\*Student t-test: \*p<0.01, \*\*p<0.0001 when compared to control; ± values are expressed as mean±SD

Table 2: Showing the mean values of bilirubin and hydroxyproline in different groups

Group (n = 10)	Total bilirubin (mg dL <sup>-1</sup> )	Direct bilirubin (mg dL <sup>-1</sup> )	Hydroxyproline HP (mg g <sup>-1</sup> )
Control	0.23±0.022	0.17±0.019	4.8±0.950
Sham operated	0.24±0.023	0.18±0.022	5.1±0.820
Bile duct ligated	0.81±0.076***	0.23±0.021**	13.8±0.1.29***
Bile duct ligated co-treated with vit E and selenium	0.29±0.025*	0.2±0.020*	7.4±0.570**

Student t-test: \*p<0.05, \*\*p<0.001, \*\*\*p<0.0001, when compared to control; ± values are expressed as mean±SD




Fig. 1: Light photomicrographs of liver sections showing immuno-localization of collagen type I: (Ps = Portal space, CV = Central Vein); A) Control group: little positive staining in the portal spaces with negative sinusoidal staining. B) BDL group: strong staining of extensive fibrous septa around the proliferated bile ductules in the dilated portal spaces with little positive sinusoidal staining in the hepatic lobule. C) BDL group co-treated with vitamin E and selenium: decreased positive staining around mild proliferated bile ductules in the portal spaces with negative sinusoidal staining in the hepatic lobule

**Immunolocalization of collagens, fibronectin and laminin**

**Livers of normal rats:** As shown in the Fig. 1-4 and the major sites of deposition of the different components were in the portal spaces (around the basement membranes of the bile ducts and blood vessels) and around the central veins in the hepatic lobules. All the components were




Fig. 2: Light photomicrographs of liver sections showing immunolocalization of collagen type IV: (Ps = Portal space, CV = Central Vein); A) Control group: strong staining was seen in the portal spaces around the basement membranes of bile ductules and blood vessels with moderate and discontinuous sinusoidal staining in the hepatic lobule. B) BDL group: strong staining of basement membranes around the proliferated bile ductules in the dilated portal spaces with positive sinusoidal staining in the hepatic lobule. C) BDL group co-treated with vitamin E and selenium: decreased positive staining around mild proliferated bile ductules in the portal spaces with positive sinusoidal staining in the hepatic lobule

visualized although the intensity of staining and the manner of distribution varied from one component to




Fig. 3: Light photomicrographs of liver sections showing immunolocalization of fibronectin: (Ps = Portal space, CV = Central Vein); A) Control group: strong staining was seen in the portal spaces around bile ducts and blood vessels with strong continuous sinusoidal staining in the hepatic lobules and slight cytoplasmic staining. B) BDL group: strong staining of extensive fibrous septa around the proliferated bile ductules in the dilated portal spaces with little positive sinusoidal staining in the hepatic lobule. C) BDL group co-treated with vitamin E and selenium: decreased positive staining around mild proliferated bile ductules in the portal spaces with strong positive sinusoidal staining in the hepatic lobule

another. The interstitial type I collagen was present in the portal tracts while the liver sinusoids showed negative staining with it. The basement membrane type IV collagen appeared in the portal tracts around blood vessels and

bile ducts. In addition, in the hepatic lobules, liver sinusoids showed a moderate and discontinuous sinusoidal staining with Col IV. Regarding the fibronectin, it was the most prominent component, which showed very strong staining in the portal spaces and a continuous sinusoidal labeling in the hepatic lobules with slight cytoplasmic staining. The laminin showed a moderate staining, especially around the bile ducts and blood vessels while in the hepatic lobules, only faint and continuous sinusoidal staining could be detected.

**Livers of bile-duct ligated rats:** As shown in the Fig. 1-4, the biliary obstruction resulted in a rapid extensive bile duct proliferation and fibrogenesis in enlarged portal areas. Infiltration of the portal tract with polymorphonuclear inflammatory cells was occasional and hepatocyte necrosis was nearly absent. Central vein areas remained unchanged. By the immunostaining technique, extensive fibrous septa were stained for collagens I and IV, fibronectin and laminin. The most intense staining was found in portal spaces. Intense staining for collagen IV and laminin was observed around bile duct proliferations. In the sinusoids, collagen 4 was continuously distributed along hepatocyte cords. Laminin and fibronectin were deposited mainly within the periportal areas and in the borders of septa. Regarding the fibronectin, a strong staining was seen in the portal spaces around the basement membranes of bile ducts and blood vessels, with strong sinusoidal staining in the hepatic lobules.

**Livers of bile-duct ligated rats co-treated with vitamin E and selenium:** As shown in the Fig. 1-4, the co-administration of vitamin E and selenium resulted in much regression of bile duct proliferation and fibrogenesis in enlarged portal areas. The distribution of collagens, fibronectin and laminin was in some extent similar to that observed in normal liver. In portal spaces, the remaining clusters of neo-formed bile ducts were still surrounded by laminin and collagen IV. In the sinusoids, the distribution of collagens I, laminin and fibronectin was similar to that in normal liver. Only collagen IV remained abundant in the sinusoids and continuously deposited all along hepatocyte cords.

Liver fibrosis is the common result of chronic hepatic injury of diverse origins such as chronic viral infections, metabolic/storage diseases (hemochromatosis), helminthic infections (schistosomiasis), chronic toxin exposure (alcohol and environmental poisons) and biliary obstruction (biliary cirrhosis). In end stage liver fibrosis or cirrhosis, the liver Extracellular Matrix (ECM) may contain up to 6-10 times more collagen and proteoglycans than in the normal state (Gressner *et al.*, 2007; Wells, 2007).




Fig. 4: Light photomicrographs of liver sections showing immunolocalization of laminin: (Ps = Portal space, CV = Central Vein); A) Control group: moderate staining was seen in the portal spaces around the basement membranes of bile ducts and blood vessels with faint continuous sinusoidal staining in the hepatic lobules. B) BDL group: strong staining of basement membranes around the proliferated bile ductules in the dilated portal spaces with moderate sinusoidal staining in the hepatic lobule. C) BDL group co-treated with vitamin E and selenium: decreased positive staining around mild proliferated bile ductules in the portal spaces with negative sinusoidal staining in the hepatic lobule

Because the ECM support of the liver parenchyma is particularly critical to its function, research that emphasizes the nature of liver ECM, the molecular

regulation of the turnover of its components and identification of liver cells responsible for the synthesis of matrix proteins are especially crucial for the ultimate design of effective therapies for liver fibrosis (Schuppan, 1990).

The use of animal models of liver fibrosis such as the administration of liver toxins  $\text{CCl}_4$  (Rojkind *et al.*, 1977), dimethylnitrosamine (Risteli *et al.*, 1976), alcohol (Mezey *et al.*, 1976) have greatly helped in the characterization of the temporal expression of various components of the ECM during fibrogenesis. It was reported that there is increase in the amounts of collagens types I-VI (Wu *et al.*, 1982; Isaka *et al.*, 1996). It was reported that in early fibrosis, the amounts of types III and IV collagens increase relative to other collagens. In late fibrosis, type I collagen predominates (Schuppan, 1990). Other components of liver ECM such as laminin, fibronectin and proteoglycans are also increased in fibrosis (Rojkind *et al.*, 1979; Geerts *et al.*, 1986; Reid *et al.*, 1992).

The reversibility of hepatic fibrosis remains the subject of controversy. In this study, it was decided to investigate fibrogenesis in cholestatic rat liver experimentally induced by common bile duct ligation. This model is morphologically characterized by marked proliferation of bile ducts and formation of enlarged fibrous septa with only limited inflammation and necrosis. Dramatic and early increase in the serum concentration of bilirubin and enzymes indicated that extrahepatic cholestasis was induced rapidly in this experimental model. Moreover, the bile duct ligation model of rats mimics best the clinical and histopathological aspects of hepatic fibrosis secondary to extrahepatic biliary obstruction in humans. This model allows researchers to study in detail all the developmental stages of hepatic fibrosis, ultimately, terminating in cirrhosis within about 4 weeks (Moritz and Snodgrass, 1972; Kountouras *et al.*, 1984).

Elevated tissue collagen levels can be measured directly or can be indirectly evaluated by the measurement of hydroxyproline levels (Stegemann and Stalder, 1967). In this study, the bile duct ligation resulted in highly significant rise in serum hydroxyproline levels when compared to the control values. It was demonstrated that liver fibrosis is accompanied by a significant increase in collagen content in this organ (Bankowski, 1994; Matsuda *et al.*, 1997). Collagen is involved in wound healing via formation of hydroxylation. However, the balance of collagen is important in wound healing. If an uncontrolled accumulation of collagen cannot be degraded by collagenases, it will result in fibrosis

formation instead of normal healthy hepatocytes. The immunohistochemical study in this research demonstrated that fibrosis is formed by interstitial collagen types I, fibronectin and basement membrane proteins. In agreement, it was reported that as bile duct cells actively proliferated, it can be expected that they were the producers of basement membrane components, i.e., collagen IV and laminin and were responsible for the accumulation of these proteins in portal areas (Miyabayashi *et al.*, 1987; Clement *et al.*, 1988).

Studies have shown that oxidative stress (lipid peroxides) is one of the mechanisms in the development of liver fibrosis (Lee *et al.*, 2001; Svegliati-Baroni *et al.*, 2001; Nieto *et al.*, 2002; Zhan *et al.*, 2003); it is worth investigating the therapeutic role of antioxidative nutrients in hepatic fibrosis. Over the years, many compounds have been studied as possible protectors for liver fibrosis, like antioxidants such as flavonoids and other phenolic derivatives. These compounds belong to a group of a bigger family known as phytochemicals. They are widely found in fruits and vegetables and while not having an energetic value are an important part of the human diet. Flavonoids exhibit a wide variety of biological properties, including hepatoprotection and inhibition of fibrosis (Middleton *et al.*, 2000). Studies demonstrating the beneficial effect of vitamin E supplementation in prevention of both enhancement of lipid peroxidation and synthesis of collagens might support this suggestion (Houglum *et al.*, 1991; Parola *et al.*, 1992; Chojkier *et al.*, 1998). However, it has been reported that treatment of bile duct-ligated rats with vitamin E completely prevented the increase in lipid peroxidation in liver and plasma but failed to prevent tissue injury histologically.

Thus, the current controversies between the performed studies to evaluate the role of oxidative stress in tissue injury induced by bile duct ligation make newer studies to be performed necessary (Baron and Muriel, 1999; Friedman, 2000; Schuppan *et al.*, 2001). Selenium is a trace element which has antioxidant properties because it is an essential component of some oxidoreductase enzymes. Selenium deficiency is associated with higher levels of ROS and lipid peroxidation which can be reversed by selenium supplementation (Luoma *et al.*, 1984; Bedwal *et al.*, 1993). In one study, administration of vitamin E together with Se and zinc resulted in a lowered 1 month mortality in patients with alcoholic hepatitis (Wenzel *et al.*, 1993). In other studies, however, administration of Se together with vitamins A and E did not affect survival of patients with alcoholic hepatitis (Stewart *et al.*, 2007).

Selenium-enriched lactobacillus decreased liver injury and lipid peroxidation induced by CCl<sub>4</sub> administration in mice (Chen *et al.*, 2005) while supplementation of the diet with vitamin E and Se decreased hepatic fibrosis produced in rats by acute and chronic CCl<sub>4</sub> administration (Shen *et al.*, 2005). In this study, co-treatment with vitamin E and selenium resulted in an evident shift to normal liver architecture and functions including normal values in serum enzyme activities and bilirubin levels suggesting that massive and selective disruption of bile ducts and death of the newly formed bile duct epithelial cells rapidly occurred.

The dosage of vitamin E and selenium used in the study was selected as previously described (Tsukamoto *et al.*, 1990; Zhang *et al.*, 1996; Li *et al.*, 2003). In accordance, it was reported that several antioxidants were scavenged free radicals (Sies *et al.*, 1992; McPherson, 1994) and these radicals in hepatic cellular structures may increase the cellular degeneration. This process may be able to affect the levels of liver enzymes due to damage of cellular membrane.

Thus, abnormal levels of the liver enzymes in plasma are usually indicative of the hepatic cellular injury in experimental animals (Albano *et al.*, 1989; Miao *et al.*, 1990; Biasi *et al.*, 1991; Parola *et al.*, 1992; Knook *et al.*, 1995). Also, the results from histology and serum enzymes showed that vitamin E and selenium reduced liver damage during liver injury and had the effect of decreasing collagen fibers thus preventing liver fibrosis. Immunostaining of different component of ECM showed that the amount of collagen fibers decreased markedly in vitamin E and selenium co-treated group when compared to the bile duct ligated group, suggesting that these elements can prevent liver damage during liver injury, by affecting cellular sources of ECM (Iredale *et al.*, 1998). In accordance, a recently published study, the combination of vitamins E and C was associated with decreased ethanol-induced hepatic glutathione peroxidase activity and hepatic fibrosis in protein-deficient rats fed with a high-fat liquid diet (Soylu *et al.*, 2006).

## CONCLUSION

In the study, oxidative stress might contribute to the pathogenesis of secondary biliary fibrosis. Antioxidants vitamins E and selenium combination attenuate the development of hepatic fibrosis in bile duct-ligated rats. Long-term, prospective studies in humans with chronic cholestatic liver diseases may be helpful to evaluate the beneficial effects of these elements.

## REFERENCES

- Albano, E., R. Carini, M. Parola, G. Bellomo, L. Goria-Gatti, G. Poli and M.U. Dianzani, 1989. Effect of carbon tetrachloride on calcium homeostasis: A critical reconsideration. *Biochem. Pharmacol.*, 38: 2719-2725.
- Aronson, D.C., J. de Haan, J. James, K.S. Bosch, A.G. Ketel, J.M. Houtkooper and H.A.S. Heijmans, 1988. Quantitative aspects of the parenchymal-stroma relationship in experimentally induced cholestasis. *Liver*, 8: 116-126.
- Bankowski, E., 1994. Collagen in liver fibrosis induced by ethanol. *Rocz. Akad. Med. Białymst.*, 39: 1-6.
- Baron, V. and P. Muriel, 1999. Role of glutathione, lipid peroxidation on acute bile-duct obstruction in the rat. *Biochem. Biophys. Acta*, 1472: 173-180.
- Bedwal, R.S., N. Nair, M.P. Sharma and R.S. Mathur, 1993. Selenium-its biological perspectives. *Med. Hypotheses*, 41: 150-159.
- Biasi, F., E. Albano, E. Chiarpotto, F.P. Corongiu and M.A. Pronzato *et al.*, 1991. *In vivo* and *in vitro* evidence concerning the role of lipid peroxidation in the mechanism of hepatocyte death due to carbon tetrachloride. *Cell Biochem. Funct.*, 9: 111-118.
- Carpino, F., E. Gaudio, G. Marinozzi, M. Melis and P.M. Motta, 1981. Scanning and transmission electron microscopic study of experimental extrahepatic cholestasis in the rat. *J. Submicrosc. Cytol.*, 13: 581-598.
- Chen, L., D.D. Pan, J. Zhou and Y.Z. Jiang, 2005. Protective effect of selenium enriched *Lactobacillus* on  $\text{CCl}_4$  induced liver injury in mice and its possible mechanisms. *World J. Gastroenterol.*, 11: 5795-5800.
- Chojkier, M., K., Houghlum, K.S. Lee and M. Buck, 1998. Long- and shortterm D-alpha-tocopherol supplementation inhibits liver collagen alpha1(I) gene expression. *Am. J. Gastrointest. Physiol.*, 275: G1480-G1485.
- Clement, B., M. Rissel, S. Peyrol, Y. Mazurier, J.A. Grimaud and A. Guillouzo, 1985. A procedure for light and electron microscopic intracellular immunolocalization of collagen and fibronectin in rat liver. *J. Histochem. Cytochem.*, 33: 407-414.
- Clement, B., P.Y. Rescan, G. Baffet, O. Loreal, D. Lehry, J.P. Campion and A. Guillouzo, 1988. Hepatocytes may produce laminin in fibrotic liver and in primary culture. *Hepatology*, 8: 794-803.
- Dahm, L.J., J.A. Hewett and R.A. Roth, 1988. Bile and bile salts potentiate superoxide anion release from activated rat peritoneal neutrophils. *Toxicol. Applied Pharmacol.*, 95: 82-92.
- Friedman, S.L., 2000. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. *J. Biol. Chem.*, 275: 2247-2250.
- Geerts, A., H.J. Geuze, J. W. Slot, B. Voss, D. Schuppan, P. Schellinck and E. Wisse, 1986. Immunological localization of procollagen III, fibronectin and heparan sulfate proteoglycan on ultrathin frozen sections of the normal rat liver. *Histochem.*, 84: 355-362.
- Geetha, A., R. Sankar, T. Marar and C.S.S. Devi, 1990.  $\alpha$ -Tocopherol reduces doxorubicin-induced toxicity in rats-histological and biochemical evidences. *Indian J. Physiol. Pharmacol.*, 34: 94-100.
- Gressner, O.A., R. Weiskirchen and A.M. Gressner, 2007. Evolving concepts of liver fibrogenesis provide new diagnostic and therapeutic options. *Comp. Hepatol.*, 6: 7-7.
- Grimaud, J.A., M. Druguet, S. Peyrol, O. Chevalier, D. Herbage and N. El Badrawy, 1980. Collagen immunotyping in human liver: Light and electron microscope study. *J. Histochem. Cytochem.*, 28: 1145-1156.
- Houghlum, K., D.A. Brenner and M. Chojkier, 1991. d-alpha-tocopherol inhibits collagen alpha 1(I) gene expression in cultured human fibroblasts. Modulation of constitutive collagen gene expression by lipid peroxidation. *J. Clin. Invest.*, 87: 2230-2235.
- Huang, Y.T., Y.C. Hsu, C.J. Chen, C.T. Liu and Y.H. Wei, 2003. Oxidativestress-related changes in the livers of bile-duct-ligated rats. *J. Biomed. Sci.*, 10: 170-178.
- Iredale, J.P., R.C. Benyon, J. Pickering, M. McCullen and M. Northrop *et al.*, 1998. Mechanisms of spontaneous resolution of rat liver fibrosis. Hepatic stellate cell apoptosis and reduced hepatic expression of metalloproteinase inhibitors. *J. Clin. Investigat.*, 102: 538-549.
- Isaka, Y., D.K. Brees, K. Ikegaya, Y. Kaneda, E. Imai, N.A. Noble and W.A. Border, 1996. Gene therapy by skeletal muscle expression of decorin prevents fibrotic disease in rat kidney. *Nat. Med.*, 2: 418-423.
- Knook, D.L., A. Bosma and W.F. Seifert, 1995. Role of vitamin A in liver fibrosis. *J. Gastroenterol. Hepatol.*, 10: 47-49.
- Kountouras, J., B.H. Billing and P.J. Scheuer, 1984. Prolonged bile duct obstruction: A new experimental model for cirrhosis in the rat. *Br. J. Exp. Pathol.*, 65: 305-311.
- Krahenbuhl, S., C. Talos, B. Lauterburg and J. Reichen, 1995. Reduced antioxidative capacity in liver mitochondria from bile duct ligated rats. *Hepatology*, 22: 607-612.
- Lee, K.S., S.J. Lee, H.J. Park, J.P. Chung and K.H. Han *et al.*, 2001. Oxidative stress effect on the activation of hepatic stellate cells. *Yonsei Med. J.*, 42: 1-8.



- Lee, S. and M.G. Clemens, 1992. Effect of  $\alpha$ -tocopherol on hepatic mixed function oxidases in hepatic ischemia reperfusion. *Hepatology*, 15: 276-281.
- Li, F., X.H. Li, W.F. Cheng and L.M. Xie, 2003. Effects of antifibrosis and antioxidation of dietary supplement vitamin E and selenium on experimental model of rats. *Yingyang Xuebao*, 25: 60-65.
- Luoma, P.V., E.A. Sotaniemi, H. Korpela and J. Kumpulainen, 1984. Serum selenium, glutathione peroxidase activity and high density lipoprotein cholesterol-effect of selenium supplementation. *Res. Commun. Chem. Pathol. Pharmacol.*, 46: 469-472.
- Martin, G.R. and R. Timpl, 1987. Laminin and other basement membrane components. *Annu. Rev. Cell Biol.*, 3: 57-85.
- Matsuda, Y., K. Matsumoto, A. Yamada, T. Ichida and H. Asakura *et al.*, 1997. Preventive and therapeutic effects in rats of hepatocytes growth factor infusion on liver fibrosis/cirrhosis. *Hepatology*, 26: 81-89.
- McPherson, A., 1994. Selenium Vitamin E and Biological Oxidation. In: *Recent Advances in Animal Nutrition*, Cole D.J. and P.J. Garnsworthy (Eds.). Butterworth and Heinemann's, Oxford, pp: 3-30.
- Mehlhorn, H., J.K. Frenkel, P. Andrews and H. Thomas, 1982. Light and electron microscopic studies on *Schistosoma mansoni* granulomas of mouse livers following treatment with praziquantel. *Trop. Med. Parasitol.*, 33: 229-239.
- Mezey, E., J.J. Potter and W.C. Maddrey, 1976. Hepatic collagen proline hydroxylase activity in alcoholic liver disease. *Clin. Chem. Acta.*, 68: 313-320.
- Miao, S., W. Bao-En, G. Annoni, S.D. Esposti, L. Biempica, M. Zern, 1990. Two rat models of hepatic fibrosis. A morphometric and molecular comparison. *Lab. Invest.*, 63: 467-475.
- Middleton, Jr. E., C. Kandaswami and T.C. Theoharides, 2000. The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease and cancer. *Pharmacol. Rev.*, 52: 673-751.
- Miyabayashi, C., T. Kojima, K. Inoue, H. Sasaki, Y. Muragaki and A. Ooshima, 1987. Ultrastructural localisation of type IV collagen, laminin and prolyl hydroxylase in biliary epithelial cells of rat liver following ligation of the common bile duct. *Gastroenterol. Jpn.*, 2: 354-369.
- Morcos, S.H., M.T. Khayyal, M.M. Mansour, S. Salen, E.A. Ishak and N.I. Girgis, 1985. Reversal of hepatic fibrosis after praziquantel therapy of murine schistosomiasis. *Am. J. Trop. Med. Hyg.*, 34: 314-321.
- Moritz, M. and P.J. Snodgrass, 1972. Serum enzymes derived from liver cell fractions. II. Responses to bile duct ligation in rats. *Gastroenterol.*, 62: 93-100.
- Nagamatsu, K. and A. Hasegawa, 1993. Effect of sodium selenite on morphine-induced hepatotoxicity in mice. *Drug. Chem. Toxicol.*, 16: 241-253.
- Naziroglu, M., M. Cay, B. Ustundag, M. Aksakal and H. Yekeler, 1999. Protective effects of vitamin E on carbon tetrachloride-induced liver damage in rats. *Cell Biochem. Funct.*, 17: 253-259.
- Nieto, N., S.L. Friedman and A.I. Cederbaum, 2002. Stimulation and proliferation of primary rat hepatic stellate cells by cytochrome P450 2E1-derived reactive oxygen species. *Hepatology*, 35: 62-73.
- Parola, M. and G. Robino, 2001. Oxidative stress-related molecules and liver fibrosis. *J. Hepatology*, 35: 297-306.
- Parola, M., G. Leonarduzzi, F. Biasi, E. Albano, M.E. Biocca, G. Poli and M.U. Dianzani, 1992. Vitamin E dietary supplementation protects against CCl<sub>4</sub> induced chronic liver damage and cirrhosis. *Hepatology*, 16: 1014-1021.
- Perez-Tamayo, R., 1979. Cirrhosis of the liver: A reversible disease. *Pathol. Annu.*, 14: 183-213.
- Poli, G. and M. Parola, 1997. Oxidative damage and fibrogenesis. *Free Radic. Biol. Med.*, 22: 287-305.
- Poli, G., 2000. Pathogenesis of liver fibrosis: Role of oxidative stress. *Mol. Aspects Med.*, 21: 49-98.
- Poli, G., E. Albano and M.U. Dianzani, 1987. The role of lipid peroxidation in liver damage. *Chem. Phys. Lipids*, 45: 117-142.
- Popper, H. and F. Schaffner, 1985. *Cholestasis, Gastroenterology*. 4th Edn., Vol. 5, W.B. Saunders, Philadelphia, pp: 2697-2731.
- Ramachandran, P. and J.P. Iredale, 2009. Reversibility of liver fibrosis. *Ann. Hepatology*, 8: 283-291.
- Reid, L.M., A.S. Fiorino, S.H. Sigal, S. Brill and P.A. Holst, 1992. Extracellular matrix gradients in the space of Disse: Relevance to liver biology. *Hepatology*, 15: 1198-1203.
- Risteli, J., L. Tuderman and K.I. Kivirikko, 1976. Intracellular enzymes of collagen biosynthesis in rat liver as a function of age and in hepatic injury produced by dimethyl-nitrosamine. *Biochem. J.*, 158: 369-376.
- Rojkind, M., M.A. Giambrone and L. Biempica, 1979. Collagen types in normal and cirrhotic liver. *Gastroenterol.*, 76: 710-719.
- Rojkind, M., M.A. Giambrone and M. Ehrenpreise, 1977. Proline oxidase activity and the availability of proline for collagen biosynthesis in livers of CCl<sub>4</sub> treated rats. *Gastroenterol.*, 73: 12-43.
- Rojkind, M. and P. Ponce-Noyola, 1982. The extracellular matrix of the liver. *Coll. Relat. Res.*, 2: 151-175.
- Schuppan, D., 1990. Structure of the extracellular matrix in normal and fibrotic liver: Collagens and glycoproteins. *Seminars Liver Dis.*, 10: 1-10.

- Schuppan, D., M. Ruehl, R. Somasundaram and E.G. Hahn, 2001. Matrix as a modulator of hepatic fibrogenesis. *Semin. Liver Dis.*, 21: 351-372.
- Shen, X.H., W.F. Cheng, X.H. Li, J.Q. Sun, F. Li, L. Ma and L.M. Xie, 2005. Effects of dietary supplementation with vitamin E and selenium on rat hepatic stellate cell apoptosis. *World J. Gastroenterol.*, 11: 4957-4961.
- Sies, H., W. Stahl and A.R. Sundquist, 1992. Antioxidant functions of vitamins: Vitamin C and E beta-carotene and other carotenoids. *Ann. N.Y. Acad. Sci.*, 669: 7-20.
- Singh, S., G. Shackleton, E. Ah-Sing, J. Chakraborty and M.E. Bailey, 1992. Antioxidant defenses in the bile duct-ligated rat. *Gastroenterol.*, 103: 1625-1629.
- Sokol, R.J., M. Devereaux, R. Khandwala and K. O'Brien, 1993. Evidence for involvement of oxygen free radicals in bile acid toxicity to isolated rat hepatocytes. *Hepatology*, 17: 869-881.
- Soylu, A.R., S. Altaner, N. Aydogdu, U.N. Basaran and O. Tarcin *et al.*, 2006. Effects of vitamins E and C supplementation on hepatic glutathione peroxidase activity and tissue Injury associated with ethanol ingestion in malnourished rats. *Curr. Ther. Res. Clin. Exp.*, 67: 118-137.
- Stegemann, H. and K. Stalder, 1967. Determination of hydroxyproline. *Clin. Chim. Acta.*, 18: 267-273.
- Stewart, S., M. Prince, M. Bassendine, M. Hudson and O. James *et al.*, 2007. A randomized trial of antioxidant therapy alone or with corticosteroids in acute alcoholic hepatitis. *J. Hepatology*, 47: 277-283.
- Svegliati-Baroni, G., S. Saccomanno, H. Van Goor, P. Jansen, A. Benedette and H. Moshage, 2001. Involvement of reactive oxygen species and nitric oxide radicals in activation and proliferation of rat hepatic stellate cells. *Liver*, 21: 1-12.
- Tsukamoto, H., M. Matsuoka and S.W. French, 1990. Experimental models of hepatic fibrosis: A review. *Semin. Liver Dis.*, 10: 56-65.
- Voss, B., J. Rauterberg, S. Allam and G. Pott, 1980. Distribution of collagen tyoe I and type III and of two collgenous components of basement membranes in the human liver. *Pathol. Res. Pract.*, 170: 50-60.
- Wells, R.G., 2007. Function and Metabolism of Collagen and other Extracellular Matrix Proteins. In: *The Textbook of Hepatology: From Basic Science to Clinical Practice*, Rhodes, J.T. (Ed.). 3rd Edn., Blackwell Publishing, Oxford.
- Wenzel, G., B. Kuklinski, C. Ruhlmann and D. Ehrhardt, 1993. Alkoholtoxische hepatitis-eine freie radikale assoziierte erkrankung letalitassenkung durch adjuvante antioxidantientherapie. *Imm. Med.*, 48: 490-496.
- Williams, R., P. Smith, E. Spicer, M. Barry and S. Sherlock, 1969. Venesection therapy in idiopathic haemochromatosis. *Q. J. Med.*, 38: 1-16.
- Wu, C.H., M.A. Giambrone, D.J. Howard, M. Rojkind and G.Y. Wu, 1982. The nature of collagen of hepatic fibrosis in advanced murine schistosomiasis. *Hepatology*, 2: 366S-371S.
- Yeong, M.L., G.I. Nicholson and S.P. Lee, 1982. Regression of biliary cirrhosis following choledochal cyst drainage. *Gastroenterol.*, 82: 332-335.
- Zhan, Y., Y. Wang, L. Wei and H. Chen, 2003. Effects of vitamin E on the proliferation and collagen synthesis of rat hepatic stellate cells treated with IL-2 or TNF-alpha. *Chin. Med. J.*, 116: 472-474.
- Zhang, M., G. Song and G.Y. Minuk, 1996. Effects of hepatic stimulator substance, herbal medicine, selenium/vitamin E and ciprofloxacin on cirrhosis in the rat. *Gastroenterology*, 110: 1150-1155.