

## Effect of Different Levels of DL-Methionine Replaced with Betafin on Some of Blood Parameters on Broiler Chickens

M. Baghaei, M. Eslami, M. Chaji, M. Mamoue and M. Bojarpour  
Department of Animal Science,

Ramin (Khuzestan) Agricultural and Natural Resources University, Mollassani, Khuzestan, Iran

**Abstract:** This experiment was conducted to evaluate the effect of dietary Betafin (Betaine anhydrous 97%) supplementation as a replacement for DL-Methionine on blood parameters of 42 days old broiler chickens. Total 300 days old Ross 308 broilers were used in a completely randomized design with 4 treatment and 5 replicates in each treatment and 15 birds/replicates and reared on the floor pens for 42 days. A basal diet was formulated as control according to NRC recommendations for starter (1-21 days) and grower (22-42 days) periods. In experimental diets, DL-Methionine levels were formulated either according to NRC (control (T<sub>1</sub>)) or at 90 (T<sub>2</sub>), 80 (T<sub>3</sub>) and 70% (T<sub>4</sub>) of the control. The incomplete levels of the Methionine in T<sub>2</sub>-T<sub>4</sub> supplemented by adding Betafin to the diets. The results indicated that by replacement of Betafin instead of DL-Methionine in diets of broiler blood parameters was not altered significantly. Although, in birds fed T<sub>4</sub>, the concentration of the all measured blood parameters except HDL was the maximum or minimum when compared to other groups.

**Key words:** Betafin, blood, broiler, DL-methionine, HDL, Iran

### INTRODUCTION

Betaine is a natural compound having important functions in animal metabolism. Chemically, betaine (glycine betaine, trimethylglycine) is a quaternary ammonium compound (Yancey *et al.*, 1982). The unique chemical properties of betaine are due to its bipolar structure and its chemically reactive methyl groups which it can donate in methylation reactions. Betaine is chemically stable and nontoxic.

Hence, betaine is not present in large quantities in most feedstuffs. Betaine plays an important function in the amino acid, lipid metabolism and antibody production in practical animal production (Fernandez *et al.*, 1998; Verreschi *et al.*, 2002; Kim *et al.*, 2003; Zulkifli *et al.*, 2004). Manifold studies available about effect of Betaine on performance and carcass characteristic of broiler chickens (Sun *et al.*, 2008; Kermanshahi, 2001; Esteve-Garcia and Mack, 2000; Schutte *et al.*, 1997) but there are limit information about effect of Betaine on hematological indices of broiler chickens.

Therefore, the purpose of this study was investigated the effect of using different levels of DL-Methionine replaced with Betafin on some of blood parameters on broiler chickens.

### MATERIALS AND METHODS

**Bird and diet:** In this study, 300 broiler chickens of the commercial Ross 308 strain were used in a completely

randomized design with 4 treatment and 5 replicates in each treatment and 15 birds/replicates and reared on the floor pens for 42 days. A basal diet was formulated as control according to NRC (1994) recommendations for starter (1-21 days) and grower (22-42 days) periods (Table 1).

Table 1: Ingredient composition (as percent of dry matter) and calculated analysis of the basal diets

Ingredients	Starter (1-21 days)	Grower (22-42 days)
Corn	58.70	61.00
Soybean meal	30.00	29.00
Wheat bran	5.00	5.00
Fish meal	2.00	0.00
Soybean oil	1.00	2.00
Oyster shell meal	1.20	1.00
DCP	1.07	1.00
Vitamin and mineral premix	0.50	0.50
DL-Methionine	0.13	0.10
L-lysine	0.15	0.25
Salt	0.25	0.10
Coccidiostat	0.00	0.05
Total	100.00	100.00
<b>Nutrient content</b>		
ME (Kcal kg <sup>-1</sup> )	2850.00	2950.00
Crude protein (%)	20.48	18.44
Crude fiber (%)	3.89	3.81

Vitamin and mineral provided per kilogram of diet: vitamin A, 360,000 IU; vitamin D3, 80,000 IU; vitamin E, 7200 IU; vitamin K3, 800 mg; vitamin B1, 720 mg; vitamin B9, 400 mg; vitamin H2, 40 mg; vitamin B2, 2640 mg; vitamin B3, 4000 mg; vitamin B5, 12000 mg; vitamin B6, 1200 mg; vitamin B12, 6 mg; Choline chloraid, 200,000 mg; Manganese, 40000 mg; Iron, 20000 mg; Zinc, 40000 mg; copper, 4000 mg; Iodine, 400 mg; Selenium, 80 mg

**Corresponding Author:** Morteza Chaji, Department of Animal Science, Ramin Agricultural and Natural Resources University, Mollassani, Khuzestan, Iran

**Table 2: The effect of treatments on serum lipid concentrations of broilers**

Treatments	Parameters					
	Glucose (mg dL <sup>-1</sup> )	Cholesterol (g dL <sup>-1</sup> )	Triglycerides (g dL <sup>-1</sup> )	HDL (mg dL <sup>-1</sup> )	LDL (mg dL <sup>-1</sup> )	VLDL (mg dL <sup>-1</sup> )
T <sub>1</sub>	218.62±2.10	121.34±2.38	27.79±1.24	41.08±0.94	74.67±3.03	5.56±0.25
T <sub>2</sub>	221.79±0.70	122.19±1.54	28.65±0.46	40.34±0.95	76.12±2.41	5.73±0.09
T <sub>3</sub>	222.53±4.19	123.21±1.08	29.16±0.26	40.07±0.55	77.31±1.25	5.83±0.05
T <sub>4</sub>	223.93±3.32	124.02±1.22	30.59±2.26	39.18±0.51	78.71±1.55	6.11±0.45
<b>p-values</b>	0.63	0.68	0.53	0.45	0.63	0.54

\*c means in each column with different superscripts are significant different (p<0.05)

In experimental diets, Methionine levels were as formulated (control (T<sub>1</sub>)) or at 90 (T<sub>2</sub>), 80 (T<sub>3</sub>) and 70% (T<sub>4</sub>) of the control. The incompletes levels of the Methionine in T<sub>2</sub>-T<sub>4</sub> supplemented by adding Betafin to the diets. During the experiment birds had *ad libitum* access to feed and water.

**Sample collection:** At 42nd day of the experimental period, 5 mL of blood was collected from wing vein from 6 birds in each treatment. Blood samples were centrifuged (at 3,000×rpm for 15 min) and serum was separated and then stored at -20°C until assayed for measuring blood parameters (glucose, cholesterol, triglycerides and High Density Lipoprotein cholesterol (HDL)) using appropriate laboratory kits (Friedewald *et al.*, 1972; Gordon *et al.*, 1977; Gowenlock *et al.*, 1988). Very Low Density Lipoprotein (VLDL) was calculated from triglycerides by dividing the factor by 5. The Low Density Lipoprotein cholesterol (LDL) was calculated by using the equation:

$$\text{LDL cholesterol} = \text{Total cholesterol} - \text{HDL cholesterol} - \text{VLDL cholesterol}$$

**Statistical analysis:** All data were analyzed using the One-Way ANOVA procedure of SAS® (SAS Institute Inc, 1998) for analysis of variance. Significant differences among treatments were identified at 5% level by Duncan’s multiple range tests.

**RESULTS AND DISCUSSION**

**Blood parameters:** The effect of experimental treatments on blood parameters are shown in Table 2. Treatments had no significant effect on serum cholesterol, glucose, triglycerides, HDL, LDL and VLDL content of blood, although in birds fed T<sub>4</sub>, the concentration of the all measured blood parameters, except HDL were different in compare with the control treatment. These results are in agreement with Konca *et al.* (2008) and Attia *et al.* (2005), they indicated that betaine supplementation had no significant effect on most of serum constituents. However, Hassan *et al.* (2005) revealed that betaine supplementation in diet (0.072-0.144% DM) caused an

increase in serum total protein and globulin but cholesterol level was not affected by betaine supplementation which results are in agreement with our results. Zhan *et al.* (2006) found that dietary betaine addition decreased serum uric acid and serum triglycerides content in broilers. In their experiment free fatty acids concentration of serum and Hormone-Sensitive Lipase (HSL) activity in abdominal fat increased. HSL is an enzyme that initials the catabolism of TG in adipocytes (Mersmann, 1998), therefore may it be the reason decrease of triglyceride. Zou and Lu (2002) showed that supplementation of betaine (600 mg kg<sup>-1</sup>) increased serum glucose in laying hens. According to the results of present study, it seems that using different levels of betafin was not able to cause beneficial changes on lipid metabolism.

**CONCLUSION**

The results suggested that addition of betafin to broiler diets had no effect on blood parameters of broilers and betafin can replace as part of methionine. There are some factors which cause to differences among reports of experiment that use betafin, including the basal diets, genetic stocks used and environmental conditions of the experiments. Therefore, further experiments should need to be conducted to determine the effect of betafin at different condition in broilers.

**REFERENCES**

Attia, Y.A., R.A. Hassan, M.H. Shehatta and S.B. Abd-El-Hady, 2005. Growth, carcass quality and serum constituents of slow growing chicks as affected by betaine addition to diets containing 2. Different levels of methionine. *Int. J. Poult. Sci.*, 4: 856-865.  
 Esteve-Garcia, E. and S. Mack, 2000. The effect of DL-methionine and betaine on growth performance and carcass characteristics in broilers. *Anim. Feed Sci. Technol.*, 87: 85-93.  
 Fernandez, C., L. Gallego and C.J. Lopez-Bote, 1998. Effect of betaine on fat content in growing lambs. *Anim. Feed Sci. Technol.*, 73: 329-338.

- Friedewald, W.T., R.I. Levy and D.S. Fredrickson, 1972. Estimation of the concentration of LDL cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin. Chem.*, 18: 499-502.
- Gordon, T., W.P. Castelli, M.C. Hjortland, W.B. Kannel and T.R. Dawber, 1977. High density lipoprotein as a protective factor against coronary heart disease: The framingham study. *Am. J. Med.*, 62: 707-714.
- Gowenlock, A.H., J.R. McMurray and D.M. McLauchlan, 1988. Varley's Practical Clinical Biochemistry. 6th Edn., CAS Publishers and Distributors, New Delhi, pp: 477-549.
- Hassan, R.A., Y.A. Attia and E.H. El-Ganzory, 2005. Growth, carcass quality and serum constituents of slow growing chicks as affected by betaine addition to diets containing 1. Different levels of choline. *Int. J. Poult. Sci.*, 4: 840-850.
- Kermanshahi, H., 2001. Betaine replacement for DL-Methionine in the performance and carcass characteristics of broiler chicks. *J. Agric. Sci. Technol.*, 3: 273-279.
- Kim, S.K., K.H. Choi and Y.C. Kim, 2003. Effect of acute betaine administration on hepatic metabolism of S-amino acids in rats and mice. *Biochem. Pharmacol.*, 65: 1565-1574.
- Konca, Y., F. Kirkpınar, S. Mert and E. Yaylak, 2008. Effects of betaine performance, carcass, bone and blood characteristics of broilers during natural summer temperatures. *J. Anim. Vet. Adv.*, 7: 930-937.
- Mersmann, H.J., 1998. Lipoprotein and hormone-sensitive lipases in porcine adipose tissue. *J. Anim. Sci.*, 76: 1396-1404.
- NRC, 1994. Nutrient Requirements of Poultry. 9th Edn., National Academy Press, Washington, DC. USA., ISBN-13: 978-0-309-04892-7.
- SAS Institute Inc., 1998. SAS/STAT User's Guide. SAS Institute Inc., Cary, North Carolina.
- Schutte, J.B., J. de Jong, W. Smink and M. Pack, 1997. Replacement value of betaine for DL-methionine in male broiler chicks. *Poult. Sci.*, 76: 321-325.
- Sun, H., W.R. Yang, Z.B. Yang, Y. Wang, S.Z. Jiang and G.G. Zhang, 2008. Effects of betaine supplementation to methionine deficient diet on growth performance and carcass characteristics of broilers. *Am. J. Anim. Vet. Sci.*, 3: 78-84.
- Verreschi, D.C., R.H.S. Souza, P.V.R. Logato, K.K. Iseki and R.T. Freitas, 2002. Effects of dietary betaine on lipid metabolism of the pacu, *Piaractus mesopotamicus*, Holmberg (1817) in laboratory conditions. *Comp. Biochem. Physiol.*, 126: 152-152.
- Yancey, P.H., M.E. Clark, S.C. Hand, R.D. Bowlus and G.N. Somero, 1982. Living with water stress: Evolution of osmolyte systems. *Science*, 217: 1214-1222.
- Zhan, X.A., J.X. Li, Z.R. Xu and R.Q. Zhao, 2006. Effects of methionine and betaine supplementation on growth performance, carcass composition and metabolism of lipids in male broilers. *Br. Poult. Sci.*, 47: 576-580.
- Zou, X.T. and J.J. Lu, 2002. Effects of betaine on the regulation of the lipid metabolism in laying hen. *Agric. Sci., China*, 1: 1043-1049.
- Zulkifli, I., S.A. Mysahra and L.Z. Jin, 2004. Effect of betaine on growth performance and carcass characteristics in growing pigs. *Asian-Aust. J. Anim. Sci.*, 17: 1700-1704.