

The Effects of Vitamin E and L-Arginine Supplementation on Antioxidant Status and Biochemical Indices of Broiler Chickens with Pulmonary Hypertension Syndrome

¹Mokhtar Fathi, ¹Kambiz Nazer Adl, ¹Yahya Ebrahim Nezhad,
¹Habib Aghdam Shahryar, ²Mohsen Daneshyar and ¹Taimor Tanha
¹Department of Animal Science, Shabestar Islamic Azad University,
Shabestar Branch, Shabestar, Iran
²Department of Animal Science, Urmia University, Urmia, Iran

Abstract: An experiment with 320, 1 day male broilers (Ross 308) was performed to evaluate the effects of Arginine (Arg) and Vitamin E (VE) supplementation on antioxidant status, some biochemical parameters and ascites mortality in broiler chickens with cold temperature induced Pulmonary Hypertension Syndrome (PHS). The chickens were distributed between 16 pens of 20 each and each 4 allocated to each of experimental treatments. Treatment 1 was the birds fed a commercial corn-soybean meal based diet (control), Treatment 2; the birds fed the control diet supplemented with 400 IU VE kg⁻¹ of feed (VE), Treatment 3; the birds fed the control diet supplemented with 0.3% Arg in drinking water and Treatment 4; the birds fed the control diet supplemented with 400 IU VE and 0.3% Arg (Arg-VE). At day 14, temperature was reduced to amplify the incidence of pulmonary hypertension. Hematological and oxidants status parameters (in serum and liver) were determined at day 21 and 42 of age. Right/total Ventricular weight ratio (RV/TV) was recorded at the end of the experiment (week 6). The results of experiment showed that Arg birds at day 42 had the lower ($p \leq 0.05$). RV/TV ratio and ascites mortality yet had improved growth performance parameters. Red Blood Cell (RBC), Hematocrit (Hct) and Hemoglobin (Hgb) was significantly lower in Arg and Arg-VE birds as compared to other birds at day 42 of age. Plasma Malondialdehyde (MDA) was significantly lower in Arg and Arg-VE birds as compared to other birds at day 42. Plasma Glutathione Peroxidase (GPX) activity was greater ($p \leq 0.05$) in both the Arg and Arg-VE birds as compared to other groups at day 42. Plasma and liver Superoxide Dismutase (SOD) activity was not affected by none of the Arg or VE supplementation. In summary, level of Arg but not for VE used in this experiment, synchronic, some change effects on MDA and TAS in plasma or hematological parameters had positive effects on cardio pulmonary performance, growth performance and ascites mortality in broiler encounter with low temperature and PHS.

Key words: Ascites, oxidative stress, vitamin E, arginine, broilers, PHS, Iran

INTRODUCTION

Pulmonary Hypertension Syndrome (PHS) is a common disorder in broiler chickens that causes considerable mortality (Julian, 1993). PHS can be initiated by an increase in metabolic rate which implies higher oxygen demands (Lorenzoni and Ruiz-Feria, 2006). In gas exchange system working close to its physiological limit, this elevation in metabolic demand can lead to cellular hypoxia (Scheele *et al.*, 1991). To avoid cellular hypoxia, an increased cardiac output must be propelled through the noncompliant pulmonary vasculature which requires increases in Pulmonary Arterial Pressure (PAP) and Right Ventricle (RV) work (Wideman Jr., 1988). Broilers with PHS die due to right-sided congestive heart failure which is

associated with central venous congestion, pressure-induced liver cirrhosis and transudation of ascetic fluid in to the abdominal cavity (Julian, 1993). By reducing the pulmonary vascular resistance, it is possible to reduce the PAP needed to propel the cardiac output required to match broilers' metabolic demands (Wideman Jr. and Kirby, 1995). Nitric Oxide (NO) is a potent endogenous pulmonary vasodilator that increases the cyclic Guanosine Monophosphate (cGMP) level and thereby reduces the intracellular Ca⁺⁺ levels in vascular smooth muscle cells (Forstermann *et al.*, 1986). Nitric oxide is produced in the pulmonary endothelium where the NO Synthase (NOS) enzyme converts L-arginine to L-citrulline (Jorens *et al.*, 1993; McQueston *et al.*, 1993). Arginine is an essential amino acid for avian species

because birds lack the enzyme carbamyl phosphate synthetase which aids in the conversion of ornithine to citrulline and thus Arg (Tamil and Ratner, 1963). It has been suggested that Arg levels normally fed to broiler chickens (NRC, 1994) are adequate for growth but may be inadequate to fully support the production of NO by avian macrophages and the pulmonary vascular endothelium (Taylor *et al.*, 1992; Dietert *et al.*, 1994; Wideman Jr. and Kirby, 1995; Wideman *et al.*, 1996; Martinez-Lemus *et al.*, 1999).

Vitamin E (VE) is known to be a powerful lipid-soluble antioxidant that scavenges lipid radicals. It has the ability to react with fatty acid peroxy radicals, the primary products of lipid peroxidation and intercepts the chain reaction preventing further radical reactions. During the antioxidant reaction, α -tocopherol is turned into a stable radical (Burton and Ingold, 1981). Vitamin E could thus help in reducing the oxidative stress in lung vessels and consequently reduce the endothelial damage. It has been reported that birds developing ascites have low levels of α -tocopherol in lung and liver, providing evidence that a compromised antioxidant status is involved in the etiology of PHS (Enkvetchakul *et al.*, 1993). Objective of this research was conducted to investigate the effects of an vasodilator source (Arg) and an antioxidant (VE) and combination on ascites disorder in broiler chickens.

MATERIALS AND METHODS

Birds and diets: A 320, 1 day old male broiler chickens (Ross 308) were used in this research. Chicks were distributed between 16 cages (replicates) randomly on arrival and each 4 were allocated to one of the treatments. Four groups including, control group (control, basal diet); vitamin E (DL α -tocopherol acetate) with 400 IU vitamin E kg^{-1} fed (HVE); arginine with 0.3% L -Arg (monohydrochloride Canada-Iran) in water (HArg) and containing both compound at same levels from HVE and Harg (HVE-HArg). All chickens were fed a basal corn-soybean diet to meet the requirement including 22.04% CP and 3,200 kcal kg^{-1} of ME (1-21 days) or 20.26% CP and 3,200 kcal ME (22-42 days). The basal diet includes Arg at 1.3% and VE at 40 IU kg^{-1} of feed for both growing periods. From day 1-14, all chicks were fed the basal diet. From day 15, the chicks received the treatment. Feed and water provided for *ad libitum* consumption.

Management and measurements: Birds reared at 32 and 30°C during week 1 and 2, respectively. For inducing ascites, the house temperature was decreased to 15°C during week 3 and temperature was maintained between 10 and 15 for the rest of the study (Iqbal *et al.*, 2001).

Average body weight gain, average feed intake and average feed conversion ratio were measured and calculated weekly from week 3 and for all experimental period. Mortality was recorded daily and all of the dead birds inspected for diagnosis of ascites. The mortality of ascites was diagnosed by the symptoms of right ventricle hypertrophy, cardiac muscle laxation, swollen and stiff liver and occurrence of clear, yellowish and colloidal fluid in the abdominal cavity (Geng *et al.*, 2004).

On days 21 and 42 of age, one chick per replicate was randomly selected after a 3 h starvation and used for blood sampling from wing vein. After taking blood samples, the birds were killed and thorax and abdomen were opened and heart was removed and used for determination of RV/TV ratio (Daneshyar *et al.*, 2009). Liver tissue for antioxidant enzymatic evaluation and inspected for signs of heart failure and ascites. Bird having RV/TV ratios $>0.29\%$ were considered to have ventricular hypertrophy (Julian, 1987). Blood samples were collected in anticoagulant (EDTA) tubes. Part of each blood sample (4 mL) immediately used for measuring total Red Blood Cell (RBC) count, Hematocrit (Hct) and Hemoglobin (Hgb). The remainder (6 mL) was centrifuged and plasma was collected and stored at -80°C until measurement of the other enzymatic and chemical analysis.

Antioxidant indices: Antioxidant enzyme activities were determined in whole blood and liver tissues. Enzymes activities (GPX, SOD) were spectrophotometrically using RANSEL glutathione peroxidase, RANSEL superoxide dismutase and RANDOX total antioxidant status kit (RANDOX Laboratories Ltd., Ardmore, Diamond Road, Crumlin, United Kingdom, BT29 4QY). Glutathione Peroxidase (GPX) activity was measured using the method described by Paglia and Valentine (1967) in which GPX activity is coupled to the oxidation of Nicotinamide Adenine Dinucleotide Phosphate (NADPH) by glutathione reductase. The oxidation of NADPH was determined at 378°C by measuring the absorbance at 340 nm. The reaction mixture consisted of 50 mmol potassium phosphate buffer (pH 7), 1 mmol EDTA, 1 mmol NaN_3 , 0.2 mmol NADPH, 1 mmol glutathione and 1 U mL^{-1} glutathione reductase. Absorbance at 340 nm was recorded for 5 min. The activity was calculated as the slope of the lines as mmol NADPH oxidized/min GPx activity was expressed as U g^{-1} protein.

Plasma Superoxide Dismutase (SOD) activity was determined by Xanthine Oxidase (XOD) enzyme. This method employs Xanthine and Xanthine Oxidase (XOD) to generate superoxide radicals which react with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyltetrazolium

chloride (INT) to form a red formazan dye. The superoxide dismutase activity is then measured by the degree of inhibition of this reaction. One unit of SOD is that which causes a 50% inhibition of the rate of reduction of INT under the conditions of the assay. Centrifuge 0.5 mL of whole blood for 10 min at 3000 rpm and then aspirate off the plasma. Then wash erythrocytes 4 times with 3 mL of 0.9% NaCl solution centrifuging for 10 min at 3000 rpm after each wash. The washed centrifuged erythrocytes should then be made up to 2.0 mL with cold redistilled water, mixed and left to stand at +4°C for 15 min. The lysate is diluted with 0.01 mol L⁻¹ phosphate buffer pH 7.0, so that the percentage of inhibition falls between 30 and 60%. Then, the lysate incubated for 5 min at 25°C with CAPS 40 mmol L⁻¹, pH 10.2, EDTA 0.94 mmol L⁻¹, Xanthine 0.05 mmol L⁻¹, INT 0.025 mmol L⁻¹, Xanthine oxidase 80 U L⁻¹. The results were express as units of activity (U g⁻¹ Hemoglobin) (RANDOX RANSOD Superoxide dismutase MANUAL). Liver Superoxide Dismutase (SOD) activity in liver was determined following the Xanthine Oxidase Method described by McCord and Fridovich (1969) with modifications. For plasma Total Antioxidant (TAS) capacity, ABTS (2, 2'-Azino-di-[3-ethylbenzthiazoline sulphonate]) is incubated with a peroxidase (metmyoglobin) and H₂O₂ to produce the radical cation ABTS^{•+}. This has a relatively stable blue-green colour which is measured at 600 nm. Antioxidants in the added sample cause suppression of this colour production to a degree which is proportional to their concentration.

For evaluation serum (20 µL) was incubated at 37°C with 1 mL metmyoglobin chromogen (6.1 mM L⁻¹), ABTS 610 µmol L⁻¹ then detected absorbance in 600 nm. Then, this component incorporate with 250 µmol L⁻¹ hydrogen peroxide (25% µmol L⁻¹) and after 3 sec detected aberrance in 600 nm. Reduce amount in catione ABTS^{•+} result in antioxidant would compared with standard. The results were expressed as units of activity (mmol L⁻¹).

MDA determination: The blood was centrifuged at 1,500×g for 5 min; plasma was collected in labeled tubes and stored at -80°C until analysis. After thawing, 500 µL of plasma was placed in a labeled glass tube and mixed with the reagents of a commercial kit for the measurement of Thiobarbituric Acid Reactive Substances (TBARS) and each tube was covered with a glass marble and incubated at 95°C for 45 min.

The tubes were removed from incubation and allowed to cool in an ice bath for 10 min. Once cooled, the tubes were centrifuged at 3,000×g for 10 min and the supernatant carefully removed from the tubes for analysis. The absorbance of the supernatants was measured at

532 nm using a UV/VIS spectrophotometer (Gildford Instrument Laboratories, Inc., Oberlin, OH) and the results were compared against a standard curve made with 100, 50, 25, 12.5 and 0 nmol mL⁻¹ of malondialdehyde dimethyl acetyl.

Statistical analysis: The data were analyzed based on a completely randomized design using the GLM procedure of SAS 9.1 Institute (SAS, 1996). Duncan's multiple range were used to separate the means when treatments means were significant (p≤0.05), thus a probability level of p≤0.05 was considered statistically significant. Data were presented as means±SEM.

RESULTS AND DISCUSSION

As showed in Table 1, Arg birds had lower RV/TV ratio and mortality. Arg supplementation causes the pulmonary arterial relaxation which reflect the improved NO synthesis because of higher substrate availability (Taylor *et al.*, 1992; Wideman Jr. and Kirby, 1995; Wideman *et al.*, 1996; Lorenzoni and Ruiz-Feria, 2006). Furthermore, it has been reported that additional availability of Arg may support the NO production and consequently, higher vasodilator capacity. Moreover, lower blood Hct was observed for Arg birds. Higher blood hematocrit is associated with sustained hypoxia (Yersin *et al.*, 1992) and correlated with ascites susceptibility (Wideman *et al.*, 1998). Decreased blood Hct by Arg supplementation results in lower blood viscosity and reduced pulmonary vascular resistance afterwards and as a result reduced Pulmonary Arterial Pressure (PAP). Since, increases in PAP represent a key step in the pathogenesis of ascites (Wideman Jr., 1988), improvements in pulmonary vasorelaxation and reducing in blood viscosity (via reducing pulmonary vascular resistance) should help reduce the mortality caused by

Table 1: RV/TV ratio and ascites mortality of broilers consuming a com-soybean meal based diet and different levels of Arginine (Arg) and Vitamine E (VE)

Group ¹	Determination time	RV/TV index (%)	Ascites mortality (%)
Control HVE	Day 42	0.320 ^a	38 ^a
HArg HVE-HArg		0.300 ^a	28 ^a
		0.260 ^b	22 ^b
		0.290 ^{ab}	26 ^{ab}
	±SEM	0.011	3
	p-value	0.037	0.025

¹Control group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹; HVE group: Basal diet plus 400 IU of VE supplementation per kg of feed; HArg group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹ and supplementation Arg (0.3% wt./vol.) in the drinking water; HVE-HArg: Basal diet supplementation with Arg (0.3% wt./vol.) in the drinking water plus 400 IU of VE supplementation per kg of feed. *Means within rows with different superscript letters are significantly different (p<0.05)

ascites (Wideman Jr. and Kirby, 1995), subsequently induced improving growth performance. Moreover, Wideman Jr. and Kirby (1995) reported that birds fed diets supplemented with 1.5% L-Arg HCl had significantly lower right; total ventricle weight ratios than control birds, presumably reflecting a reduction in pulmonary arterial pressure in Arg-supplemented birds. Later, it was reported that supplemental dietary Arg permitted broilers to exhibit flow-dependent pulmonary vasodilatation when a pulmonary artery snare was tightened to force the entire cardiac output through one lung. The reduced pulmonary vascular resistance allowed Arg-supplemented broilers to maintain a low pulmonary arterial pressure at high pulmonary blood flow rates (Wideman *et al.*, 1996) which resembles the inherent capability of jungle fowl that are known to be highly resistant to PHS (Wideman *et al.*, 1998; Kochera-Kirby *et al.*, 1999). The reducing in PAP, pulmonary vascular resistance and cardiac output cause decrease hypertrophy and subsequently reducing the RV/TV. The L-Arg-NO mechanism also facilitates vasorelaxation of pulmonary artery rings isolated from domestic fowl (Martinez-Lemus *et al.*, 1999).

The efficacy of supplemental Arg for reducing the incidence of PHS in field trials has been highly variable and dose response influences of Arg on PHS have not been demonstrated (Wideman Jr. and Kirby, 1995). The reasons for this variability are not known but it is possible that under some circumstances supplemental dietary Arg is catabolized by kidney arginase instead of providing additional substrate for NO synthesis, especially when increased kidney arginase activity is induced by diets high in Lys or CP (O'Dell and Savage, 1966; Stutz *et al.*, 1972). Arginase converts Arg into ornithine and urea. It has been reported that between 40-60% of urea excreted by birds is from Arg metabolism and as birds cannot synthesize ornithine almost all of the plasma ornithine in the birds is also derived from Arg metabolism (Nesheim, 1968; Austic and Nesheim, 1970; Stutz *et al.*,

1972; Chu and Nesheim, 1979). Urea production increases with high levels of Arg and an excess of urea reportedly inhibited the action of nitric oxide synthase in a macrophage cell line (Prabhakar *et al.*, 1997) (Table 2). Data from antioxidants enzymes activities, TAS and MDA in plasma is shown in Table 3. There is no significant difference in these parameters at day 21 in plasma. But at day 42, the GPX activity in HArg and HArg-HVE groups was greater than other groups, contemporary HArg and HArg-HVE groups had lower TAS and lower MDA value than the control and HVE. There are no significant differences in SOD activation among groups. Endothelial cells synthesize nitric oxide from L Arginine in a reaction catalyzed by the enzyme Nitric Oxide Synthase (NOS). During NO production by NOS, produce amount of superoxide and hydrogen peroxide. So, the antioxidant system should neutralize these free radicals, hence the lower TAS in Harg and Harg-HVE groups can be

Table 2: Red Blood Cell (RBC), Hemoglobin (Hgb) and Hematocrit (Hct) of broilers consuming a corn-soybean meal based diet and different levels of arginine (Arg) and Vitamine E (VE)

Group ¹	Determination time	RBC (10 ⁶ dL ⁻¹)	Hgb (g dL ⁻¹)	Hct (%)
Control	Day 21	2.420	8.570	34.27
		2.390	8.170	33.55
		2.060	8.120	32.46
HVE		2.280	7.350	29.55
		0.150	0.450	1.950
		0.090	0.080	0.100
HArg	±SEM	0.150	0.450	1.950
		0.090	0.080	0.100
		0.090	0.080	0.100
HVE-HArg	Day 42	2.800 ^b	11.20 ^a	39.30 ^{ab}
		3.290 ^a	12.15 ^a	45.62 ^a
		2.431 ^b	8.520 ^b	33.67 ^b
	±SEM	2.400 ^b	8.270 ^b	33.52 ^b
		0.180	0.480	1.500
		0.041	0.035	0.025
	p-value			

¹Control group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹; HVE group: Basal diet plus 400 IU of VE supplementation per kg of feed; HArg group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹ and supplementation Arg (0.3% wt./vol.) in the drinking water; HVE-HArg: Basal diet supplementation with Arg (0.3% wt./vol.) in the drinking water plus 400 IU of VE supplementation per kg of feed. *Means within rows with different superscript letters are significantly different (p<0.05)

Table 3: Plasma Malondialdehyde (MDA), Total Antioxidant capacity (TAS) and Glutathione Peroxidase (GPX) and Superoxide Desmutase (SOD) activities of broilers consuming a corn-soybean meal based diet and different levels of Arginine (Arg) and Vitamine E (VE)

Group ¹	Determination time	MDA (nmol mL ⁻¹)	GPX activity (U mg ⁻¹ protein)	SOD activity (U mg ⁻¹ protein)	TAS (mmol L ⁻¹)
Control	Day 21	2.500	39.320	1371.000	0.710
		3.600	40.770	1800.000	0.740
		3.100	43.050	1527.000	0.970
HVE		3.600	42.860	1309.000	0.830
		0.260	2.500	195.000	0.110
		0.091	0.087	0.071	0.068
HArg	±SEM	0.260	2.500	195.000	0.110
		0.091	0.087	0.071	0.068
		0.091	0.087	0.071	0.068
HVE-HArg	Day 42	6.270 ^a	30.720 ^b	913.000	1.360 ^a
		6.770 ^a	33.650 ^b	906.000	1.300 ^{ab}
		2.270 ^b	43.670 ^a	882.000	0.940 ^b
	±SEM	3.120 ^b	42.350 ^a	1061.000	1.010 ^b
		0.400	1.500	125.000	0.150
		0.041	0.034	0.065	0.027
	p-value				

¹Control group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹; HVE group: Basal diet plus 400 IU of VE supplementation per kg of feed; HArg group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹ and supplementation Arg (0.3% wt./vol.) in the drinking water; HVE-HArg: Basal diet supplementation with Arg (0.3% wt./vol.) in the drinking water plus 400 IU of VE supplementation per kg of feed. *Means within rows with different superscript letters are significantly different (p<0.05)

Table 4: Malondialdehyde (MDA) and Glutathione Peroxidase (GPX) and Superoxide Desmotase (SOD) activities in liver tissue of broilers consuming a corn-soybean meal based diet and different levels of Arginine (Arg) and Vitamine E (VE)

Group ¹	Determination time	MDA (nmol mL ⁻¹)	GPX (U mg ⁻¹ protein)	SOD (Umg ⁻¹ protein)
Control	Day 21	1.320	0.250	8.600
HVE		1.970	0.240	7.550
		1.450	0.250	7.750
		1.970	0.260	8.250
HArg	±SEM	0.190	0.011	0.490
HVE-HArg	p-value	0.078	0.091	0.082
	Day 42	2.600	0.250	8.250
		1.950	0.290	9.450
		2.100	0.290	9.100
		2.420	0.300	9.070
	±SEM	0.300	0.020	0.510
	p-value	0.072	0.067	0.082

¹Control group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹; HVE group: Basal diet plus 400 IU of VE supplementation per kg of feed; HArg group: Basal diet with 1.2% Arg and 40 IU of VE kg⁻¹ and supplementation Arg (0.3% wt./vol.) in the drinking water; HVE-HArg: Basal diet supplementation with Arg (0.3% wt./vol) in the drinking water plus 400 IU of VE supplementation per kg of feed. *Means within rows with different superscript letters are significantly different (p<0.05)

justified this lowering. Beside, Arg induce increase GPX activity and amplification of antioxidant system. So, its result reducing in lipid and protein per oxidation and reducing MDA. Contents of Table 4 showing that there are no significant differences in MDA value, GPX and SOD activities in liver among treatment groups at both periods 21 and 42. It seeming that hematological and plasma antioxidant actives changes in response to systemic hypoxemia and oxidative stress may occur more rapidly than liver tissue.

The lack of consistency in the efficacy of supplemental Arg on reducing the incidence of PHS may be explained by the multiple independent components that affect PHS susceptibility in addition to the maintenance of an inappropriately elevated vascular tone. These factors include allow anatomical pulmonary vascular capacity, inadequate surface area available for gas exchange and a cardiac output that steadily increases in support of metabolic requirements (Wideman Jr. and Kirby, 1995). Additionally, kidney arginase activity may differ among strains of birds or among individual birds within a population (Nesheim, 1968; Chu and Nesheim, 1979). If so, birds with high kidney arginase activity (high rate of Arg degradation) may benefit from supplemental Arg whereas birds with a low kidney arginase activity may not respond to increased levels of Arg. The production of NO is modulated and affected by a wide variety of factors, making it difficult to achieve the predictable results solely by increasing substrate availability *in vivo*.

These results supported hypothesis that supplemental Arg improve problems associated with pulmonary hypertension by improve NO synthesis and

improving in pulmonary arterial relaxation (Taylor *et al.*, 1992; Wideman Jr. and Kirby, 1995; Wideman *et al.*, 1996). On the other hand, we found that VE supplementation did not improve or augment the effects of Arg on hematological, antioxidant enzymes activities, growth performance and problems associated with ascites. It appears that VE supplementation at the levels used in this research negated the ability of Arg to improve pulmonary hypertension under cold stress. It was anticipated that vitamin E might improve vasodilation by protecting the endothelium from free radical damage and a healthier endothelium might have more functional cells with higher constitutive levels of NOS which could contribute to a higher NO production. We further anticipated that vitamin E might improve vasodilation by protecting NO from direct free radical attack increasing its life span and avoiding the formation of peroxynitrite (Beckman and Koppenol, 1996). The levels of MDA in blood were measured as an indicator of lipid peroxidation and oxidative exposure to low temperature. Levels of MDA did differ among treatments at day 42 (Table 2). However, there was a tendency for the birds in the control and HVE groups to have higher MDA values than birds in the other groups (p = 0.05). It has been documented that VE may become prooxidative agent under some circumstances. For instance, vitamin E can react with organic peroxides interrupting the chain reaction of lipid peroxidation. This reaction involves the formation of tocopheroxyl radicals which are regenerated by means of hydrogen donors. If this reduction is incomplete, the tocopheroxyl radicals can initiate oxidative processes (Mukay, 1993; Schneider, 2005).

CONCLUSION

These results agree with recent epidemiological studies in humans in which the use of VE in patients to prevent cardiovascular diseases has had null or even negative effects in high-risk patients (Kris-Etherton *et al.*, 2004). There is evidence of chemical differences between the isomers of VE that could be important to consider for future research. Although, γ -tocopherol is considered a less potent chain-breaking antioxidant compared with α -tocopherol, its molecular arrangement appears to make it better to trap lipophilic electrophiles such as reactive nitrogen oxide species (Kamal-Eldin and Appelqvist, 1996). This feature may be important to rapidly decrease molecules like peroxynitrite, a potent oxidant believed to cause direct tissue injury (Beckman and Koppenol, 1996). Furthermore, γ -tocopherol has antiinflammatory properties and a less hydrophobic condition than α -tocopherol (Jiang *et al.*, 2000, 2001).

REFERENCES

- Austic, R.E. and C. Nesheim, 1970. Role of kidney arginase in variations of the arginine requirements of chicks. *J. Nutr.*, 100: 855-867.
- Beckman, J.S. and W.H. Koppenol, 1996. Nitric oxide, superoxide and peroxynitrite: The good, the bad and ugly. *Am. J. Physiol. Cell. Physiol.*, 271: C1424-C1437.
- Burton, G.W. and K.U. Ingold, 1981. Autoxidation of biological molecules. 1. Antioxidant activity of vitamin E and related chain-breaking phenolic antioxidants *in vitro*. *J. Am. Chem. Soc.*, 103: 6472-6477.
- Chu, S.H.W. and M.C. Nesheim, 1979. The relationship of plasma arginine and kidney arginase activity to Arginine degradation in chickens. *J. Nutr.*, 109: 1752-1758.
- Daneshyar, M., H. Kermanshahi and A. Golian, 2009. Changes of biochemical parameters and enzyme activities in broiler chickens with cold-induced ascites. *Poult. Sci.*, 88: 106-110.
- Dietert, R.R., K.A. Golemboski and R.E. Austic, 1994. Environment-immune interactions. *Poult. Sci.*, 73: 1062-1076.
- Enkvetchakul, B., W. Bottje, N. Anthony, R. Moore and W. Huff, 1993. Compromised antioxidant status associated with ascites in broilers. *Poult. Sci.*, 72: 2272-2280.
- Forstermann, U., A. Mulsch, E. Bohme and R. Russe, 1986. Stimulation of soluble guanylate cyclase by an acetylcholine-induced endothelium-derived factor from rabbit and canine arteries. *Circ. Res.*, 58: 531-538.
- Geng, A.L., Y.M. Guo and Y. Yang, 2004. Reduction of ascites mortality in broilers by coenzyme Q10. *Poult. Sci.*, 83: 1587-1593.
- Iqbal, M., D. Cawthon, R.F. Wideman Jr. and W.G. Bottje, 2001. Lung mitochondrial dysfunction in pulmonary hypertension syndrome. I. Site-specific defects in the electron transport chain. *Poult. Sci.*, 80: 485-495.
- Jiang, Q., I. Elson-Schwab, C. Courtemanche and B.N. Ames, 2000. Gamma-tocopherol and its major metabolite, in contrast to γ -tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc. Nat. Acad. Sci. USA.*, 97: 11494-11499.
- Jiang, Q., S. Christen, M.K. Shigenaga and B.N. Ames, 2001. γ -tocopherol, the major form of vitamin E in the US diet, deserves more attention. *Am. J. Clin. Nutr.*, 74: 714-722.
- Jorens, P.G., P.A. Vermeire and A.G. Herman, 1993. L-arginine-dependent nitric oxide synthase: A new metabolic pathway in the lung and airways. *Eur. Respir. J.*, 6: 258-266.
- Julian, R.J., 1987. The effect of increased sodium in the drinking water on right ventricle hypertrophy, right ventricle failure and ascites in broiler chickens. *Avian Pathol.*, 16: 61-71.
- Julian, R.J., 1993. Ascites in poultry. *Avian Pathol.*, 22: 419-454.
- Kamal-Eldin, A. and L.A. Appelqvist, 1996. The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids*, 31: 671-701.
- Kochera-Kirby, Y., N.B. Anthony, J.D. Hughes Jr., R.W. McNew, J.D. Kirby and R.F. Wideman Jr., 1999. Electrocardiographic and genetic evaluation of giant jungle fowl, broilers and their reciprocal crosses following unilateral bronchus occlusion. *Poult. Sci.*, 78: 125-134.
- Kris-Etherton, P.M., A.H. Lichtenstein, B.V. Howard, D. Steinberg and J.L. Witztum, 2004. Antioxidant vitamin supplements and cardiovascular disease. *Circulation*, 110: 637-641.
- Lorenzoni, A.G. and C.A. Ruiz-Feria, 2006. Effects of vitamin E and L-arginine on cardiopulmonary function and ascites parameters in broiler chickens reared under subnormal temperatures. *Poult. Sci.*, 85: 2241-2250.
- Martinez-Lemus, L.A., R.K. Hester, E.J. Becker, J.S. Jeffrey and T.W. Odom, 1999. Pulmonary artery endothelium-dependent vasodilation is impaired in a chicken model of pulmonary hypertension. *Am. J. Physiol.*, 277: R190-R197.
- McCord, J.M. and I. Fridovich, 1969. Superoxide dismutase: An enzymic function for erythrocyte (hemocuprein). *J. Biol. Chem.*, 244: 6049-6055.
- McQueston, J.A., D.N. Cornfield, I.F. McMurtry and S.H. Abman, 1993. Effects of oxygen and exogenous L-arginine on EDRF activity in fetal pulmonary circulation. *Am. J. Physiol.*, 264: H865-H871.
- Mukay, K., 1993. Synthesis and Kinetic Study of Antioxidant and Prooxidant Actions of Vitamin E Derivates. In: *Vitamin E in Health and Disease*, Packer, L. and J. Fuchs (Eds.). Marcel Dekker, New York, pp: 97-119.
- NRC, 1994. *Nutrient Requirements of Poultry*. 9th Rev. Edn., National Academy Press, Washington, DC., USA.
- Nesheim, M.C., 1968. Kidney arginase activity and lysine tolerance in strains of chickens selected for a high or low requirement for arginine. *J. Nutr.*, 95: 79-87.
- O'Dell, B.L. and J.E. Savage, 1966. Arginine-lysine antagonism in the chick and its relationship to dietary cations. *J. Nutr.*, 90: 364-370.
- Paglia, D.E. and W.N. Valentine, 1967. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J. Lab. Clin. Med.*, 2: 158-169.

- Prabhakar, S.S., G.A. Zeballos, M. Montoya-Zavala and C. Leonard, 1997. Urea inhibits inducible nitric oxide synthase in macrophage cell line. *Am. J. Physiol.*, 273: C1882-C1888.
- SAS, 1996. SAS User's Guide. Version 6.12, SAS Institute Inc., Cary, NC, USA.
- Scheele, C.W., W. De Wit, M.T. Frankenhuis and P.F. Vereijken, 1991. Ascites in broilers. 1. Experimental factors evoking symptoms related to ascites. *Poult. Sci.*, 70: 1069-1083.
- Schneider, C., 2005. Chemistry and biology of vitamin E. *Mol. Nutr. Food Res.*, 49: 7-30.
- Stutz, M.W., J.E. Savage and B.L. O'Dell, 1972. Cation anion balance in relation to arginine metabolism in the chick. *J. Nutr.*, 102: 449-457.
- Tamil, H. and S. Ratner, 1963. A study of ornithine, citrulline and arginine synthesis in growing chick. *Arch. Biochem. Biophys.*, 102: 259-269.
- Taylor, R.L., R.E. Austic and R. Dietert, 1992. Dietary arginine influences rous sarcoma growth in a major histocompatibility B complex progressor genotype. *Proc. Soc. Exp. Biol. Med.*, 199: 38-41.
- Wideman, Jr. R.F., 1988. Ascites in poultry. *Monsanto Nutr. Update*, 6: 1-7.
- Wideman, R.F. Jr. and Y.K. Kirby, 1995. Evidence of a ventilation-perfusion mismatch during acute unilateral pulmonary artery occlusion in broilers. *Poult. Sci.*, 74: 1209-1217.
- Wideman, R.F., T. Wing, Y.K. Kirby, M.F. Forman, N. Marson, C.D. Tacker and C.A. Ruiz-Feria, 1998. Evaluation of minimally invasive indices for predicting ascites susceptibility in three successive hatches of broilers exposed to cool temperatures. *Poult. Sci.*, 77: 1565-1573.
- Wideman, R.F., Y.K. Kirby, C.D. Tackett, N.E. Marson and R.W. McNew, 1996. Cardio-pulmonary function during acute unilateral occlusion of the pulmonary artery in broilers fed diets containing normal or high levels of arginine-HCl. *Poult. Sci.*, 75: 1587-1602.
- Yersin, A.G., W.E. Huff, L.F. Kubena, M.A. Elissalde, R.B. Harvey, D.A. Witzel and L.E. Giroir, 1992. Changes in hematological, blood gas and serum biochemical variables in broilers during exposure to simulated high altitude. *Avian Dis.*, 36: 189-196.