

## Seasonal Variation of Heavy Metals in Shrimp *Penaeus kerathurus* (Forsk., 1775) from Izmir Bay, Turkey

Gurel Turkmen

Faculty of Fisheries, Ege University, 35100 Bornova, Izmir, Turkey

**Abstract:** Seasonal changes in heavy metals (Cd, Cu, Fe, Pb, Zn) concentrations in muscle tissues of both male and female caridean shrimp (*Penaeus kerathurus*) from Izmir bay, Turkey were measured for a year period by using flame and graphite furnace AAS. The relationships in various heavy metal concentrations were compared according to sex and seasons. Heavy metal content varied with type of metals, seasons and sex. The highest concentration of heavy metal in the muscle tissue of *P. kerathurus* for Cd, Pb, Cu, Fe and Zn was registered in Summer months. Mean Fe level in male shrimps was considerably higher (30.19 compared to 20.23  $\mu\text{g g}^{-1}$  dry weight) in October. Regarding Cu, the same case could be observed (27.58  $\mu\text{g g}^{-1}$  dry weight in males compared to 19.02 in females). In the present study, heavy metals accumulated in the shrimp muscle in the order Zn>Fe>Cu>Pb>Cd. Concentrations of metals ranged within the recommended limits for human consumption and not represent a health risk.

**Key words:** Heavy metals, *Penaeus kerathurus*, seasonal changes, dry weight, Izmir bay

---

### INTRODUCTION

In the marine environment, toxic metals are potentially accumulated in sediments and marine organisms and subsequently transferred to man through the food chain. Thus, it has become increasingly important to determine and assess levels of heavy metals in marine organisms because of nutritional and safety conditions. This is true especially for edible marine organisms as they are a potential dietary source of protein (Pourang and Dennis, 2005). *Penaeus kerathurus* (Caridean shrimp) is distributed in coastal areas of the Marmara, Aegean and Mediterranean seas and it is commercially one of the most important shrimp species in coastal fisheries in Turkey. This species is fished for all along the above mentioned sea coasts because of its large size and excellent taste. It is also a target species for fishermen using trammel nets in Izmir bay in Western Turkey (Turkmen and Yilmazyerli, 2006).

Izmir bay is one of the great natural bays of the Mediterranean. The main urban settlement around this bay is the city of Izmir (88,000 ha and a population of 3,965,232 (2011 census)). The bay has been divided into three sections (outer, middle and inner) according to the physical characteristics of the different water masses (Kontas, 2012). Izmir is an important industrial and commercial centre and a cultural focal point. The streams and hundreds of small domestic discharge outlets, flow to the bay. The main industries in the region include food processing, beverage manufacturing and bottling, tanneries, oil, soap and paint production, chemical industries, paper and pulp factories, textile industries, metal processing and timber processing

(Kucuksezgin *et al.*, 2006). Since, heavy metal pollution has been increasing in the area, shrimp can accumulate heavy metals from seawater and/or sediment to concentrations that may exceed the levels of safety for consumers. Therefore, it is important to carry out a study to determine heavy metal concentrations accumulated in the shrimp *Penaeus kerathurus*. There have been a large number of detailed studies on heavy metals related to tissue, season or sexes in marine shrimps recently particularly in heavily polluted areas of Mediterranean and Marmara sea in Turkey (Canli *et al.*, 2001; Kargin *et al.*, 2001; Cogun *et al.*, 2005; Yilmaz and Yilmaz, 2007; Firat *et al.*, 2008). However, a few papers have been published to date concerning heavy metal concentrations in *P. kerathurus* in Izmir bay (Kontas, 2012) and Iskenderun bay in Turkey (Balkas *et al.*, 1982). But no data are available on heavy metals concentrations related to season or sex differences of *P. kerathurus* from Turkish sea coast. The objective of the present study was to provide information on the seasonal concentration of Cd, Cu, Fe, Pb and Zn in the edible part of shrimp *P. kerathurus* from the Izmir Bay and compare them with other species and previous studies. There is increasing awareness that trace metal concentrations in crustaceans can differ not only among different tissues but also between males and females. To the knowledge, no study to date has focused on analyzing seasonal variation of heavy metals in male and female shrimp of *Penaeus kerathurus*. Additionally, the relationships between seasonal heavy metal concentration and male and female individuals were examined. This data is important as a background for the estimation of the future impact of metal pollutants in this area.




Fig. 1: Izmir bay and location of sampling site

## MATERIALS AND METHODS

Caramote shrimps were captured from Izmir bay (Fig. 1) in Summer (July) and Autumn (October) of 2010 and Winter (January) and Spring (April) of 2011 by a commercial shrimp trammel net. Immediately after capture, the shrimp were stored on ice in an insulated box and transferred to the laboratory and frozen  $-20^{\circ}\text{C}$  until dissected. Male and female shrimps were grouped in sex and size. Ten samples from each sex were obtained for each season. Total shrimp length and weight were measured before dissection. Metal analyses were done in the Central Fishers Research Institute of Ministry of Food, Agriculture and Livestock in Trabzon. For analysis, edible part of each shrimp were dissected, weighed and dried at  $120^{\circ}\text{C}$  until they reached a constant weight. For each sample, between 0.3 and 0.5 g of tissue was weighed and placed in a Teflon digestion vessel, 7 mL of concentrated (65%) nitric acid ( $\text{HNO}_3$ ) and 1 mL of (30%) hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) added and digested in a microwave digester (ETHOS 1, Milestone Inc.). The sample in the vessel containing concentrated nitric acid was then subjected to a microwave program as follows: Step 1:  $25-200^{\circ}\text{C}$  for 10 min at 1,000 W; Step 2:  $200^{\circ}\text{C}$  for 10 min at 1,000 W (Digestion Application Note: DG-FO-17, ETHOS 1, Milestone Inc.). Digests were diluted with deionized water to 25 mL in acid washed standard flask. Analyses were performed by flame (Fe, Zn) and graphite furnace (Cd, Cu, Pb,) atomic absorption spectrophotometry (GBC 905 model). Following absorption lines were used; Fe 248.3 nm, Zn 213.9 nm, Cd 228.8 nm, Cu 324.7 nm, Pb 283.3 nm. The results were

expressed on a dry weight basis ( $\mu\text{g g}^{-1}$  dw). Replicates (five) were utilized to analysis of metals concentration. The accuracy of the determination procedure was assessed by the analysis of certified reference material (CRM, Dorm-3) (Fish protein) in triplicate for each batch of analysis. Recoveries were consistently in the range 93.1-104.3%. In order to compare with other shrimp and evaluate human health risk on a wet weight (ww) basis the moisture content ( $75.8 \pm 1.2\%$ ) of shrimp species was evaluated and the data was converted to wet weight using conversion factor of 4.13 for muscle. Differences in heavy metal levels between seasons and sexes were compared by one way ANOVA (Kruskal Wallis one way Analysis of Variance). The levels of significance were  $p < 0.05$ .

## RESULTS AND DISCUSSION

The results show that the concentrations of zinc, iron, copper, lead and cadmium in shrimp muscles were significantly affected by the seasons in Izmir bay except iron and copper for male shrimp ( $p < 0.05$ ) (Table1). The highest concentration of heavy metal in the muscle tissue of *P. kerathurus* for Cd, Pb, Cu, Fe and Zn was registered in Summer months. In shrimps, high metal concentrations were observed mainly during the Summer (Mantelatto *et al.*, 1999; Kargin *et al.*, 2001; Cogun *et al.*, 2005). In addition, Kontas (2012) found that metal concentrations in suspended particulate matter increased during Summer season in Izmir bay. In this study, Zinc, Fe and Cu were the most abundant elements in the different samples analyzed throughout the year whereas Pb and Cd were the least abundant. The order of metal

Table 1: Mean±SD of some body measurements and heavy metal concentrations ( $\mu\text{g g}^{-1}$  dw) between seasons and sexes of *P. kerathurus* from Izmir bay

TL-W/Metals	Sex	April	July	October	January
Total Length (cm)	M	14.2±1.5	14.8±1.8	13.1±1.3	13.8±1.4
	F	16.0±2.2	17.8±2.5	15.1±1.8	15.8±2.0
Weight (g)	M	20.9±4.3	22.6±4.0	19.7±4.2	20.2±3.5
	F	34.6±6.6	44.8±7.8	25.9±5.8	26.8±5.6
Cd	M	0.082±0.030 <sup>ab</sup>	0.220±0.074 <sup>ba</sup>	0.082±0.030 <sup>ab</sup>	0.020±0.006 <sup>ca</sup>
	F	0.110±0.035 <sup>ab</sup>	0.188±0.069 <sup>ba</sup>	0.102±0.025 <sup>ab</sup>	0.023±0.007 <sup>ca</sup>
Pb	M	0.42±0.15 <sup>ab</sup>	0.81±0.21 <sup>ba</sup>	0.39±0.18 <sup>ab</sup>	0.36±0.15 <sup>ab</sup>
	F	0.39±0.18 <sup>ab</sup>	0.91±0.22 <sup>ba</sup>	0.35±0.16 <sup>ab</sup>	0.42±0.19 <sup>ab</sup>
Cu	M	19.11±2.95 <sup>ab</sup>	28.47±3.22 <sup>ba</sup>	27.58±3.28 <sup>ba</sup>	17.93±1.39 <sup>ab</sup>
	F	20.90±3.55 <sup>ab</sup>	26.37±3.34 <sup>ba</sup>	19.02±3.86 <sup>ab</sup>	20.36±3.25 <sup>ab</sup>
Fe	M	20.06±6.06 <sup>ab</sup>	32.47±6.45 <sup>ba</sup>	30.19±5.62 <sup>ba</sup>	21.78±4.51 <sup>ab</sup>
	F	23.97±3.31 <sup>ab</sup>	31.26±8.89 <sup>ba</sup>	20.23±3.06 <sup>ab</sup>	21.18±5.51 <sup>ab</sup>
Zn	M	46.40±4.13 <sup>ab</sup>	63.45±6.26 <sup>ba</sup>	48.01±6.07 <sup>a</sup>	43.09±5.14 <sup>ab</sup>
	F	45.69±4.49 <sup>ab</sup>	60.81±6.41 <sup>ba</sup>	46.94±4.19 <sup>a</sup>	45.31±4.43 <sup>ab</sup>

<sup>a-c</sup>Correspond to the comparison of each group of interest along the seasons; <sup>A,B</sup>Correspond to group comparisons between sexes

concentrations in muscle tissue of both male and female was Zn>Fe>Cu>Pb>Cd. This results can be explained as a result of various biological factors and many decades of intense industrial, anthropogenic and agricultural activities carried out in this area.

Zn being an essential element for normal growth and metabolism of animals, exhibited highest accumulation in the shrimp muscle when compared with the other four metals. Zn concentrations in the muscle ranged from 38-73  $\mu\text{g g}^{-1}$  dw. Although, there were no significant differences in the zinc levels of different sexes, zinc concentrations varied significantly between seasons in the studied shrimp samples (Table 1) ( $p<0.05$ ). The high concentration of Zn is associated with the activity of enzymes played a role in the enzymatic processes of in aquatic animals (Cogun *et al.*, 2005). In general, zinc concentrations in muscle tissues were much higher than other metals in shrimps (Table 2). Iron, an oligo element plays a vital role in the enzymatic and respiratory processes of aquatic animals. The iron contents of shrimp changed from 20.06-32.4  $\mu\text{g g}^{-1}$  dw. The lowest ( $p<0.05$ ) iron content was found in the Spring and Winter. Iron concentrations did not vary significantly in muscles of both sexes except in October. Copper is an essential trace metal for animal metabolism but at high levels is a very toxic substance to aquatic life. Copper was present at relatively high concentrations in the muscle tissue with only significant differences between males and females in the Autumn (Table 1). The high levels of this element can be due in part to its high capacity of hemocyanin for binding Cu to a respiratory pigment present in large amounts in the gills of crustaceans (Paez-Osuna and Tron-Mayen, 1995). The available evidence leads us to postulate that *P. kerathurus* males and females differ in their physiological requirements for this metal. The high concentrations detected in males were probably due to the greater development of their gonads in relation to males during a phase characterized by an accelerated

metabolic rate with somatization of metal absorption. Gonads maturation in males and females of *P. kerathurus* at Izmir bay occurs during Spring and Summer with a high incidence in Summer and involves the simultaneous increase of organic constituents and ovary mass (Turkmen and Yilmazyerli, 2006). A similar pattern was described for *Pleoticus muelleri* studied in San Jorge Gulf, Argentina by Jeckel *et al.* (1996) with copper concentrations tending to increase in the male gonad in parallel to increases of these trace metals in the digestive gland indicating that their levels in the male reproductive system reflect at least partly, their levels of intake. However, Rainbow (1998) reported that decapods usually regulate their body concentrations of the essential elements Zn, Cu and Fe to approximately constant levels while they are net accumulators of the non-essential elements Cd and Pb.

Lead is a neurotoxin that causes behavioral deficits in aquatic organisms' and cause decreases in survival, growth rates and metabolism. There is often little accumulation of Pb in marine and freshwater species (Mitra *et al.*, 2012). Lead is higher concentration can occur in aquatic organisms close to anthropogenic sources. Most of the lead appears to be accumulated in the gill and liver and lower concentrations of the element were found in the muscle tissue in shrimp (Cogun *et al.*, 2005). Highest concentrations in both of male and female of lead were observed in July (Table 1). Cadmium is regarded as a priority pollutant because of its toxicity to organisms in the aquatic environment. The concentrations of Cd in *P. kerathurus* showed a wide variability, being the highest in Summer displaying a similarity between in Spring and Autumn and being the lowest in Winter season (Table 1).

Metal concentrations in *P. kerathurus* of Izmir bay were compared to previous studies performed in the same and other areas of the world (Table 2). Uysal and Tuncer and Kontas (2012) studied fish and shrimp of the Izmir

Table 2: Comparison of heavy metal concentrations in *P. kerathurus* and shrimp species (belonging to family Penaeidae) from Turkey and different areas of the world as well as maximum allowable concentrations of several heavy metals in shrimp for human consumption

Location	Species	Cd	Cu	Fe	Pb	Zn
Bay of Izmir, Turkey <sup>1</sup>	<i>Penaeus kerathurus</i> **	0.20	6.14	4.45	0.96	14.62
Mediterranean, Turkey <sup>2</sup>	<i>Penaeus kerathurus</i> **	0.03	7.4	3.1	0.34	13.2
Bay of Izmir, Turkey <sup>3</sup>	<i>Penaeus kerathurus</i> **	-	4.75	6.70	-	10.80
Present study	<i>Penaeus kerathurus</i> *	0.103±0.077	22.03±4.50	25.76±7.10	0.50±0.27	49.96±8.67
Present study	<i>Penaeus kerathurus</i> **	0.025±0.018	5.33±1.09	6.24±1.72	0.12±0.07	12.10±2.10
Mar Menor, Spain <sup>4</sup>	<i>Penaeus kerathurus</i> **	0.5	12.0	12.2	0.4	22.2
Mediterranean <sup>5</sup>	<i>Penaeus kerathurus</i> **	-	5.2±2.7	-	-	22±16
Amrovikos Gulf, Greece <sup>6</sup>	<i>Penaeus kerathurus</i> *	-	19.0-25.2	-	-	-
Mediterranean <sup>7</sup>	<i>Penaeus kerathurus</i> **	0.16	1.5	-	0.46	-
Mediterranean, Spain <sup>8</sup>	<i>Penaeus kerathurus</i> **	0.03-0.05	-	-	0.05-0.79	-
Lesina Lagoon, Italy <sup>9</sup>	<i>Penaeus kerathurus</i> *	0.16	15.83	-	0.59	50.60
Sea of Marmara, Turkey <sup>10</sup>	<i>Parapenaeus longirostris</i> *	0.77±0.31	24.5±3.85	56.20±34.82	4.82±2.08	40.42±11.24
Yskenderun Bay, Turkey <sup>11</sup>	<i>Penaeus semisulcatus</i> *	2.7-5.0	27.9-41.9	8.7-16.9	15.4-28.6	50.1-63.1
Mediterranean, Turkey <sup>12</sup>	<i>Penaeus japonicus</i> *	0.64-0.91	15.71-22.59	52.3-113.4	3.70-5.40	18.0-24.7
Yskenderun Bay, Turkey <sup>13</sup>	<i>Metapenaeus monoceros</i> *	0.5-1.1	20.6-30.7	60.7-75.3	10.3-21.6	60.4-71.3
Mexico <sup>14</sup>	<i>Penaeus californiensis</i> *	ND	18.2	28	-	74
Sundrbans, India <sup>15</sup>	<i>Penaeus indicus</i> *	ND-4.50	14.50-63	-	8.5-12	20.50-87.50
Persian Gulf <sup>16</sup>	<i>Penaeus merguensis</i> *	0.07±0.03	17.86±11.53	17.72±9.72	-	40.20±15.21
Persian Gulf <sup>16</sup>	<i>Metapenaeus affinis</i> *	0.11±0.06	17.37±9.89	22.15±16.84	-	46.06±15.22
Bay of Bengal <sup>17</sup>	<i>Penaeus monodon</i> *	0.2-0.3	12.2-21.3	9.1-15.7	0.8-1.3	24.2-35.7
Pacific coast, Mexico <sup>18</sup>	<i>Penaeus vannamei</i> *	0.57	23.3	180.1	-	60.6
Persian Gulf, Iran <sup>19</sup>	<i>Penaeus semisulcatus</i> **	0.133±0.010	3.418±0.711	-	-	8.977±0.959
Mediterranean, Algeria <sup>20</sup>	<i>Parapenaeus longirostris</i> *	0.65	117.75	142.50	1.27	98.17
Gulf of Mexico <sup>21</sup>	<i>Penaeus setiferus</i> *	3.64	17.6	102.1	4.22	106
Gulf of California <sup>22</sup>	<i>Litopenaeus stylirostris</i> *	0.66	25.4	-	5.3	57.8
Chittagong Bangladesh <sup>23</sup>	<i>Metapenaeus monoceros</i> *	0.7	33.1	43.0	3.6	40.8
Arabian sea <sup>24</sup>	<i>Penaeus japonicus</i> **	0.47	-	837	-	7.11
Saronikos gulf, Greece <sup>25</sup>	<i>Parapenaeus longirostris</i> **	0.05	6.30	7.40	2.27	13.40
Ubatuba bay, Brazil <sup>26</sup>	<i>Xiphopenaeus kroyeri</i> *	2.09	48.97	-	34.49	67.83
WHO <sup>19**</sup>		0.2	10	-	-	1000
European Commission <sup>27**</sup>		0.5	-	-	0.5	-
UK (MAFF) <sup>19**</sup>		0.2	20	-	2.0	50
IAEA-407 <sup>28*</sup>		0.19	3.28	146	0.12	67.1
FDA <sup>29**</sup>		3	60	-	1.5	200
Turkish Food Codex <sup>30**</sup>		0.5	20	-	0.5	50

<sup>1</sup>Uysal and Tuncer (1982); <sup>2</sup>Balkas *et al.* (1982); <sup>3</sup>Kontas (2012); <sup>4</sup>Leon *et al.* (1982); <sup>5</sup>UNEP/FAO (1986); <sup>6</sup>Panayotidis and Florou (1989); <sup>7</sup>Medpol (1992); <sup>8</sup>Pastor *et al.* (1994); <sup>9</sup>D'Adamo *et al.* (2008); <sup>10</sup>Kurun *et al.* (2007); <sup>11</sup>Çogun *et al.* (2005); <sup>12</sup>Canli *et al.* (2001); <sup>13</sup>Kargin *et al.* (2001); <sup>14</sup>Paez-Osuna and Tron-Mayen (1995); <sup>15</sup>Mitra *et al.* (2012); <sup>16</sup>Pourang and Amini (2001); <sup>17</sup>Hossain and Khan (2001); <sup>18</sup>Paez-Osuna and Fernandez (1995); <sup>19</sup>Pourang *et al.* (2005); <sup>20</sup>Abdenour *et al.* (2000); <sup>21</sup>Vazquez *et al.* (2001); <sup>22</sup>Frias-Espicuetta *et al.* (2007); <sup>23</sup>Khan and Alam (1994); <sup>24</sup>Tariq *et al.* (1993); <sup>25</sup>Voutsinou-Taliadouri (1982); <sup>26</sup>Mantelatto *et al.* (1999); <sup>27</sup>European Commission (2011); <sup>28</sup>IAEA-407 (2003); <sup>29</sup>FDA (2009); <sup>30</sup>Turkish Food Codex (2008); \*µg g<sup>-1</sup> dry weight; \*\*µg g<sup>-1</sup> wet weight; ND: Not Detected

bay and found concentrations (µg g<sup>-1</sup> ww) of Cd, Cu, Fe, Pb and Zn in the edible muscle of *P. kerathurus* at 0.20, 6.14-4.75, 4.45-6.70, 0.96 and 14.62-10.80, respectively. In the investigation, metal concentrations (considering dry weight) of, except for Cu, Fe and Zn were lower than these. The mean Cd value [(0.103 µg g<sup>-1</sup> dw) (conceding, 0.024 µg g<sup>-1</sup> ww)] is lower than the 0.20 µg g<sup>-1</sup> ww reported by Uysal and Tuncer. The results for higher concentrations of copper, iron and zinc than other metals (Cd and Pb) in shrimp were similar to findings previously recorded for shrimp other waters of the world. The levels of almost all the metals measured in this study are relatively lower than those recorded in shrimp from other region of Turkey and the world (Table 2).

The information on the differences between the metal content of the sexes is more limited than for the other factors so far considered. The faster growing sex (usually the female) can be expected to contain lower

concentrations of metals but not necessarily a smaller total body burden (Pourang *et al.*, 2005). In this study, highly significant differences between sexes in Fe and Cu contents could be detected in October (Table 1). Mean Fe level in male shrimps was considerably higher (30.19 compared to 20.23 µg g<sup>-1</sup> dw). Regarding Cu the same case could be observed (27.58 µg g<sup>-1</sup> dw in males compared to 19.02 in females). Frenet and Alliot (1985) reported that bioaccumulation factor of copper in the prawn *Palaemonetes varians* was greater in males and juveniles. Paez-Osuna and Tron-Mayen (1995) observed sex based differences in concentrations of Zn, Ni, Cu and Cd in some tissues of shrimp *Penaeus californiensis*. In the case of Zn and Cu, the higher concentrations were observed in muscle of males.

Almost all metals measured in this study are relatively lower than the values recorded in shrimp species from other regions of the world. The findings of other studies are shown in the Table 2 and are compared

with the concentrations reported in this study and elsewhere in the world. Patterns of distribution of trace elements in edible tissues of aquatic organisms are very important from different aspects, especially from health of sea food consumers' point of view. Considering limit value for human consumption of Cd, Cu, Fe, Pb and Zn, these concentrations ranged within the recommended limits for human consumption and not represent a health risk (WHO, European Commission, UK (MAFF), FDA, Turkish Food Codex) (WHO, 2006, Table 2). Comparing the present data with guidelines and limits, it can be seen that the mean values of Cu and Pb in the muscle for both sexes were only higher than levels of IAEA. Cd concentrations in the species ranged from 0.02-0.20  $\mu\text{g g}^{-1}$  dw. Cd concentration in *P. kerathurus* was below the local legislative limit of 0.5  $\mu\text{g g}^{-1}$  ww and within EU regulations for crustaceans. Cu concentrations throughout the study ranged from 17.93-27.58  $\text{mg } \mu\text{g g}^{-1}$  dw. The lowest and highest Cu level was detected in January and October month. The Cu concentrations found in the species were below the local legislative limit of 20  $\mu\text{g g}^{-1}$  ww. The Turkish legislation establishes maximum levels for two of the metals studied, above which human consumption is not permitted; 0.5  $\mu\text{g g}^{-1}$  ww for Pb and 50  $\mu\text{g g}^{-1}$  ww for Zn.

### CONCLUSION

The knowledge of heavy metal concentrations in shrimp are very important with respect to nature management, human consumption of these species and to determine the most useful biomonitor species and the most polluted area. Information on the distribution pattern of toxic heavy metal pollutants in aquatic environment becomes important so as to know the accumulation of such pollutants in the organisms and final transfer to man through sea foods. A comparison of the present results with data reported for shrimp from other marine environments in general heavy metal concentrations in the shrimp samples are not higher in Izmir bay. Based on the sample analyzed, metal concentrations found in the muscles proved to be the acceptable values for human consumption.

### REFERENCES

Abdenmour, C., B.D. Smith, M.S. Boulakoud, B. Samraoui and P.S. Rainbow, 2000. Trace metals in marine, brackish and freshwater prawns (Crustacea, Decapoda) from Northeast Algeria. *Hydrobiologia*, 432: 217-227.

Balkas, T.I., S. Tugrel and I. Salihoglu, 1982. Trace metal levels in fish and crustacea from Northeastern Mediterranean coastal waters. *Mar. Environ. Res.*, 6: 281-289.

Canli, M., M. Kalay and O. Ay, 2001. Metal (Cd, Pb, Cu, Zn, Fe, Cr, Ni) concentrations in tissues of a fish *Sardina pilchardus* and a prawn *Peaenus japonicus* from three stations on the Mediterranean Sea. *Bull. Environ. Contam. Toxicol.*, 67: 75-82.

Cogun, H., T.A. Yuzereroglu, F. Kargin and O. Firat, 2005. Seasonal variation and tissue distribution of heavy metals in shrimp and fish species from the Yumurtalik coast of Iskenderun Gulf, Mediterranean. *Bull. Environ. Contam. Toxicol.*, 75: 707-715.

D'Adamo, R., M. Di Stasio, A. Fabbrocini, F. Petitto, L. Roselli and M.G. Volpe, 2008. Migratory crustaceans as biomonitors of metal pollution in their nursery areas. The Lesina lagoon (SE Italy) as a case study. *Environ. Monit. Assess.*, 143: 15-24.

European Commission, 2011. Commission regulation (EU) No 420/2011 of 29 April 2011 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs. *Official J. Eur. Union*, L111: 3-6.

FDA, 2009. Guide for the Control of Molluscan Shellfish. National Shellfish Sanitation Program, U.S. Food and Drug Administration, USA., Pages: 547.

Firat, O., G. Gok, H.Y. Cogun, T.A. Yuzereroglu and F. Kargin, 2008. Concentrations of Cr, Cd, Cu, Zn and Fe in crab *Charybdis longicollis* and shrimp *Penaeus semisulcatus* from the Iskenderun Bay, Turkey. *Environ. Monit. Assess.*, 147: 117-123.

Frenet, M. and A. Alliot, 1985. Comparative bioaccumulation of metals in *Palaemonetes varians* in polluted and non-polluted environments. *Mar. Environ. Res.*, 17: 19-44.

Frias-Espericueta, M.G., F.C. Izaguirre-Fierro, E.F. Valenzuela-Quinonez, J.I. Osuna-Lopez and D. Voltolina *et al.*, 2007. Metal content of the gulf of California blue shrimp *Litopenaeus stylirostris* (Stimpson). *Bull. Environ. Contam. Toxicol.*, 79: 214-217.

Hossain, M.S. and Y.S.A. Khan, 2001. Trace metals in penaeid shrimp and spiny lobster from the bay of Bengal. *Sci. Asia*, 27: 165-168.

IAEA-407, 2003. Trace elements and methylmercury in fish tissue. Reference Sheet, International Atomic Energy Agency. <http://www.iaea.org/nael/refmaterial/iaea407.pdf>.

Kargin, F., A. Donmez and H.Y. Cogun, 2001. Distribution of heavy metals in different tissues of the shrimp *Penaeus semisulcatus* and *Metapenaeus monoceris* from the Iskenderun Gulf. *Tukey: Seasonal variations. Environ. Contam. Toxicol.*, 66: 102-109.

- Kontas, A., 2012. A case study of trace metals in suspended particulate matter and biota before wastewater treatment plant from the Izmir Bay, Turkey. *Environ. Monit. Assess.*, 184: 2605-2616.
- Kucuksezgin, F., A. Kontas, O. Altay, E. Uluturhan and E. Darilmaz, 2006. Assessment of marine pollution in Izmir Bay: Nutrient, heavy metal and total hydrocarbon concentrations. *Environ. Int.*, 32: 41-51.
- Kurun, A., H. Balkis and N. Balkis, 2007. Accumulations of total metal in dominant shrimp species (*Palaemon adspersus*, *Palaemon serratus*, *Parapenaeus longirostris*) and bottom surface sediments obtained from the Northern Inner Shelf of the Sea of Marmara. *Environ. Monit. Assess.*, 135: 353-367.
- MEDPOL, 1992. Assessment of the state of pollution of the Mediterranean Sea by radioactive substances. MAP Technical Reports Series No. 62, Athens, Greece, pp: 60.
- Mantelatto, F.L.M., W.E.P. Avelar, D.M.L. Silva, A.C. Tomazelli, J.L.C. Lopez and T. Shuhama, 1999. Heavy metals in the shrimp *Xiphopenaeus kroyeri* (Heller, 1862) (Crustacea, Penaeidae) from Ubatuba Bay, Sao Paulo, Brazil. *Bull. Environ. Contam. Toxicol.*, 62: 152-159.
- Mitra, A., P. Barua, S. Zaman and K. Banerjee, 2012. Analysis of trace metals in commercially important crustaceans collected from UNESCO protected world heritage site of Indian Sundarbans. *Turk. J. Fish. Aquat. Sci.*, 12: 53-66.
- Paez-Osuna, F. and L. Tron-Mayen, 1995. Distribution of heavy metals in tissues of the shrimp *Penaeus californiensis* from the Northwest coast of Mexico. *Bull. Environ. Contam. Toxicol.*, 55: 209-215.
- Paez-Osuna, F. and R.C. Fernandez, 1995. Trace metals in the Mexican shrimp *Penaeus vannamei* from estuarine and marine environments. *Environ. Pollut.*, 87: 243-247.
- Panayotidis, P. and E. Florou, 1989. Bioconcentration of Cu and Cd in benthic organisms and fish of the Amvrakikos Gulf. *Proceedings of the Conference on Environmental Science and Technology*, September, 1989, Mytilini, Greece, pp: 377-387.
- Pastor, A., F. Hernandez, M.A. Peris, J. Beltran, J.V. Sancho and M.T. Castello, 1994. Levels of heavy metals in some marine organisms from the western Mediterranean area (Spain). *Mar. Pollut. Bull.*, 28: 50-53.
- Pourang, N. and G. Amini, 2001. Distribution of trace elements in tissues of two shrimp species from Persian Gulf and effects of storage temperature on elements transportation. *Water, Air Soil Pollut.*, 129: 229-243.
- Pourang, N. and J.H. Dennis, 2005. Distribution of trace elements in tissues of two shrimp species from the Persian Gulf and roles of metallothionein in their redistribution. *Environ. Int.*, 31: 325-341.
- Pourang, N., J.H. Dennis and H. Ghourchian, 2005. Distribution of heavy metals in *Penaeus semisulcatus* From Persian Gulf and possible role of metallothionein in their redistribution during storage. *Environ. Monitoring Assess.*, 100: 71-88.
- Rainbow, P.S., 1998. Phylogeny of Trace Metal Accumulation in Crustaceans. In: *Metal Metabolism in Aquatic Environments*, Langston, W.J. and M. Bebianno (Eds.). Chapman and Hall, London, pp: 285-319.
- Tariq, J., M. Jaffar, M. Ashraf and M. Moazzam, 1993. Heavy metal concentrations in fish, shrimp, seaweed, sediment and water from Arabian Sea, Pakistan. *Mar. Pollut. Bull.*, 26: 644-647.
- Turkmen, G. and H. Yilmazyerli, 2006. Some biological aspects of *Melicertus kerathurus* (Forsk., 1775) (Decapoda, Penaeidae) inhabiting Izmir Bay (Aegean Sea), Turkey. *Crustaceana*, 79: 583-591.
- UNEP/FAO, 1986. Assessment of the present state of pollution by cadmium, copper, zinc and lead in the Mediterranean Sea. Document UNEP/WG. 144/11, UNEP, Athens, Greece, pp: 41.
- Vazquez, F.G., V.K. Sharma, Q.A. Mendoza and R. Hernandez, 2001. Metals in fish and shrimp of the Campeche sound, Gulf of Mexico. *Bull. Environ. Contam. Toxicol.*, 67: 756-762.
- WHO, 2006. *Guidelines for Drinking-Water Quality*. 3rd Edn., World Health Organization, Geneva Pages: 515.
- Yilmaz, A.B. and L. Yilmaz, 2007. Influences of sex and seasons on levels of heavy metals in tissues of green tiger shrimp (*Penaeus semisulcatus* de Ham, 1844). *Food Chem.*, 101: 1664-1669.