

The Effect of Cu⁺², Fe⁺² and Cr⁺³ in Mineral Additives Enriched with Biosorption Process Form on Chosen Parameters of *in vitro* Caecal Fermentation in Laying Hens (Lohmann Brown)

¹Maciej Janeczek, ²Katarzyna Chojnacka, ³Nezir Yasar Toker, ¹Ewa Pecka, ¹Albert Czerski, ²Zuzanna Witkowska, ¹Aleksander Chroszcz, ¹Wojciech Zawadzki and ⁴Mariusz Korczyński

¹Department of Biostructure and Animal Physiology, Faculty of Veterinary Medicine, Wrocław University of Environmental and Life Sciences, 51-631 Wrocław, Poland

²Department of Biochemistry, Faculty of Veterinary Medicine, Istanbul University, 34320 Avcılar, Istanbul, Turkey

³Institute of Inorganic Technology and Mineral Fertilizers, Wrocław University of Technology, I-26, Smoluchowskiego 25, 50-372 Wrocław, Poland

⁴Department of Animal Hygiene and Welfare, Faculty of Biology and Animal Husbandry, Wrocław University of Environmental and Life Sciences, Chelmonskiego 38, C51-631 Wrocław, Poland

Abstract: Trace element additives with Cu (II), Fe (III) and Cr (II) were tested *in vitro* in a Hen Caecum Model. The levels of Short-Chain Fatty Acid (SCFA) were estimated in the 4th and 6th h of fermentation. Measurements of hydrogen and methane as fermentation end products were conducted. Any negative influence on the fermentation process were noted. The study observed a higher production of SCFA in the experimental group than in the control group. In the experiment an increase in propionate acid was observed except in the control and Cr (II) groups. It seems to be true that fermentation in experimental groups was more intensive and faster than in the control one. A result of intensive fermentation was a pH decrease in experimental groups but this was the within range value. The lack of a correlation was shown between the level of acetate and methane and hydrogen concentration.

Key words: Poultry, hen, biosorption, caecum, fermentation, organic sorbent, trace elements

INTRODUCTION

Trace element supplementation is currently one of the most important aspects of animal nutrition (Garg *et al.*, 2008; Pechova *et al.*, 2009; Janeczek *et al.*, 2012). Microelement deficiencies cause insufficiencies leading to production decrease both in animals and humans. Traditionally, trace elements are introduced as inorganic salts or organic mineral products (e.g., amino acid chelates, proteinates and polysaccharides). In poultry nutrition, the most popular form of trace element supplementation are inorganic salts. It is well known that in this form microelements have low bioavailability (Fairweather-Tait, 1999; Nicholson *et al.*, 1999; Caussy *et al.*, 2003). Additionally in the earlier mentioned case a large emission of the trace elements occurred into the environment (Gustafson and Olsson, 2004). On the other hand, the organic forms like helats are relative expensive and can have a negative influence on the

alimentary tract (Chowdhury *et al.*, 2004). In recent years the concept of the control of trace elements level in animal products like eggs, meat and milk has been the subject of discussion. This concept is known as biofortification (Johns and Byzaguirre, 2007). It seems that biofortification could potentially be an important instrument to solve the global problem of hidden hunger (Kennedy *et al.*, 2003; Zhao *et al.*, 2009; Janeczek *et al.*, 2012).

Promising but very rarely discussed in the literature, seem to be alternative products in which trace metals can be bound via naturally occurring processes like biosorption (Hossain *et al.*, 1998; Mrvcic *et al.*, 2007; Dobrzanski *et al.*, 2008; Michalak and Chojnacka, 2008; Michalak *et al.*, 2009; Chojnacka, 2010). Biosorption is defined as the passive binding and removal process of metal ions or metalloid species by biological material (Hossain *et al.*, 1998; Chojnacka, 2007a, b). The mechanism of biosorption is an ion exchange and complex formation process in the donor-acceptor system of

electrons from functional groups derived from polymers and macromolecules that are elements of the cell wall. The formation of complexes involves organic compounds containing atoms with free electron pairs, a coordination bond that is formed by oxygen, nitrogen and sulfur (Janeczek *et al.*, 2012). It is well known that ion exchange is the dominant biosorption mechanism (Crist *et al.*, 1981; Schiewer and Volesky, 1995; Kadukova and Vircikova, 2005). The natural biomass contains light metal ions, e.g., K⁺, Na⁺, Ca²⁺ and Mg²⁺ that are bound to the functional groups on the cell wall. In the biosorption process these ions are exchanged for other metal ions and their concentration in the solution after biosorption is higher than before the process. At the moment light metal ions and protons are released their binding sites are taken by the microelement ions from the solution (Schiewer and Volesky, 1995; Davis *et al.*, 2003). Metal ion binding to the cell wall because of physical adsorption occurs due to electrostatic interactions and Van der Waals forces. In the reaction of metal ions with the components of the cell wall extracellular precipitation may also occur (Veglio and Beolchini, 1997). The main factor responsible for the biosorption is cell wall composition.

The results of the use of biomass enriched with biosorption processes are interesting. In Cu (II) and Fe (II) supplementation as a source of micronutrients in laying hen diet, the following results were observed: increased shell strength lower breakage, higher egg mass lower egg white mass lower parameter L form CIE Color Model of yolk was and higher parameter a eggs fortified with Fe, Zn and Mn were obtained. Eggs from the experimental group were richer in all the examined micronutrients and macronutrients when compared to eggs from the control group (Zielinska, 2010). The preliminary results on the supplements obtained using this method demonstrated that microelements in that form are widely available. Promising results were obtained in the studies on laying hens where using a diet supplement containing Cr⁺³ obtained via biosorption resulted in a considerable increase in the concentration of that element in egg shells was demonstrated (Chojnacka, 2007a; Michalak *et al.*, 2009). The fermentation processes in the large intestine are of a high significance in poultry. The fermentation is led by a range of microorganisms. Dietary manipulations of animals results in changes in intestinal flora. The products of fermentation are Short Chain Fatty Acid (SCFA) which may provide 30-40% of metabolic energy (Engelhardt, 1995; Tsukahara and Ushida, 2000; Donalson *et al.*, 2008; Mista *et al.*, 2011). It is already known that a correlation exists between microorganism populations and SCFA production (Marounek *et al.*, 1996; Van der Wielen *et al.*, 2000; Donalson *et al.*, 2008). The

influence of the new mineral feed additives based on biomass that is enriched with microelements in biosorption on animal health including digestive tract status is completely unknown. In poultry, caecal fermentation has decided importance for metabolic energy production and animal health. The biosorbents affecting production of caecal metabolites are not known, yet. The aim of this study was to characterize the influence of microelement supplementation on fermentation parameters (SCFA, methane and hydrogen) in the caecum of hens during *in vitro* investigation.

MATERIALS AND METHODS

Methods of manufacturing diet supplements with microelements: Soybean meal (Vetos, Zebowice near Jawor) was enriched with microelement ions Cu (II), Fe (II) and Cr (III) via biosorption using inorganic salts allowed by law as feed additives: CuSO₄·5H₂O, FeCl₂·4H₂O, Cr(NO₃)₃·9H₂O (POCh, Gliwice). Biosorption was conducted using a column reactor with a bed of a volume of 0.1 dm³. Water of pH 5.0 demineralised using 0.1M NaOH/HCl (POCh, Gliwice) was applied in the process. The reaction of water was controlled using a Mettler-Toledo pH-meter (Seven Multi, Switzerland) equipped with an InLab413 electrode with temperature compensation. The biosorption process was conducted at a temperature of 20°C to bed saturation controlling the concentration of solution flowing out of the column. The enriched biomass was air dried for 48 h.

The elemental composition of the samples collected was determined using an ICP-OES plasma spectrometer (Varian Vista-MPX; Varian, PaloAlto, USA) in the Chemical Laboratory of Multi-elemental Analysis at Wroclaw University of Technology, accredited by ILAC-MRA and Polish Accreditation Center (PCA) (No. AB 696).

The Control group (C) and three experimental groups were distinguished (Table 1). The availability of a given element from biological preparations was examined in the three groups (Fe, Cr and Cu). The biological preparations were added separately to the prepared feed mixtures (NJT-214, Tasomix®).

The composition of mixtures was established so that it did not contain an addition of a given microelement

Table 1: Content of microelements in non enriched soybean meal before and after biosorption (N = 3)

Microelement	Groups	Microelement content in soybean meal before and after enrichment (mg g ⁻¹)	
		Non enriched soybean meal	Enriched soybean meal
Fe (II)	Fe	0.235±0.015	16.337±0.228
Cr (III)	Cr	0.022±0.002	20.588±0.212
Cu (II)	Cu	0.065±0.004	15.690±0.370

$\bar{x} \pm 8D$ shows the \pm values

added in the form of biological preparation (groups Fe, Cr, Cu) and the need for other microelements was fulfilled using inorganic salts. The other components of feed mixtures were the same for all the groups. The requirement for a given microelement was fulfilled 100% by the biological preparation in the experimental groups. The availability of each of the preparations was examined against the control mixture which was a feed mixture containing all microelements in the form of inorganic salts.

In each experimental group, 0.530 g of preparation with Cu (II), 2.755 g of preparation with Fe (II), 4.259 g of preparation with Cr (III) was added to 1 kg of the feed (Table 2). Laying hens microelement requirement:

- Cr: 2 mg kg⁻¹
- Fe: 45 mg kg⁻¹
- Cu: 8 mg kg⁻¹

Animals: The research material consisted of the amount of caecum collected from 50 hens from Hy-Line Brown line, aged 22 weeks. The animals were maintained in a furnished Battery Cage System under controlled microclimatic conditions. The experiments were carried out according to Local Bioethic Committee Permission No. 129/2010.

The obtained mixtures (Table 2) were used as an addition for the incubated content in the group C-control group the complete mixture, group feed with an addition of iron in an organic form group Cr-feed with chromium addition, Cu-group feed with copper addition.

In vitro caecal fermentation: The caecum was collected during dissection and the ingesta was obtained from it. After mixing the ingesta collected from each animal, 5 g of it was divided into serum bottles of a volume of 125 mL (Sigma-Aldrich®). Next, 40 mL of a suitable buffer of pH 7.3 was added to each bottle in order

to dilute the ingesta (Janssen *et al.*, 2009). The samples were subjected to incubation after addition to 1 g of supplement at a temperature of 39°C. The anaerobic conditions in the bottle were obtained after CO₂ passing. The bottles were tightly closed using a crowner (Restek) with rubber plugs and aluminum caps (Sigma-Aldrich®). The samples placed in the bottles were subjected to 4 and 6 h fermentation *in vitro* in a shaker with water bath. About 48 samples were conducted in total 6 with each feed mixture in particular fermentation hours.

Analysis of collected fermentation products: After incubation, the gas contained in the serum bottles was subjected to analysis in order to examine bacteria activity in the ingesta. The analysis of the produced hydrogen and methane was conducted using gas chromatography on a gas chromatograph (Agilent Technologies 7890A GC System) with TCD and FID detector. After collection of gas samples, measurement of pH of the ingesta was performed and then the sample was centrifuged and 4 M formic acid was added (0.1, 2 mL of solution) in order to inhibit the fermentation processes. The samples of liquid ingesta were subjected to analysis using a gas chromatograph (Agilent Technologies 7890A GC System) with an FID detector in order to determine total SCFA concentration and the percentage contribution of particular acids: acetic, propionic, isobutyric, butyric, isovaleric and valeric.

Statistical analysis: Statistical analyses were done using multi variate analyses of variance using Statistica 9.0 Software (StatSoft Poland, Krakow, Poland). Significant differences were determined using Duncan's test.

RESULTS AND DISCUSSION

The level of active acidity in the caecum content of the hens (Table 3) with the application of iron as a

Table 2: Chemical components of layer ration substrates for *in vitro* fermentation, control group (Tasomix®)

Ingredient content of diets	Groups			
	C	Fe	Cr	Cu
Ground corn (percentage of DM)	29.99	29.99	29.99	29.99
Triticale (percentage of DM)	15.00	18.00	15.00	18.00
Soybean meal (percentage of DM)	13.70	12.10	13.70	12.10
Wheat (percentage of DM)	12.00	10.60	12.00	10.60
Chalk (percentage of DM)	8.440	8.440	8.440	8.440
Decoction wheat-com (percentage of DM)	6.000	6.000	6.000	6.000
Pszenmix (percentage of DM)	4.500	4.400	4.500	4.400
Sunflower meal (percentage of DM)	4.200	4.300	4.200	4.300
Fats (percentage of DM)	2.300	2.400	2.300	2.400
MPU 2% Nioska St.Tow. (percentage of DM)	2.000	2.000 (no Fe)	2.000	2.000 (no Cu)
Dried full blood (percentage of DM)	1.800	1.700	1.800	1.700
Mycifix select (percentage of DM)	0.050	0.050	0.050	0.050
L-Lysine (percentage of DM)	0.020	0.020	0.020	0.020
Biosorbent addition for 1 kg of feed (g)	-	4.259 Fe (II)	4.259 Cr (III)	0.530 Cu (II)
Lucantin pigment-red	-	-	-	-

Table 3: Production of short-chain fatty acids and levels of the active acidity of the caecal contents of chickens of different mix forage and fermentation time *in vitro*

Groups	Fermentation time (h)	Statistical symbol	pH	SCFA mmol kg ⁻¹ caecal	Acetate	Propionate	Izobutyrate	Butyrate	Izovalerate	Valerate
					mmol kg ⁻¹ caecal (%)					
C	4	\bar{x}	6.79 ^A	135.28 ^{Bbc}	51.50	30.81	2.56 ^B	12.10	1.36 ^b	1.54 ^{ab}
		SD	0.10	28.66	2.49	2.47	0.53	2.35	0.36	0.70
	6	\bar{x}	6.78 ^A	158.43 ^{ab}	52.20	29.68	3.23 ^B	11.33 ^c	1.60 ^{bc}	1.92 ^{bc}
		SD	0.05	19.74	2.62	1.57	0.67	1.85	0.52	0.33
Fe	4	\bar{x}	7.05 ^B	134.97 ^{Bbc}	49.43	31.57	3.05 ^{Ba}	12.50	1.46 ^c	1.93 ^{bc}
		SD	0.46	5.43	2.53	2.31	0.43	2.45	0.45	0.07
	6	\bar{x}	6.76 ^A	200.80 ^{bc}	51.11	33.15 ^b	3.11 ^{Ba}	10.00 ^{aa}	1.21	1.39 ^{aa}
		SD	0.36	67.09	8.74	8.57	0.68	1.22	0.46	0.24
Cr	4	\bar{x}	7.32 ^B	222.36	46.81	34.05	3.65 ^B	11.84 ^{bc}	1.56 ^{bc}	2.05 ^{bc}
		SD	0.44	104.28	4.91	5.93	2.28	3.41	1.04	0.31
	6	\bar{x}	7.20 ^B	223.95 ^{bc}	44.64	30.53 ^b	6.75 ^A	8.80 ^{bc}	7.74	1.51 ^{Bc}
		SD	0.27	98.15	6.82	1.40	4.68	6.66	11.16	1.06
Cu	4	\bar{x}	6.79 ^A	136.62 ^{Bbc}	51.88	28.22 ^a	2.15 ^{Bb}	14.06 ^{Bab}	1.22	2.02 ^{bc}
		SD	0.02	25.85	1.06	0.61	0.29	0.20	0.36	0.50
	6	\bar{x}	6.67 ^A	205.61 ^A	52.05	29.49	2.89 ^B	12.50 ^{bc}	1.27	1.74
		SD	0.03	13.95	1.90	1.57	0.13	0.56	0.21	0.04

^{a,b}Statistically significant differences ($p \leq 0.05$); ^{A,B}Highly statistically significant differences ($p \leq 0.01$)

mineral supplement between 4th and 6th h of fermentation was subject to decrease ($p \leq 0.01$) (of about 0.3 of pH value). The above phenomenon was also observed in other experimental groups however, a decrease between fermentation hours was considerably lower and was from 0.1-0.12 pH. At the 4th h of fermentation, a higher ($p \leq 0.01$) pH content in hens ingesta was observed in the samples after iron and chromium addition (Fe 4 h and Cr 4 h) when compared to the samples with copper addition and the control group (C 4 h and Cu 4 h). At the 6 h of fermentation, the pH value of ingesta after chromium addition (Cr 6 h) was on a higher ($p \leq 0.01$) level when compared to the other groups (C 6 h, Fe 6 h and Cu 6 h).

The lowest level of volatile fatty acids (134.97 mmol kg⁻¹) in hen caecum was noted at the 4th h of fermentation with iron addition (Fe 4 h). The highest in turn (223.95 mmol kg⁻¹) was observed at the 6th h of fermentation when using chromium addition (Cr 6 h). At the 4th h of fermentation of caecum content with the addition of a mineral supplement in the form of chromium (Cr 4 h), an increase in SCFA content of 87.08 mmol kg⁻¹ was observed when compared to the control group (C 4 h). After 6 h fermentation of hen ingesta, the supplements used led to an increase in LKT production. An increased ($p \leq 0.05$) general level of acids of 65.52 mmol kg⁻¹ when compared to the control group (C 6 h) was noted with chromium application (Cr 6 h). An increase in SCFA production up to 30% was noted at the 6th h of fermentation of caecum content between the group with iron and copper addition (Fe 6 h, Cu 6 h) and the control group (C 6 h).

In *in vitro* conditions at the 6th h of fermentation of caecum content of the hens, the applied mineral supplements caused a decrease in the percentage contribution (mmol kg⁻¹ %) of acetic acid when compared to the control group. The lowest level of acetic acid was noted at the 4th and 6th h of fermentation using chromium

addition (46.81 and 44.64% mmol kg⁻¹, respectively). In the case of chromium addition, a decrease in acetic acid contribution was observed at the 6th h of fermentation when compared to the 4th h. The reverse tendency was noted in the other experimental groups. A decrease in propionic acid production by caecal bacteria at the 6th h of fermentation was noted in the control group and in the group with chromium addition when compared to the production at the 4th h of fermentation. The reverse relationship was noted in the other experimental groups.

An addition of iron and chromium led to an increase in propionic acid production in the examined caecum content while copper addition caused a decrease in that acid contribution when compared to the control group irrespective of the fermentation time.

At the 4th h of fermentation, a lower contribution of isobutyl acid was noted with a concurrent increase in the level of butyl acid when compared to the 6th h of fermentation irrespective of the feed additive used.

At the 4th h of fermentation, a lower level of isobutyl acid in caecum content was observed after an addition of copper (Cu 4 h) when compared to the control group (C 4 h). The reverse tendency was noted in the other experimental groups. At the 6th h of fermentation, the content of butyl acid in caecum content after copper addition (Cu 6 h) was at a higher level when compared to the control group (C 4 h). The reverse tendency was noted in the other experimental groups.

A lower ($p \leq 0.05$) level of butyl acid (even by 20%) was noted in the caecum content after iron addition at the 6th h of fermentation (Fe 6 h) when compared to the experimental group with an addition of copper and chromium (Cu 6 h, Cr 6 h).

At the 4th h of fermentation iron (Fe 4 h) influenced a higher ($p \leq 0.05$) contribution of isovaleric acid when compared to the ingesta of the control group (C 4 h).

An increase in the level of valeric acid was observed at the 4th h of caecum content fermentation with mineral supplement addition. At the 6th h of fermentation the contribution of valeric acid after iron application (Fe 6 h) was lower when compared to the control group (C 6 h; $p \leq 0.01$) and in relation to the group where chromium was applied (Cr 6 h; $p \leq 0.05$).

Hydrogen production at the 4th h of fermentation (Fig. 1) was at the highest level ($0.33 \text{ mmol kg}^{-1}$) in the control group while lowest with iron application ($0.18 \text{ mmol kg}^{-1}$). Both in the control group and in the groups where an addition of chromium and copper was used, an increase (even of about 54%-Cr) in hydrogen production was noted between the 4th and 6th h of fermentation. An increase in methane production of 47.65 and 13.38% in the control group and Cu group, respectively (Fig. 2) was observed between the 4th and

noted in the control group (C 6 h) when compared to caecum content production with chromium (Cr 6 h) and copper (Cu 6 h) application. The methane content in the samples collected at the 4th h of fermentation was at a higher level with iron application (Fe 4 h) when compared to the other groups.

In the caecum of hens, microbial fermentation is a source of SCFA. This process can be modified by a great number of factors (Marounek *et al.*, 1999; Shanmugavelu *et al.*, 2006). The SCFA play an important role in animal metabolism and they have also a significant importance in physiological equivalence in caecum as a biotope. Too low production of SCFA favors intestinal mucous colonization by pathogens like *Salmonella* sp. (Barnes *et al.*, 1979; Corrier *et al.*, 1995; Kubena *et al.*, 2001). It seems that the microflora of caecum and fermentation profile are important aspects of animal health.

The pH level in caecum should be of 5.65-7.8 (Jozefiak *et al.*, 2004). It has been proved that lower pH (pH-5.8) has a negative influence on acetate and propionate acid production (Meimandipour *et al.*, 2010). In the investigation pH level was in the physiological range of value but a decrease in its level was observed between 4th and 6th h. This pH decrease was higher in the experimental than control groups. Probably, this effect was caused by an increase in fermentation product concentration (Mista *et al.*, 2011). It should be taken into consideration that feed additives were generated on the basis of the soy pellets which contain carbohydrates and its role in pH decrease in caecum is well known (Berggren *et al.*, 1993). The results presented in this research do not indicate that low pH changes influence the qualitative composition of SCFA.

Plant polysaccharides are used in SCFA and gases like ammonia, carboxydate, methane and hydrogen in caecal production (Engelhardt, 1995; Tsukahara and Ushida, 2000; Donalson *et al.*, 2008). In the investigation the influence of microelement additives on SCFA increase was presented. The mean value of SCFA concentration/kg caecal content in a hen equals $107\text{-}151 \text{ mmol kg}^{-1}$ (Jozefiak *et al.*, 2004). According to Tsukahara and Ushida (2000) the SCFA production in caecum is time dependent, i.e., $32.1\text{-}35.2 \text{ mmol kg}^{-1}$ in the 1st h, $128.4\text{-}140.8 \text{ mmol kg}^{-1}$ in the 4th h and $192.6\text{-}211.2 \text{ mmol kg}^{-1}$ in the 6th h. In the investigation, the level of SCFA in the 6th h of fermentation was higher than in the control group but not in a statistically important manner except fermentation with Cr. It should be pointed out that the level of SCFA was higher in the group with Cr than in other experimental and control groups after 4 h of fermentation. The results indicate that the effect of biosorbent additive introduction was fermentation increase (pH level decrease and SCFA concentration level increase).




Fig. 1: Concentration of hydrogen (mmol kg^{-1} caecal content)




Fig. 2: Concentration of methane (mmol kg^{-1} caecal content). ^{a,b}Statistically significant differences ($p \leq 0.05$); ^{A,B}highly statistically significant differences ($p \leq 0.01$)

6th h of fermentation. The reverse tendency was noted in the other experimental groups. At the 6th h of fermentation, higher ($p \leq 0.05$) methane production was

SCFA production is an important parameter of microbial fermentation in caecum. It is already known that caecal SCFA are a significant factor in preventing caecal *Salmonella* colonization (Barnes *et al.*, 1979). Especially important is the propionate acid level because of the proven negative correlation between its caecal concentration and *Salmonella* colonization in young chickens (Saengkerdsub *et al.*, 2006). The relationship between propionate concentration and some microbial agents, e.g., *Salmonella typhimurium* and Enterobacteriaceae has been shown (Van der Wielen *et al.*, 2000). It was proved that the optimal proportion between acetate, propionate and butyrate acids should equal 3-5:2:1 (Marounek and Rada, 1998; Saengkerdsub *et al.*, 2006; Donalson *et al.*, 2008; Meimandipour *et al.*, 2010). Some other researchers suggest optimal concentrations of acetate, propionate, izobutyrate, butyrate, izovalerate and valerate of 48-65, 12-32, 0.3-3, 6-14, 2-3 and 3-10% in caecum (Marounek and Rada 1998; Saengkerdsub *et al.*, 2006; Donalson *et al.*, 2008). In the investigation similar proportions were observed. In the case of additional Fe and Cu, the concentration of propionate increased in the 6th h of fermentation but in Cr decreased.

The production of gas as an end product of caecal fermentation provides a reproducible *in vitro* method for determining the potential effects of feed additives on gut flora (Saengkerdsub *et al.*, 2006). Guo *et al.* (2003) results proved gut gas production increased after polysaccharides intake (e.g., yeast). Methane production level is an indicator of carbohydrate fermentation and it has a positive correlation with SCFA production. The reduction of acetate production as a result of methanogenesis can be energy-efficient (Varadyova *et al.*, 2000). High levels of hydrogen can be the result of its accumulation during intensive reduction processes in acetate production (Marounek and Rada, 1998). In the investigation significant changes of acetate concentration were not proved. The lack of correlation between levels of acetate and methane and hydrogen concentration was stated. However, a higher level of hydrogen concentration after 6 h of fermentation in group with Cr addition was observed.

It is known that hydrogen and methane production per hour of fermentation should be of 0.07 and 11.4 mmol kg⁻¹ of gut content (Saengkerdsub *et al.*, 2006). In 4 months old chickens the ratio between hydrogen and methane is 1:9 (Marounek and Rada, 1998). In this study the earlier mentioned ratio was similar. It seems to be true that fermentation in the group with Fe (III) additive was more intensive and faster than in other groups (Fig. 2).

The preliminary results of the investigations suggest that the introduction of trace element additives generated with biosorption do not have a negative influence on microbial fermentation in caecum of laying hens.

CONCLUSION

According to the analyzed parameters, it seems that the Cu (II) addition in the new form had the most positive effect on fermentation. Because of the positive accumulation of Fe (III), Cu (II) in technological processes high bioavailability and low emission to the environment, biosorbents can be an alternative source of trace elements for animal feeding.

ACKNOWLEDGEMENT

The research was granted by Polish Ministry of Science and Higher Education, Grant No. N N 209068140.

REFERENCES

- Barnes, E.M., C.S. Impey and B.J. Stevens, 1979. Factors affecting the incidence and anti-salmonella activity of the anaerobic caecal flora of the young chick. *J. Hyg. (Lond.)*, 82: 263-283.
- Berggren, A.M., I.M.E. Bjorck, E.M.G.L. Nyman and B.O. Eggum, 1993. Short chain fatty acid content and pH in caecum of rats given various sources of carbohydrate. *J. Sci. Food Agric.*, 63: 397-406.
- Caussy, D., M. Gochfeld, E. Gurzau, C. Neagu and H. Ruedel, 2003. Lessons from case studies of metals: Investigating exposure, bioavailability and risk. *Ecotoxicol. Environ. Saf.*, 56: 45-51.
- Chojnacka, K., 2007a. Biosorption and bioaccumulation of microelements by *Riccia fluitans* in single and multi-metal system. *Bioresour. Technol.*, 98: 2919-2925.
- Chojnacka, K., 2007b. The Application of Biosorption and Bioaccumulation of Toxic Metals in Environmental Pollution Control. In: *Hazardous Materials and Wastewater: Treatment, Removal and Analysis*, Lewinsky, A.A. (Ed.). NOVA Science Publisher, New York, USA., pp: 309-359.
- Chojnacka, K., 2010. Biosorption and bioaccumulation the prospects for practical applications. *Environ. Inter.*, 36: 299-307.
- Chowdhury, S.D., I.K. Paik, H. Namkung and H.S. Limb, 2004. Responses of broiler chickens to organic copper fed in the form of copper-methionine chelate. *Anim. Feed Sci. Technol.*, 115: 281-293.

- Corrier, D.E., D.J. Nisbet, C.M. Scanlan, A.G. Hollister and J.R. Deloach, 1995. Control of *Salmonella typhimurium* colonization in broiler chicks with a continuous-flow characterized mixed culture of cecal bacteria. *Poult. Sci.*, 74: 916-924.
- Crist, R.H., K. Oberholser, N. Shank and M. Nguyen, 1981. Nature of bonding between metallic ions and algal cell walls. *Environ. Sci. Technol.*, 5: 1212-1217.
- Davis, T.A., B. Volesky and A. Mucci, 2003. A review of the biochemistry of heavy metal biosorption by brown algae. *Water Res.*, 37: 4311-4330.
- Dobrzanski, Z., M. Korczynski, K. Chojnacka, H. Gorecki and S. Opalinski, 2008. Influence of organic forms of copper, manganese and iron on bioaccumulation of these metals and zinc in laying hens. *J. Elementol.*, 13: 309-319.
- Donalson, L.M., W.K. Kim, V.I. Chalova, P. Herrera and J.L. McReynolds *et al.*, 2008. *In vitro* fermentation response of laying hen cecal bacteria to combinations of fructooligosaccharide prebiotics with alfalfa or a layer ration. *Poult. Sci.*, 87: 1263-1275.
- Engelhardt, W., 1995. Absorption of Short-Chain Fatty Acids from the Large Intestine. In: *Physiological and Clinical Aspects of Short-Chain Fatty Acids*, Cummings J.H., J.L. Rombeau and T. Sakata (Eds.). Cambridge University Press, Cambridge, MA., USA., pp: 149-170.
- Fairweather-Tait, S.J., 1999. The importance of trace element speciation in nutritional sciences. *Fresenius J. Anal. Chem.*, 363: 536-540.
- Garg, A.K., V. Mudgal and R.S. Dass, 2008. Effect of organic zinc supplementation on growth, nutrient utilization and mineral profile in lambs. *Anim. Feed Sci. Technol.*, 144: 82-96.
- Guo, F.C., B.A. Williams, R.P. Kwakkel and M.W. Verstegen, 2003. *In vitro* fermentation characteristics of two mushroom species, an herb and their polysaccharide fractions, using chicken cecal contents as inoculum. *Poult. Sci.*, 82: 1608-1615.
- Gustafson, G.M. and I. Olsson, 2004. Partitioning of nutrient and trace elements in feed between body retention, faeces and urine by growing dairy-breed steers. *Acta Agric. Scand. Sect. A: Anim. Sci.*, 54: 10-19.
- Hossain, S.M., S.L. Barreto and C.G. Silva, 1998. Growth performance and carcass composition of broilers fed supplemental chromium from chromium yeast. *Anim. Feed Sci. Technol.*, 71: 217-228.
- Janeczek, M., K. Chojnacka, A. Chroszcz and A. Zielinska, 2012. New Generation of Dietary Supplements with Microelements for Livestock Possibilities and Prospects. In: *Livestock: Rearing, Farming Practices and Diseases*, Javed, M.T. (Ed.). NOVA Science Publisher, New York, USA pp: 17-44.
- Janssen, P.W.M., R.G. Lentle, C. Hulls, V. Ravindran and A.M. Amerah, 2009. Spatiotemporal mapping of the motility of the isolated chicken caecum. *J. Comp. Physiol. B: Biochem. Syst. Environ. Physiol.*, 179: 593-604.
- Johns, T. and P.B. Eyzaguirre, 2007. Biofortification, biodiversity and diet: A search for complementary applications against poverty and malnutrition. *Food Policy*, 32: 1-24.
- Jozefiak, D., A. Rutkowski and S.A. Martin, 2004. Carbohydrate fermentation in the avian ceca: A review. *Anim. Feed Sci. Technol.*, 113: 1-15.
- Kadukova, J. and E. Vircikova, 2005. Comparison of differences between copper bioaccumulation and biosorption. *Environ. Int.*, 31: 227-232.
- Kennedy, G., G. Nantel and P. Shetty, 2003. The scourge of hidden hunger: Global dimensions of micronutrient deficiencies. *Food Nutr. Agric.*, 32: 8-16.
- Kubena, L., Y. Kwon, J. Byrd, C. Woodward and R. Moore *et al.*, 2001. Drinking water treatment and dietary treatment effects on *Salmonella enteritidis* in leghorn hens during forced molt. *Poult. Sci.*, 80 (Suppl. 1): 88-88.
- Marounek, M. and V. Rada, 1998. Age effect on *in vitro* fermentation pattern and methane production in the caeca of chickens. *Physiol. Res.*, 47: 259-263.
- Marounek, M., O. Suchorska and O. Savka, 1999. Effect of substrate and feed antibiotics on *in vitro* production of volatile fatty acids and methane in caecal contents of chickens. *Anim. Feed Sci. Technol.*, 80: 223-230.
- Marounek, M., V.M. Rada and V. Benda, 1996. Effect of Ionophores and 2-bromoethanesulphonic acid in hen caecal methanogenic cultures. *J. Animal Feed Sci.*, 5: 425-431.
- Meimandipour, A., M. Shuhaimi, A.F. Soleimani, K. Azhar and M. Hair-Bejo *et al.*, 2010. Selected microbial groups and short-chain fatty acids profile in a simulated chicken cecum supplemented with two strains of *Lactobacillus*. *Poult. Sci.*, 89: 470-476.
- Michalak, I. and K. Chojnacka, 2008. The application of macroalga *Pithophora varia* wille enriched with microelements by biosorption as biological feed supplement for livestock. *J. Sci. Food Agric.*, 88: 1178-1186.
- Michalak, I., K. Chojnacka and P. Glavic, 2009. The possibilities of the application of feed additives from macroalgae in sustainable mineral animal feeding. *Am. J. Applied Sci.*, 6: 1458-1466.
- Mista, D., B. Kioliczewska, W. Zawadzki, E. Pecka and M. Steininger *et al.*, 2011. The effect of Linola and W92/72 transgenic flax seed on the rabbit caecal fermentation at *in vitro* study. *Pol. J. Vet. Sci.*, 14: 557-564.

- Mrvčić, J., D. Stanzer, V. Stehlik-Tomas, D. Skevin and S. Grba, 2007. Optimization of Bioprocess for production of copper-enriched biomass of industrially important *Sacharomyces cerevisiae*. *J. Biosci. Bioeng.*, 103: 331-337.
- Nicholson, F.A., B.J. Chambers, J.R. Williams and R.J. Unwin, 1999. Heavy metal contents of livestock feeds and animal manures in England and Wales. *Bioresour. Technol.*, 70: 23-31.
- Pechova, A., L. Misurova, L. Pavlata and R. Dvorak, 2009. The influence of supplementation of different forms of zinc in goats on the zinc concentration in blood plasma and milk. *Biol. Trace Elem. Res.*, 132: 112-121.
- Saengkerdsud, S., W.K. Kim, R.C. Anderson, D.J. Nisbet and S.C. Ricke, 2006. Effects of nitrocompounds and feedstuffs on *In vitro* methane production in chicken cecal contents and fluid. *Anaerobe*, 12: 85-92.
- Schiewer, S. and B. Volesky, 1995. Modeling of the proton-metal ion exchange in biosorption. *Environ. Sci. Technol.*, 29: 3049-3058.
- Shanmugavelu, S., G. Ruzickova, J. Zrustova and J.D. Brooker, 2006. A fermentation assay to evaluate the effectiveness of antimicrobial agents on gut microflora. *J. Microbiol. Methods*, 67: 93-101.
- Tsukahara, T. and K. Ushida, 2000. Effects of animal or plant protein diets on cecal fermentation in guinea pigs (*Cavia porcellus*), rats (*Rattus norvegicus*) and chicks (*Gallus gallus domesticus*). *Comp. Biochem. Physiol. A, Mol. Integr. Physiol.*, 127: 139-146.
- Van der Wielen, P.W.J.J., S. Biesterveld, S. Notermans, H. Hofstra, B.A. Urlings and F. Vanknapen, 2000. Role of volatile fatty acids in development of ceca microflora in broiler chickens during growth. *Applied Environ. Microbiol.*, 66: 2536-2540.
- Varadyova, Z., I. Zelenak and P. Siroka, 2000. *In vitro* study of the rumen and hindgut fermentation of fibrous materials (meadow hay, beech sawdust, wheat straw) in sheep. *Anim. Feed Sci. Technol.*, 83: 127-138.
- Veglio, F. and F. Beolchini, 1997. Removal of metals by biosorption: A review. *Hydrometallurgy*, 44: 301-316.
- Zhao, F.J., Y.H. Su, S.J. Dunham, M. Rakszegi, Z. Bedo, S.P. McGrath and P.R. Shewry, 2009. Variation in mineral micronutrient concentrations in grain of wheat lines of diverse origin. *J. Cereal Sci.*, 49: 290-295.
- Zielinska, A., 2010. Elaboration of production process of new generation of mineral feed additives based on microalgae biomass. Ph.D. Thesis, Wrocław University of Technology.