

Characteristics of *Clostridium perfringens* Isolated from Chicken in Korea

Young Ju Lee

College of Veterinary Medicine, Kyungpook National University, 702-701 Daegu, Republic of Korea

Abstract: *Clostridium perfringens* (*C. perfringens*) is major cause of Necrotic Enteritidis (NE) in poultry. *C. perfringens* types A and C are the only types linked to the disease observed in avian species and toxins alpha (encoded by *cpa*), beta (*cpb*), beta-2 (*cpb2*), enterotoxin (*cpe*) and *netB* are the most significant in disease. Researchers report here the toxinotyping of NE-producing and commensal isolates of *C. perfringens* from chickens in Korea. All isolates recovered from NE lesion or normal flora were type A (*cpa* positive, *cpb* negative, *etx* negative, *cpI_o* negative). The *cpb2* and *luxS* showed no significant difference between NE-producing and commensal isolates but of the 27 *netB* positive isolates, 25 (36.2%) had been recovered from 69 chickens with NE while only 2 (4.9%) isolates were recovered from 41 chickens with normal flora.

Key words: *Clostridium perfringens*, toxinotyping, flora, lesion, chickens

INTRODUCTION

Clostridium perfringens (*C. perfringens*) are spore-forming bacteria that can be found in the soil, litter, feed, intestine of all animals but can cause Necrotic Enteritidis (NE) in poultry. The presence of intestinal damage produced by coccidia are considered predisposing factors that increases *C. perfringens* proliferation and the incidence of NE (Al-Sheikhly and Al-Saieg, 1980; Annett *et al.*, 2002).

C. perfringens types A, B, C, D and E are currently recognized as the major pathogenic contributors. Types A and C are the only types linked to the disease observed in avian species (Tschirdewahn *et al.*, 1991; Cooper and Songer, 2009). The *C. perfringens* strains that infect avian species produce many extracellular toxins however, toxins alpha (encoded by *cpa*), beta (*cpb*), beta-2 (*cpb2*), enterotoxin (*cpe*) and *netB* are the most significant (Engstrom *et al.*, 2003; Martin and Smyth, 2009).

Researchers report here the toxinotyping of NE-producing and commensal isolates of *C. perfringens* from chickens in Korea. In addition, researchers have investigated these toxin-encoding genes to genotype *C. perfringens* isolated from healthy or disease chickens in order to identify an association between the possession of toxin encoding genes and disease.

MATERIALS AND METHODS

Bacterial isolates and growth conditions: Isolates for the present study had been obtained from chickens by swabbing of small intestine of 110 birds which died during

rear on broiler breeder farm or commercial broiler farms. Sixty nine isolates were recovered from the NE-producing chickens. These birds has originated from 1 broiler breeder farm and 3 broiler farms which were experiencing an increased mortality rate and showed clinical NE characterized by confluent mucosal necrosis of large part of the small intestine, covered with a yellow-brown or bile-stained pseudomembrane or depression in the mucosal surface with discolored, amorphous material adhering to the mucosal surface. Forty one chickens originated 4 broiler farms showed no evidence of NE but 41 isolates were subsequently recovered for examination based on the comparison the toxinotype between NE-producing and commensal isolates. All swabs were streaked onto 5% sheep blood agar and incubated overnight at 37°C under anaerobic conditions. Resultant colonies which exhibited a double zone of hemolysis and which failed to grow under aerobic conditions were tentatively identified as *C. perfringens*. Isolates were preserved in preservation media (Brucella broth; BBL, Mississauga, Canada) and 50% buffered glycerol (1 M MgSO₄, 1 M Tris pH 8 and H₂O) and were kept at -80°C until use.

Preparation of DNA: Template DNAs for PCR were extracted using a QIAamp DNA Mini kit (Qiagen, France) according to manufacturer's directions. The DNA concentration was measured spectrophotometrically at A₂₆₀.

Toxinotyping by polymerase chain reaction: All isolates which had been tentatively identified as *C. perfringens*

Table 1: Primers used for toxinotyping in this study

Target genes	Primer	Sequence (5'-3')	Amplicon size (bp)	Annealing temperature	References
<i>cpa</i>	Forward	GTTGATAGCGCAGGACATGTTAAG	324	55	Garmory <i>et al.</i> (2000)
	Reverse	CATGTAGTCATCTGTCCAGCATC			
<i>cpb</i>	Forward	ACTATACAGACAGATCATTCAACC	195	55	Garmory <i>et al.</i> (2000)
	Reverse	TTAGGAGCAGTTAGAACTACAGAC			
<i>cpb2</i>	Forward	AGATTTTAAATATGATCCTAAC	567	53	Bueschel <i>et al.</i> (2003)
	Reverse	CCAATACCCCTTCACCAAATACTC			
<i>cpe</i>	Forward	GGAGATGGTTGGATATTAGG	233	53	Bueschel <i>et al.</i> (2003)
	Reverse	GGACCAGCAGTTGTAGATA			
<i>etx</i>	Forward	GCGGTGATATCCATCTATTTC	655	53	Garmory <i>et al.</i> (2000)
	Reverse	CCACTTACTTGTCTACTAAC			
<i>cpI_o</i>	Forward	ACTACTCTCAGACAAGACAG	446	53	Garmory <i>et al.</i> (2000)
	Reverse	CTTTCCTTCTATACTATAACG			
<i>luxS</i>	Forward	ACAAAGGTTAAGGCACCATATGT	385	53	Ohtani <i>et al.</i> (2002)
	Reverse	ACCTGTTTGGCATGACTCTAGCT			
<i>netB</i>	Forward	GCTGGTGCTGGAATAAATGC	384	53	Keyburn <i>et al.</i> (2008)
	Reverse	TCGCCATTGAGTAGTTTCCC			

were tested for the presence of *cpa*, *cpb*, *etx*, *cpI_o*, *cpb2*, *cpe* and *luxS* which encode, respectively for alpha toxin, beta toxin, epsilon, Iota, beta-2 toxin, enterotoxin and *AI-2* and *netB* gene. All primers are shown in Table 1.

DNA sequencing: All products obtained following PCR for *netB* were purified using the QIAquick PCR purification kit (Qiagen, Valencia, CA) according to the manufacturer's instructions. The purified PCR products were sequenced then sequences were compared in the GenBank database using Basic Local Alignment Search Tool (BLAST) through the internet (<http://www.ncbi.nlm.nih.gov/BLAST>) with an Accession No. EU143239 (Keyburn *et al.*, 2008).

RESULTS AND DISCUSSION

Toxinotyping: All presumptive *C. perfringens* isolates were confirmed as *C. perfringens* by toxinotyping. All isolates recovered from NE lesion or normal flora were type A (*cpa* positive, *cpb* negative, *etx* negative, *cpI_o* negative, Table 2).

Distribution of *cpb2* among chicken isolates: Of the 82 *cpb2* positive isolates from chickens, 53 (76.8%) isolates had been recovered from 69 chickens with NE while 26 (63.4%) isolates were recovered from 41 chickens which did not have evidence of NE at necropsy examination by experienced personnel.

Distribution of *luxS* among chicken isolates: Of the 110 *luxS* positive isolates, 68 (98.6%) and 40 (97.6%) isolates had been recovered from 69 chickens with NE and 41 chickens with normal flora, respectively.

Distribution of *netB* among chicken isolates: Of the 27 *netB* positive isolates, 25 (36.2%) had been recovered

from 69 chickens with NE while only 2 (4.9%) isolates were recovered from 41 chickens with normal flora.

In this study, *C. perfringens* isolates from NE-producing or healthy chickens were genotype A, a genotype which has previously been associated with NE in chicken (Park *et al.*, 1996). *C. perfringens* type A commonly occurs in North America and is one of two types associated with NE (Songer, 1996). Alpha toxin was found in all bacterial cultures from the field isolates, normal flora and post-challenge isolates. In the present study, alpha toxin was recurrently found to be the only major toxin pointing to *C. perfringens* type A as the main type noted, apart from a few exceptions that are described later in this manuscript.

Toxin genes for *cpb*, *cpe* and *etx* were absent in all field strains, normal flora and post-challenge isolates. Toxins *cpb* or *etx* are associated with *C. perfringens* types other than type A and would have changed the toxinotype noted. Toxin gene *cpe* is known to cause human food poisoning and may be found in type A and type C *C. perfringens* but is not commonly seen in poultry (Gholamiandekhordi *et al.*, 2006; Songer, 1996; Tschirdewahn *et al.*, 1991).

It has been suggested that potentially virulent toxins *cpb2* and *netB* could contribute to disease effects in field isolates (Bueschel *et al.*, 2003; Keyburn *et al.*, 2008). In the present research, these genes were also found in varying degrees in normal flora. However, no obvious NE was noted in the birds used for normal flora samples despite the presence of at least one of these toxin genes in all of the isolates. The evidence of the level of virulence of these toxin genes remains inconclusive. The results agree with those of previous studies which suggested that *cpb2* and *netB* did not necessarily have a direct correlation with the incidence of NE (Gholamiandekhordi *et al.*, 2006; Martin and Smyth, 2009). Positive PCR results for both *netB* and *cpb2* toxin genes in the field isolates could play a role in pathogenesis and

Table 2: *C. perfringens* PCR toxinotyping profile between to necrotic enteritis producing and normal small intestine

Isolates origin	Toxin genes							
	<i>cpa</i>	<i>cpb</i>	<i>cpb2</i>	<i>cpe</i>	<i>etx</i>	<i>cp10</i>	<i>luxS</i>	<i>netB</i>
NE-producing chickens (n = 69) ^a	69 (100) ^b	0 (0)	53 (76.8)	0 (0)	0 (0)	0 (0)	68 (98.6)	25 (36.2)
Healthy chickens (n = 41)	41 (100)	0 (0)	26 (63.4)	0 (0)	0 (0)	0 (0)	40 (97.6)	2 (4.9)
Total	110 (100)	0 (0)	79 (71.8)	0 (0)	0 (0)	0 (0)	109 (99.1)	27 (24.5)

^an = number; ^bNo. of isolates included (%)

Table 3: *C. perfringens* PCR toxinotyping patterns between necrotic enteritis producing and normal small intestine origin

Patterns of toxin gene included	No. of strains by origin (%)	
	NE-producing chickens (n = 69) ^a	Healthy chickens (n = 41)
<i>cpa-luxS-netB-cpb2</i>	18 (26.1)	1 (2.4)
<i>cpa-luxS-netB</i>	6 (8.7)	1 (2.4)
<i>cpa-luxS-cpb2</i>	23 (33.3)	26 (63.4)

^an = number

may have enhanced necrosis in the intestine. However, some studies have shown that β -2 toxin may not be significantly associated with NE (Bueschel *et al.*, 2003; Gholamiandekhordi *et al.*, 2006; Keyburn *et al.*, 2008). It should be noted that the occurrence of a gene does not necessarily determine phenotype of toxin production which may be a stronger determinant of virulence (Abildgaard *et al.*, 2010) (Table 3).

Positive findings for *luxS* were expected based on the level of conservation of this gene between bacterial types (Bassler, 1999). As a quorum-sense-related gene, *luxS* is reliant on predisposing factors to increase colonization by causing a shift in the gut ecology and bacterial numbers before allowing toxin production. In the case of alpha toxin, it was found that *luxS* had only a slight stimulatory effect and was not completely dependent on *luxS* for toxin production (Ohtani *et al.*, 2002). However, in the results using NE-producing strains there was no difference between *luxS* findings in normal flora versus NE-producing strains.

CONCLUSION

This study provides a baseline for prevalence of *C. perfringens* isolated from NE-producing or healthy chickens. Therefore, more research into much better correlation between gene expression of toxin and the pathogenic capacity of *C. perfringens* may be beneficial for an epidemiological approach of NE-producing farms.

ACKNOWLEDGEMENT

This research was supported by Kyungpook National University Research Fund, 2012.

REFERENCES

- Abildgaard, L., T.E. Sondergaard, R.M. Engberg, A. Schramm and O. Hojberg, 2010. *In vitro* production of necrotic enteritis toxin B, NetB, by netB-positive and netB-negative clostridium perfringens originating from healthy and diseased broiler chickens. *Vet. Microbiol.*, 144: 231-235.
- Al-Sheikhly, F. and A. Al-Saieg, 1980. Role of Coccidia in the occurrence of necrotic enteritis of chickens. *Avian Dis.*, 24: 324-333.
- Annett, C.B., J.R. Viste, M. Chirino-Trejo, H.L. Classen, D.M. Middleton and E. Simko, 2002. Necrotic enteritis: Effect of barley, wheat and corn diets on proliferation of *Clostridium perfringens* type A. *Avian Pathol.*, 31: 598-601.
- Bassler, B.L., 1999. How bacteria talk to each other: Regulation of gene expression by quorum sensing. *Curr. Opin. Microbiol.*, 2: 582-587.
- Bueschel, D.M., B.H. Jost, S.J. Billington, H.T. Trinh and J.G. Songer, 2003. Prevalence of *cpb2*, encoding beta2 toxin in *Clostridium perfringens* field isolates: Correlation of genotype with phenotype. *Vet. Microbiol.*, 94: 121-129.
- Cooper, K.K. and J.G. Songer, 2009. Necrotic enteritis in chickens: A paradigm of enteric infection by *Clostridium perfringens* type A. *Anaerobe*, 15: 55-60.
- Engstrom, B.E., C. Fermer, A. Lindberg, E. Saarinen, V. Baverud and A. Gunnars-Son, 2003. Molecular typing of isolates of *Clostridium perfringens* from healthy and diseased poultry. *Vet. Microbiol.*, 94: 225-235.
- Gamory, H.S., N. Chanter, N.P. French, D. Bueschel, J.G. Songer and R.W. Titball, 2000. Occurrence of *Clostridium perfringens* beta2-toxin amongst animals, determined using genotyping and subtyping PCR assays. *Epidemiol. Infect.*, 124: 61-67.
- Gholamiandekhordi, A.R., R. Ducatelle, M. Heyndrickx, F. Haesebrouck and F. Van Immerseel, 2006. Molecular and phenotypical characterization of *Clostridium perfringens* isolates from poultry flocks with different disease status. *Vet. Microbiol.*, 113: 143-152.

- Keyburn, A.L., J.D. Boyce, P. Vaz, T.L. Bannam and M.E. Ford *et al.*, 2008. NetB, a new toxin that is associated with avian necrotic enteritis caused by *Clostridium perfringens*. *PLoS Pathog.*, Vol. 4. 10.1371/journal.ppat.0040026.
- Martin, T.G. and J.A. Smyth, 2009. Prevalence of netB among some clinical isolates of *Clostridium perfringens* from animals in the United States. *Vet. Microbiol.*, 136: 202-205.
- Ohtani, K., H. Hayashi and T. Shimizu, 2002. The *luxS* gene is involved in cell-cell signalling for toxin production in *Clostridium perfringens*. *Mol. Microbiol.*, 44: 171-179.
- Park, K.Y., S.U. Lee, H.S. Yoo and J.G. Yeh, 1996. Toxigenic type of *Clostridium perfringens* isolated from chicken in Korea. *Korean J. Vet. Res.*, 36: 829-837.
- Songer, J.G., 1996. Clostridial enteric diseases of domestic animals. *Clin. Microbiol. Rev.*, 9: 216-234.
- Tschirdewahn, B., S. Notermans, K. Wernars and F. Untermann, 1991. The presence of enterotoxigenic *Clostridium perfringens* strains in faeces of various animals. *Int. J. Food Microbiol.*, 14: 175-178.