

Atherosclerosis and Animal Models

¹Guangwei Zhang, ^{2,3}Yulong Chen, ^{2,3}Yan Lin and ^{2,3}Enqi Liu

¹Xi'an Medical University, 28 Xianning West Road, Beilin,

²Cardiovascular Research Center,

³Laboratory Animal Center, School of Medicine, Xi'an Jiaotong University,
Xi'an, 710061 Shaanxi, China

Abstract: Cardiovascular disease is currently the major cause of mortality and morbidity in developing and developed countries. Atherosclerosis is the core of cardiovascular disease and is complex inflammatory process. Many animal species have been used to investigate the pathogenesis of atherosclerosis. However, the most useful animal models have been restricted to genetically modified mice or larger animal models including rabbits, pig and non-human primates. Mouse models are widely used for atherosclerosis research due to genetic manipulation. Murine atherosclerosis are different from that of human. Thus, the models are used for investigation of biological processes of atherosclerosis. Rabbits do not develop spontaneous atherosclerosis but they are highly responsive to cholesterol diet and develop plaque in a short time. Pig and monkey are better suitable models because their plasma levels and atherosclerotic plaque are similar to that of human. However, these models are not widely used. Relatively high cost and ethical concern may be major causes.

Key words: Atherosclerosis, animal model, mouse, rabbits, cardiovascular disease

INTRODUCTION

Cardiovascular disease is currently the major cause of mortality and morbidity in developing and developed countries. Its incidence is expected to increase in the future (Bentzon and Falk, 2010). Cardiovascular disease is complex multifactor pathologies which including genetic and environmental factors. Atherosclerosis is the core of cardiovascular disease leading to myocardial infarctions, stroke and peripheral vascular disease encountered in most human populations. Atherosclerosis is complex inflammatory process (Zadelaar *et al.*, 2007). So, yet its mechanism is not clear. However, development of animal modes provide us platform for studying pathophysiology and molecular mechanism of atherosclerosis. Additionally, animal models are also essential tools to evaluate effectiveness of preventative and therapeutic strategies. Researchers will summarize the common models of atherosclerosis in this review. It will be useful to further understand development of the models of atherosclerosis.

MICE MODEL

Wild-type mice are resistant to atherosclerosis due to high levels of HDL and low levels of LDL and VLDL. Mice need to be genetically modified to study atherosclerosis. Some of the most widely used

strains are Apolipoprotein E (ApoE)-deficient mice, LDL receptor-deficient mice and ApoE*3-Leiden (E3L).

ApoE as a constituent of plasma lipoprotein is synthesized in the liver and macrophages and serves as a ligand for LDL receptor and LDL Receptor-related Proteins (LRP). Deletion of the ApoE in mice increase levels of plasma LDL and VLDL due to the failure of LDL receptor and LRP-mediated clearance of these lipoprotein. On a chow diet, mice exhibit marked hypercholesterolemia (a total cholesterol level >400 mg dL^{-1}) and they develop extensive atherosclerotic lesion widely distributed throughout the aorta. Western diet could accelerate formation of atherosclerotic lesion via rapidly elevating cholesterol level (>1000 mg dL^{-1}) (Nakashima *et al.*, 1994). The ApoE^{-/-} mice contains the entire spectrum of lesions observed during atherogenesis. The sequential events involved in lesion formation which is similar to those in human. Fatty streaks were observed in 3 months old mice ApoE^{-/-} fed with chow diet. While fed chow, foam cell lesions first appear at 8-10 weeks of age. After 15 weeks, intermediate lesions are present, containing spindle shaped cells (mostly smooth muscle cells) and beyond 20 weeks, fibrous plaques are evident containing smooth muscle cells, extracellular matrix and an overlying fibrous cap. Predilection sites for atherosclerotic lesion are the aortic root followed by the aortic arch, the brachiocephalic trunk, left carotid and subclavian and

coronary arteries (Hu *et al.*, 2005; Bentzon and Falk, 2010). In older ApoE^{-/-}-mice, hemorrhage has been observed in lesions suggesting some degree of lesion instability (Rosenfeld *et al.*, 2000). However, high levels of plasma cholesterol is major characters in the mice (Scalia *et al.*, 2001). Furthermore, the most plasma cholesterol is confined to VLDL and not to LDL particles as in human. In addition, ApoE exerts antiatherogenic properties via antioxidant, antiinflammatory, antiplatelet and up-regulation of NO (Grainger *et al.*, 2004; Ali *et al.*, 2005; Davignon, 2005; Raffai *et al.*, 2005). The immunomodulatory effects of ApoE is important in development of atherosclerosis. It inhibits T-cell proliferation (Davignon, 2005). Macrophage derived ApoE has an independent role in lesion development, perhaps related to the promotion of reverse cholesterol transport (Mazzone and Reardon, 1994). However, the transfer of bone marrow from a mouse expressing ApoE into ApoE^{-/-}-recipient reduced atherosclerosis but also reduced plasma lipid level (Linton and Fazio, 1999).

In human, mutations in the gene for the LDL receptor cause familial hypercholesterolemia. The absence of the LDL receptor influences lipoprotein uptake and clearance, resulting in high levels of plasma VLDL and LDL in mice (Teupser *et al.*, 2003; Barcat *et al.*, 2006; Bentzon and Falk, 2010). LDL receptor^{-/-}-mice displays a moderate increase of plasma cholesterol level and develop atherosclerosis slowly on chow diet (Bentzon and Falk, 2010). On chow, limited lesion only develop in older LDL receptor^{-/-}-mice (Teupser *et al.*, 2003; Barcat *et al.*, 2006). However, the severity of hypercholesterolemia and atherosclerotic lesions requires the feeding of a high-fat, high-cholesterol diet (Knowles and Maeda, 2000). High-fat diet could induce accumulation of larger VLDL/remnant lipoproteins in addition, plasma cholesterol levels are much higher than in chow-fed ApoE^{-/-}-mice (Hartvigsen *et al.*, 2007). Lesions in the LDL receptor^{-/-}-mice have a greater preponderance of foam cells than in chow-fed ApoE^{-/-}-mice. The systematic pathological analysis of lesion development were not reported in LDL receptor^{-/-}-mice (Linton *et al.*, 1999). The advantage of this model is that LDL receptor dose not have the multitude of functions of ApoE.

The ApoE*3-Leiden mutation is a rare dominant negative mutation in the human *ApoE3* gene. ApoE*3-Leide transgenic mice have been generated by introducing a human ApoE*3-Leiden gene construct into C57B1/6 mice. ApoE*3-Leide mice are very useful to explored the mechanisms by which the ApoE isoforms influence lipid metabolism and atherosclerosis (Hofker *et al.*, 1998; Pendse *et al.*, 2009). ApoE*3-Leide mice show elevations of plasma cholesterol and

triglycerides on chow diet. High-fat, high-sugar and high-cholesterol could accelerate this process (Van Vlijmen *et al.*, 1994). In addition, ApoE*3-Leide mice also represent Dysbetalipoproteinemia Model in which plasma cholesterol and triglycerides are mainly VLDL and LDL-sized lipoprotein fraction (Hofker *et al.*, 1998). This model develop atherosclerotic lesion with all the characteristics of human pathology, varying from fatty streak to mild, moderate and severe plaques (Leppanen *et al.*, 1998; Lutgens *et al.*, 1999).

Three mouse models have been used extensively as a springboard for study of other genes affecting atherogenesis. The murine products take the form of either double knockout mice or overexpression a gene. LDL receptor^{-/-}-mice coupled with an ApoB-editing deficiency or combined with human ApoB 100 transgenic mice show a larger increase in plasma LDL cholesterol and develop atherosclerosis on a low-fat diet (Powell-Braxton *et al.*, 1998; Sanan *et al.*, 1998). ApoE*3-Leide mice crossbred with human Cholesteryl Ester Transfer Protein (CETP)-expressing mice display an elevated basal cholesterol level and an even more human-like lipoprotein profile. CETP expression can shift the distribution of cholesterol from HDL toward VLDL/LDL (Rensen and Havekes, 2006; Westerterp *et al.*, 2006). ApoE^{-/-}-mice overexpressing human ABCA1 do not protect against plaque development (Singaraja *et al.*, 2002). Complete absence of ABCA1 in the setting of ApoE deficiency resulted in plasma lipid reduction and tissue foam cell accumulation but no increase in atherosclerosis. However, selective absence of ABCA1 in macrophages resulted in markedly increased atherosclerosis with no effect on plasma lipids (Aiello *et al.*, 2002). ApoE^{-/-}-mice over-expressing ApoA-I show decreased lesion size with increased HDL formation (Paszty *et al.*, 1994; Benoit *et al.*, 1999). SR-A deletion in the ApoE^{-/-}-mice reduced atherosclerosis despite increased serum cholesterol (Suzuki *et al.*, 1997). Its overexpression reduced serum cholesterol but did not promote atherosclerosis (Van Eck *et al.*, 2000). SR-BI deletion in the ApoE^{-/-}-mice increased atherosclerosis compared with its overexpression (Zhang *et al.*, 2003). The double knockout ApoE^{-/-}-RAG^{-/-}-dose not prevent atherosclerosis (Daugherty *et al.*, 1997). The ApoE^{-/-}-mice overexpressing human C-reactive protein further promotes atherosclerosis development (Paul *et al.*, 2004) P-selectin^{-/-}-ApoE^{-/-}-double knockout mice also reduced size of aortic sinus lesion (Dong *et al.*, 2000) gene transfer of P53 to ApoE^{-/-}-mice did not alter plaque size but it cause a shift toward vulnerable plaque with a decrease in cap-to-intima area ratio, cap breaks and intraplaque hemorrhages (Von der Thusen *et al.*, 2002).

RABBIT MODEL

The rabbit is sensitive to dietary cholesterol (Dornas *et al.*, 2010). It as atherosclerosis model can supply useful information which is the progression and regression of atherosclerosis in human. Although, the rabbit express CETP not similar to mice is have low plasma levels of hepatic lipase and lack ApoA-II. Lipoprotein and atherosclerosis response were modulated by feeding cholesterol diets. On chow, the rabbit is not inherently prone to atherosclerosis due to low levels of plasma cholesterol (Finking and Hanke, 1997). High-cholesterol diet (0.5-4%) rapidly increases plasma cholesterol levels (>1000 mg dL⁻¹) and leads to the formation of lesion-enriched foam cell. This contributed to the mechanisms the lipid induce macrophage accumulation and other aspects of atheroma inflammation (Bustos *et al.*, 1998; Aikawa *et al.*, 2002). The development of advanced and complex atherosclerotic plaque requires long periods of cholesterol feeding. However, long term cholesterol feeding increases hepatic toxicity and induces a massive inflammatory response that does not resemble the chronic low-grade inflammatory response associated to human atherosclerosis. In addition, the formation of advanced lesion in rabbit depends on the age. Aged rabbits with 3-4.5 years old, frequently develop fibrous plaques whereas young animal (4 months of age) do not show advanced lesion (Dornas *et al.*, 2010). Rupture of the atherosclerotic plaque is known to be the major cause of thrombosis and subsequent clinical manifestations of atherosclerosis (Libby, 1995; Lee and Libby, 1997; Shiomi *et al.*, 2004). Models of plaque rupture in rabbit was been developed via the combination of aggressive vascular injury associated to a cholesterol diet. The aortic plaque had the three features of vulnerable plaque: lipid-rich core, accumulation of macrophages and a thin fibrous cap (Shimizu *et al.*, 2009). In addition, the lesion is commonly induced using an intravenously injected balloon containing Russel's viper venom and histamine a procedure that will result in plaque rupture and thrombosis (Johnstone *et al.*, 2001). The implantation of balloon catheter in the thoracic aorta also lead to lesion rupture (Rekhter *et al.*, 1998).

Some rabbits carry genetic mutation that lead to hyperlipidemia and atherosclerosis. The most widely used hyperlipidemic rabbit in atherosclerosis research is the Watanabe Heritable Hyperlipidemic (WHHL) rabbit. WHHL rabbit carry an inactivating mutation in the gene encoding the LDL receptor and exhibit hypercholesterolemia with increased plasma levels of LDL and VLDL reduced plasma levels of HDL. The WHHL rabbit develop aortic atherosclerotic plaque ranging from

fatty streaks to advanced lesion and exhibit severe coronary atherosclerosis (Shiomi *et al.*, 1992). Transgenic rabbits provide a important pathway for researching atherosclerosis development. Rabbits with liver-specific expression of human apolipoprotein A-I exhibit smaller atherosclerosis lesions than nontransgenic rabbit (Rekhter *et al.*, 2000; Broeders *et al.*, 2002). Rabbits that overexpress human lecithin-cholesterol acyltransferase have reduced atherosclerosis compared to rabbits-fed a cholesterol diet (Aikawa *et al.*, 1998). Furthermore, WHHL rabbits that express human apolipoprotein, exhibit accelerated atherosclerosis development and more complex lesions than nontransgenic rabbits. Additionally, human lipoprotein lipase overexpression in WHHL rabbit leads to enhanced aortic atherosclerosis (Abela *et al.*, 1995; Corseaux *et al.*, 2000).

PIG MODEL

Although, rodents and rabbits have been used extensively for atherosclerosis research these animal model limits the quantities of new agents required for *in vivo* screening due to different lipid physiology from that in humans and small size. Thus, larger Animal Models were pay attention for atherosclerosis research.

Pig is suitable atherosclerosis model. Many characteristics of pig resemble that of human such as development, morphology and function of the cardiovascular system, lipoprotein profiles and metabolism. Additionally, it develops spontaneous atherosclerotic lesions and may develop sudden death when under stress. Currently there is no single and golden standard animal models of vulnerable plaque but pig model are probably the best way to recreate human plaque instability. It is worth mentioning that pig can offer the ability to evaluate coronary arteries rather than central aortic in smaller animals (Badimon *et al.*, 1985; Fuster *et al.*, 1985; Palazon *et al.*, 1998; Royo *et al.*, 2000).

Atherosclerosis can be generally quite slow and spontaneously occurs by regular chow or high cholesterol diet induction in most arterial beds of pigs (Skold *et al.*, 1966; Fuster *et al.*, 1991; Casani *et al.*, 2005). Furthermore, the atherosclerotic plaque distribution and composition is similar to that in human (White *et al.*, 1992; White and Bloor, 1992; Gal *et al.*, 1992). This phenomenon may attribute to human like lipoprotein metabolism of pigs (Marzetta and Rudel, 1986). Additionally, atherosclerotic plaque first happen in coronary arteries of pigs. After gradually occlusion of coronary arteries, pigs may develop the coronary restenosis syndrome as human (Swindle, 1998).

Like in other animal models, atherosclerotic plaque development in pigs can be accelerated by high cholesterol diet. The high cholesterol diet firstly, induce hypercholesterolemia in pigs. Almost all pigs can develop fatty streaks localized in abdominal aorta and a lesser extent in the coronary arteries after 50 days with high cholesterol diet induction. Moreover, lesion composition was similar to early stage human atherosclerosis (Casani *et al.*, 2005). After 100-150 days by the diet induction, lesion severity was higher degree than that of induction for 50 days. Pigs employed tend to be within the prepuberty period in long period study. However, their lesions are not as severe as those developed in adult human. Although, older pigs can be used to obtain relative severity plaque as adult human, management becomes more difficult due to their considerable weight (Jorgensen, 2006). Thus, miniature pigs are preferable to atherosclerotic animal models because of their small size and growth rate.

Both diabetes and hypercholesterolemia are major risk factors for atherosclerosis. Moreover, these disease frequently co-exist in patients with metabolic syndrome. However, the animal models which investigate mechanism underlying diabetes accelerated atherosclerosis are scarce. People have found that pig are suitable model for atherosclerosis accompanying with diabetes. The combination of hypercholesterolemia and diabetes accelerates atherosclerotic lesion and makes animal handling easier and leads to the formation of advanced human like atherosclerotic lesion in pigs (Gerrity *et al.*, 2001; Mohler III *et al.*, 2008; Wilensky *et al.*, 2008).

The angioplasty and stenting have become established treatments for coronary artery disease. The part stent implantation (20-30%) initially appeared promising in so-called ideal lesion. However, there is still a high rate of vessel reclosure resulting in the need for repeat procedure (Edelman *et al.*, 1998). Compared to other animal models, pig coronary stenting is a suitable model because injury response is similar to human vessels with an adaptive response (Rodgers *et al.*, 1990). Moreover, size and anatomic distribution of pigs coronary arteries are similar to those of human. Furthermore, angiography, intravascular ultrasound, instrumentation and stent deployment in the pig are all similar to the clinical situation (Gershlick and Baron, 1998).

NON-HUMAN PRIMATES

Non-human primates models for atherosclerosis research include Cebus, Woolly, squirrel, African Green, Rhesus and Cynomolgus monkey. Although, these models lipoprotein metabolism is similar to human,

atherosclerosis susceptibility are different among them (Vesselinovitch *et al.*, 1974; Davis *et al.*, 1984; Mott *et al.*, 1992; Wolfe *et al.*, 1994). Rhesus, Cynomolgus, Cebus monkey are extensively researched because of distinct advantages. These models can develop spontaneous atherosclerosis in some arteries including the coronary arteries. High cholesterol diet greatly accelerates atherosclerosis development. Moreover, low fat diet can regress coronary atherosclerosis. The regression was associated with a reduction in cholesterol content of the lesion, consistent with a reduction in the number of foam cells (Mann *et al.*, 1953; Taylor *et al.*, 1954; Taylor *et al.*, 1962; Bullock *et al.*, 1969; Clarkson *et al.*, 1979; Clarkson *et al.*, 1984; Beere *et al.*, 1992). Additionally, male were more affected than female in non human primates (Bullock *et al.*, 1969). Older animals are more susceptible to atherosclerosis but young animals are resistant to atherosclerosis (Clarkson, 1998). Ethical concern limited utility for research of non human primates.

CONCLUSION

A variety of small and large animal models have been used to study the atherogenic process. No model is ideal. Each model has its own advantages and limitations with respect to manipulation of the atherogenic process and modeling human atherosclerosis or lipoprotein profile. The choice of the suitable animal model depends on the nature of the research that is to be performed.

ACKNOWLEDGEMENT

This research was partly supported by the National Natural Science Foundation of China (81070250) and by a Public Service Platform Grant of Shaanxi Province (2010FWPT-1).

REFERENCES

- Abela, G.S., P.D. Picon, S.E. Friedl, O.C. Gebara and A. Miyamoto *et al.*, 1995. Triggering of plaque disruption and arterial thrombosis in an atherosclerotic rabbit model. *Circulation*, 91: 776-784.
- Aiello, R.J., D. Brees, P.A. Bourassa, L. Royer and S. Lindsey *et al.*, 2002. Increased atherosclerosis in hyperlipidemic mice with inactivation of ABCA1 in macrophages. *Arterioscler., Thromb. Vasc. Biol.*, 22: 630-637.
- Aikawa, M., E. Rabkin, Y. Okada, S.J. Voglic and S.K. Clinton *et al.*, 1998. Lipid lowering by diet reduces matrix metalloproteinase activity and increases collagen content of rabbit atheroma: A potential mechanism of lesion stabilization. *Circulation*, 97: 2433-2444.

- Aikawa, M., S. Sugiyama, C.C. Hill, S.J. Voglic and E. Rabkin *et al.*, 2002. Lipid lowering reduces oxidative stress and endothelial cell activation in rabbit atheroma. *Circulation*, 106: 1390-1396.
- Ali, K., M. Middleton, E. Pure and D.J. Rader, 2005. Apolipoprotein E suppresses the type I inflammatory response *in vivo*. *Circ. Res.*, 97: 922-927.
- Badimon, L., P. Steele, J.J. Badimon, E.J. Bowie and V. Fuster, 1985. Aortic atherosclerosis in pigs with heterozygous von Willebrand disease. Comparison with homozygous von Willebrand and normal pigs. *Arterioscler. Thromb. Vasc. Biol.*, 5: 366-370.
- Barcat, D., A. Amadio, A. Palos-Pinto, D. Daret, P. Benlian, M. Darmon and A.M. Berard, 2006. Combined hyperlipidemia/hyperalphalipoproteinemia associated with premature spontaneous atherosclerosis in mice lacking hepatic lipase and low density lipoprotein receptor. *Atherosclerosis*, 188: 347-355.
- Beere, P.A., S. Glagov and Ch.K. Zarins, 1992. Experimental atherosclerosis at the carotid bifurcation of the cynomolgus monkey. Localization, compensatory enlargement and the sparing effect of lowered heart rate. *Arterioscler. Thromb. Vasc. Biol.*, 12: 1245-1253.
- Benoit, P., F. Emmanuel, J.M. Caillaud, L. Bassinet and G. Castro *et al.*, 1999. Somatic gene transfer of human ApoA-I inhibits atherosclerosis progression in mouse models. *Circulation*, 99: 105-110.
- Bentzon, J.F. and E. Falk, 2010. Atherosclerotic lesions in mouse and man: Is it the same disease? *Curr. Opin. Lipidol.*, 21: 434-440.
- Broeders, M.A., G.J. Tangelder, D.W. Slaaf, R.S. Reneman and M.G. Oude Egbrink, 2002. Hypercholesterolemia enhances thromboembolism in arterioles but not venules: Complete reversal by L-arginine. *Arterioscler. Thromb. Vasc. Biol.*, 22: 680-685.
- Bullock, B.C., T.B. Clarkson, N.D.M. Lehner, H.B. Lofland Jr. and R.W. St. Clair, 1969. Atherosclerosis in *Cebus albifrons* monkeys: III. Clinical and pathologic studies. *Exp. Mol. Pathol.*, 10: 39-62.
- Bustos, C., M.A. Hernandez-Presa, M. Ortego, J. Tunon and L. Ortega *et al.*, 1998. HMG-CoA reductase inhibition by atorvastatin reduces neointimal inflammation in a rabbit model of atherosclerosis. *J. Am. Coll. Cardiol.*, 32: 2057-2064.
- Casani, L., S. Sanchez-Gomez, G. Vilahur and L. Badimon, 2005. Pravastatin reduces thrombogenicity by mechanisms beyond plasma cholesterol lowering. *Thromb. Haemost.*, 94: 1035-1041.
- Clarkson, T.B., 1998. Nonhuman primate models of atherosclerosis. *Lab. Anim. Sci.*, 48: 567-569.
- Clarkson, T.B., M.G. Bond, B.C. Bullock, K.J. McLaughlin and J.K. Sawyer, 1984. A study of atherosclerosis regression in *Macaca mulatta*: V. Changes in abdominal aorta and carotid and coronary arteries from animals with atherosclerosis induced for 38 months and then regressed for 24 or 48 months at plasma cholesterol concentrations of 300 or 200 mg dL⁻¹. *Exp. Mol. Pathol.*, 41: 96-118.
- Clarkson, T.B., N.D.M. Lehner, W.D. Wagner, R.W. St. Clair, M.G. Bond and B.C. Bullock, 1979. A study of atherosclerosis regression in *Macaca mulatta*: I. Design of experiment and lesion induction. *Exp. Mol. Pathol.*, 30: 360-385.
- Corseaux, D., T. Meurice, I. Six, L. Rugeri and M.D. Ezekowitz *et al.*, 2000. Basic fibroblast growth factor increases tissue factor expression in circulating monocytes and in vascular wall. *Circulation*, 101: 2000-2006.
- Daugherty, A., E. Pure, D. Delfel-Butteiger, S. Chen, J. Leferovich, S.E. Roselaar and D.J. Rader, 1997. The effects of total lymphocyte deficiency on the extent of atherosclerosis in apolipoprotein E^{-/-} mice. *J. Clin. Invest.*, 100: 1575-1580.
- Davignon, J., 2005. Apolipoprotein E and atherosclerosis: Beyond lipid effect. *Arterioscler. Thromb. Vasc. Biol.*, 25: 267-269.
- Davis, H.R., D. Vesselinovitch and R.W. Wissler, 1984. Histochemical detection and quantification of macrophages in rhesus and cynomolgus monkey atherosclerotic lesions. *J. Histochem. Cytochem.*, 32: 1319-1327.
- Dong, Z.M., A.A. Brown and D.D. Wagner, 2000. Prominent role of P-selectin in the development of advanced atherosclerosis in ApoE-deficient mice. *Circulation*, 101: 2290-2295.
- Dornas, W.C., T.T.D. Oliveira, L.E.F. Augusto and T.J. Nagem, 2010. Experimental atherosclerosis in rabbits. *Arquivos Brasileiros Cardiologia*, 95: 272-278.
- Edelman, M.D., R. Elazer and M.D. Rogers, 1998. Pathobiologic responses to stenting. *Am. J. Cardiol.*, 81: 4E-6E.
- Finking, G. and H. Hanke, 1997. Nikolaj Nikolajewitsch Anitschkow (1885-1964) established the cholesterol-fed rabbit as a model for atherosclerosis research. *Atherosclerosis*, 135: 1-7.
- Fuster, V., J.T. Lie, L. Badimon, J.A. Rosemark, J.J. Badimon and E.J. Bowie, 1985. Spontaneous and diet-induced coronary atherosclerosis in normal swine and swine with von Willebrand disease. *Arterioscler. Thromb. Vasc. Biol.*, 5: 67-73.
- Fuster, V., L. Badimon, J.J. Badimon, J.H. Ip and J.H. Chesebro, 1991. The porcine model for the understanding of thrombogenesis and atherogenesis. *Mayo Clin. Proc.*, 66: 818-831.

- Gal, D., S.K. Chokshi, M. Mosseri, R.H. Clarke and J.M. Isner, 1992. Percutaneous delivery of low-level laser energy reverses histamine-induced spasm in atherosclerotic Yucatan Microswine. *Circulation*, 85: 756-768.
- Gerrity, R.G., R. Natarajan, J.L. Nadler and T. Kimsey, 2001. Diabetes-induced accelerated atherosclerosis in swine. *Diabetes*, 50: 1654-1665.
- Gershlick, A.H. and J. Baron, 1998. Dealing with in-stent restenosis. *Heart*, 79: 319-323.
- Grainger, D.J., J. Reckless and E. McKilligin, 2004. Apolipoprotein E modulates clearance of apoptotic bodies *in vitro* and *in vivo*, resulting in a systemic proinflammatory state in apolipoprotein E-deficient mice. *J. Immunol.*, 173: 6366-6375.
- Hartvigsen, K., C.J. Binder, L.F. Hansen, A. Rafia and J. Juliano *et al.*, 2007. A diet-induced hypercholesterolemic murine model to study atherogenesis without obesity and metabolic syndrome. *Arterioscler. Thromb. Vasc. Biol.*, 27: 878-885.
- Hofker, M.H., B.J.M. van Vlijmen and L.M. Havekes, 1998. Transgenic mouse models to study the role of APOE in hyperlipidemia and atherosclerosis. *Atherosclerosis*, 137: 1-11.
- Hu, W., P. Polinsky, E. Sadoun, M.E. Rosenfeld and S.M. Schwartz, 2005. Atherosclerotic lesions in the common coronary arteries of ApoE knockout mice. *Cardiovasc. Pathol.*, 14: 120-125.
- Johnstone, M.T., R.M. Botnar, A.S. Perez, R. Stewart, W.C. Quist, J.A. Hamilton and W.J. Manning, 2001. *In vivo* magnetic resonance imaging of experimental thrombosis in a rabbit model. *Arterioscler. Thromb. Vasc. Biol.*, 21: 1556-1560.
- Jorgensen, L., 2006. The role of platelets in the initial stages of atherosclerosis. *J. Thromb. Haemost.*, 4: 1443-1449.
- Knowles, J.W. and N. Maeda, 2000. Genetic modifiers of atherosclerosis in mice. *Arterioscler. Thromb. Vasc. Biol.*, 20: 2336-2345.
- Lee, R.T. and P. Libby, 1997. The unstable atheroma. *Arterioscler. Thromb. Vasc. Biol.*, 17: 1859-1867.
- Leppanen, P., J.S. Luoma, M.H. Hofker, L.M. Havekes and S.Y. Herttuala, 1998. Characterization of atherosclerotic lesions in apo E3-leiden transgenic mice. *Atherosclerosis*, 136: 147-152.
- Libby, P., 1995. Molecular bases of the acute coronary syndromes. *Circulation*, 91: 2844-2850.
- Linton, M.F. and S. Fazio, 1999. Macrophages, lipoprotein metabolism and atherosclerosis: Insights from murine bone marrow transplantation studies. *Curr. Opin. Lipidol.*, 10: 97-105.
- Linton, M.F., V.R. Babaev, L.A. Gleaves and S. Fazio, 1999. A direct role for the macrophage low density lipoprotein receptor in atherosclerotic lesion formation. *J. Biol. Chem.*, 274: 19204-19210.
- Lutgens, E., M. Daemen, M. Kockx, P. Doevendans and M. Hofker *et al.*, 1999. Atherosclerosis in APOE*3-Leiden transgenic mice: From proliferative to atheromatous stage. *Circulation*, 99: 276-283.
- Mann, G.V., S.B. Andrus, A. McNally and F.J. Stare, 1953. Experimental atherosclerosis in Cebus monkeys. *J. Exp. Med.*, 98: 195-218.
- Marzetta, C.A. and L.L. Rudel, 1986. A species comparison of low density lipoprotein heterogeneity in nonhuman primates fed atherogenic diets. *J. Lipid. Res.*, 27: 753-762.
- Mazzone, T. and C. Reardon, 1994. Expression of heterologous human apolipoprotein E by J774 macrophages enhances cholesterol efflux to HDL3. *J. Lipid. Res.*, 35: 1345-1353.
- Mohler III, E.R., L. Sarov-Blat, Y. Shi, D. Hamamdziec and A. Zalewski *et al.*, 2008. Site-specific atherogenic gene expression correlates with subsequent variable lesion development in coronary and peripheral vasculature. *Arterioscler. Thromb. Vasc. Biol.*, 28: 850-855.
- Mott, G.E., E.M. Jackson, C.A. McMahan and H.C. McGill Jr., 1992. Dietary cholesterol and type of fat differentially affect cholesterol metabolism and atherosclerosis in baboons. *J. Nutr.*, 122: 1397-1406.
- Nakashima, Y., A.S. Plump, E.W. Raines, J.L. Breslow and R. Ross, 1994. ApoE-deficient mice develop lesions of all phases of atherosclerosis throughout the arterial tree. *Arterioscler. Thromb. Vasc. Biol.*, 14: 133-140.
- Palazon, C.P., J. Alfon, P. Gaffney, M. Berrozpe, T. Royo and L. Badimon, 1998. Effects of reducing LDL and increasing HDL with gemfibrozil in experimental coronary lesion development and thrombotic risk. *Atherosclerosis*, 136: 333-345.
- Paszy, C., N. Maeda, J. Verstuyft and E.M. Rubin, 1994. Apolipoprotein AI transgene corrects apolipoprotein E deficiency-induced atherosclerosis in mice. *J. Clin. Invest.*, 94: 899-903.
- Paul, A., K.W.S. Ko, L. Li, V. Yechoor and M.A. McCrory *et al.*, 2004. C-reactive protein accelerates the progression of atherosclerosis in apolipoprotein E-deficient mice. *Circulation*, 109: 647-655.
- Pendse, A.A., J.M. Arbones-Mainar, L.A. Johnson, M.K. Altenburg and N. Maeda, 2009. Apolipoprotein E knockout and knock-in mice: Atherosclerosis, metabolic syndrome and beyond. *J. Lipid. Res.*, 50: S178-S182.

- Powell-Braxton, L., M. Veniant, R.D. Latvala, K.I. Hirano and W.B. Won *et al.*, 1998. A mouse model of human familial hypercholesterolemia: Markedly elevated low density lipoprotein cholesterol levels and severe atherosclerosis on a low-fat chow diet. *Nat. Med.*, 4: 934-938.
- Raffai, R.L., S.M. Loeb and K.H. Weisgraber, 2005. Apolipoprotein E promotes the regression of atherosclerosis independently of lowering plasma cholesterol levels. *Arterioscler. Thromb. Vasc. Biol.*, 25: 436-441.
- Rekhter, M.D., G.W. Hicks, D.W. Brammer, C.W. Work and J.S. Kim *et al.*, 1998. Animal Model that mimics atherosclerotic plaque rupture. *Circ. Res.*, 83: 705-713.
- Rekhter, M.D., G.W. Hicks, D.W. Brammer, H. Hallak and E. Kindt *et al.*, 2000. Hypercholesterolemia causes mechanical weakening of rabbit atheroma. *Circ. Res.*, 86: 101-108.
- Rensen, P.C.N. and L.M. Havekes, 2006. Cholesteryl ester transfer protein inhibition: Effect on reverse cholesterol transport? *Arterioscler. Thromb. Vasc. Biol.*, 26: 681-684.
- Rodgers, G.P., S.T. Minor, K. Robinson, D. Cromeens and S.C. Woolbert *et al.*, 1990. Adjuvant therapy for intracoronary stents. Investigations in atherosclerotic swine. *Circulation*, 82: 560-569.
- Rosenfeld, M.E., P. Polinsky, R. Virmani, K. Kauser, G. Rubanyi and S.M. Schwartz, 2000. Advanced atherosclerotic lesions in the innominate artery of the ApoE knockout mouse. *Arterioscler. Thromb. Vasc. Biol.*, 20: 2587-2592.
- Royo, T., J. Alfon, M. Berrozpe and L. Badimon, 2000. Effect of gemfibrozil on peripheral atherosclerosis and platelet activation in a pig model of hyperlipidemia. *Eur. J. Clin. Invest.*, 30: 843-852.
- Sanan, D.A., D.L. Newland, R. Tao, S. Marcovina and J. Wang *et al.*, 1998. Low density lipoprotein receptor-negative mice expressing human apolipoprotein B-100 develop complex atherosclerotic lesions on a chow diet: No accentuation by apolipoprotein(a). *Proc. Natl. Acad. Sci. USA.*, 95: 4544-4549.
- Scalia, R., M.E. Gooszen, S.P. Jones, M. Hoffmeyer and D.M. Rimmer III *et al.*, 2001. Simvastatin exerts both anti-inflammatory and cardioprotective effects in apolipoprotein E-deficient mice. *Circulation*, 103: 2598-2603.
- Shimizu, T.N., K. Kai, Y. Morimoto, M. Ishihara, H. Oishi, M. Kikuchi and H. Arai, 2009. Simple rabbit model of vulnerable atherosclerotic plaque. *Neurologia Medico-Chirurgica*, 49: 327-332.
- Shiomi, M., T. Ito, M. Shiraishi and Y. Watanabe, 1992. Inheritability of atherosclerosis and the role of lipoproteins as risk factors in the development of atherosclerosis in WHHL rabbits: Risk factors related to coronary atherosclerosis are different from those related to aortic atherosclerosis. *Atherosclerosis*, 96: 43-52.
- Shiomi, M., T. Ito, S. Yamada, S. Kawashima and J. Fan, 2004. Correlation of vulnerable coronary plaques to sudden cardiac events: Lessons from a myocardial infarction-prone Animal Model (the WHHLMI rabbit). *J. Atheroscler. Thromb.*, 11: 184-189.
- Singaraja, R.R., C. Fievet, G. Castro, E.R. James and N. Hennuyer *et al.*, 2002. Increased ABCA1 activity protects against atherosclerosis. *J. Clin. Invest.*, 110: 35-42.
- Skold, B.H., R. Getty and F.K. Ramsey, 1966. Spontaneous atherosclerosis in the arterial system of aging swine. *Am. J. Vet. Res.*, 27: 257-273.
- Suzuki, H., Y. Kurihara, M. Takeya, N. Kamada and M. Kataoka *et al.*, 1997. A role for macrophage scavenger receptors in atherosclerosis and susceptibility to infection. *Nature*, 386: 292-296.
- Swindle, M.M., 1998. Defining appropriate health status and management programs for specific-pathogen-free swine for xenotransplantation. *Ann. N. Y. Acad. Sci.*, 862: 111-120.
- Taylor, C.B., G.E. Cox, B.J. Hall-Taylor and L.G. Nelson, 1954. Atherosclerosis in areas of vascular injury in monkeys with mild hypercholesterolemia. *Circulation*, 10: 613-613.
- Taylor, C.B., G.E. Cox, P. Manalo-Estrella, J. Southworth and D.E. Patton *et al.*, 1962. Atherosclerosis in rhesus monkeys II. Arterial lesions associated with hypercholesterolemia induced by dietary fat and cholesterol. *Arch. Pathol.*, 74: 16-34.
- Teupser, D., A.D. Persky and J.L. Breslow, 2003. Induction of atherosclerosis by low-fat, semisynthetic diets in LDL receptor-deficient C57BL/6 and FVB/NJ mice: Comparison of lesions of the aortic root, brachiocephalic artery and whole aorta (en face measurement). *Arterioscler. Thromb. Vasc. Biol.*, 23: 1907-1913.
- Van Eck, M., M.P. De Winther, N. Herijgers, L.M. Havekes, M.H. Hofker, P.H. Groot and T.J. Van Berkel, 2000. Effect of human scavenger receptor class A overexpression in bone marrow-derived cells on cholesterol levels and atherosclerosis in ApoE-deficient mice. *Arterioscler. Thromb. Vasc. Biol.*, 20: 2600-2606.

- Van Vlijmen, B.J., A.M. van den Maagdenberg, M.J. Gijbels, H. van der Boom and H. HogenEsch *et al.*, 1994. Diet-induced hyperlipoproteinemia and atherosclerosis in apolipoprotein E3-Leiden transgenic mice. *J. Clin. Invest.*, 93: 1403-1410.
- Vesselinovitch, D., G.S. Getz, R.H. Hughes and R.W. Wissler, 1974. Atherosclerosis in the rhesus monkey fed three food fats. *Atherosclerosis*, 2: 303-321.
- Von der Thusen, J.H., B.J. van Vlijmen, R.C. Hoeber, M.M. Kockx, L.M. Havekes, T.J. van Berkel and E.A. Biessen, 2002. Induction of atherosclerotic plaque rupture in apolipoprotein E^{-/-} mice after adenovirus-mediated transfer of p53. *Circulation*, 105: 2064-2070.
- Westerterp, M., C.C. van der Hoogt, W. de Haan, E.H. Offerman and G.M. Dallinga-Thie *et al.*, 2006. Cholesteryl ester transfer protein decreases high-density lipoprotein and severely aggravates atherosclerosis in APOE*3-Leiden mice. *Arterioscler. Thromb. Vasc. Biol.*, 26: 2552-2559.
- White, F.C. and C.M. Bloor, 1992. Coronary vascular remodeling and coronary resistance during chronic ischemia. *Am. J. Cardiovasc. Pathol.*, 4: 193-202.
- White, F.C., S.M. Carroll, A. Magnet and C.M. Bloor, 1992. Coronary collateral development in swine after coronary artery occlusion. *Circ. Res.*, 71: 1490-1500.
- Wilensky, R.L., Y. Shi, E.R. III Mohler, D. Hamamdžić and M.E. Burgert *et al.*, 2008. Inhibition of lipoprotein-associated phospholipase A₂ reduces complex coronary atherosclerotic plaque development. *Nat. Med.*, 14: 1059-1066.
- Wolfe, M.S., J.K. Sawyer, T.M. Morgan, B.C. Bullock and L.L. Rudel, 1994. Dietary polyunsaturated fat decreases coronary artery atherosclerosis in a pediatric-aged population of African green monkeys. *Arterioscler. Thromb.*, 4: 587-597.
- Zadelaar, S., R. Kleemann, L. Verschuren, J. de Vries-Van der Weij, J. van der Hoorn, H.M. Princen and T. Kooistra, 2007. Mouse models for atherosclerosis and pharmaceutical modifiers. *Arterioscler. Thromb. Vasc. Biol.*, 27: 1706-1721.
- Zhang, W., P.G. Yancey, Y.R. Su, V.R. Babaev, Y. Zhang, S. Fazio and M.F. Linton, 2003. Inactivation of macrophage scavenger receptor class B type I promotes atherosclerotic lesion development in apolipoprotein E-deficient mice. *Circulation*, 108: 2258-2263.