

Effect of *Lactobacillus delbrueckii* on Jejunum Innate Immune-Related Gene Expression in Mice

^{1,2}Jie-Lin Duan, ³Shao-Ping He, ³Guan Yang, ^{1,2}Jie Yin, ¹Wen-Kai Ren, ³Jian-Wen Deng, ³Yu Gong, ³Feng-Ming Chen, ¹Tie-Jun Li and ³Xing-Guo Huang

¹Key Laboratory of Agro-Ecological Processes in Subtropical Region, Chinese Academy of Sciences, Hunan Provincial Engineering Research Center of Healthy Livestock, Institute of Subtropical Agriculture, Scientific Observing and Experimental Station of Animal Nutrition and Feed Science in South-Central, Ministry of Agriculture, Changsha, 410125 Hunan, China

²University of Chinese Academy of Science, 100049 Beijing, China

³College of Animal Science and Technology, Hunan Agricultural University, Changsha, 410128 Hunan, China

Abstract: Probiotics, regarded as live microorganisms, play important roles in regulating intestinal inflammatory responses and innate immunity. Although, intestinal immunomodulation of probiotics have been enormously conducted into data, the effects of *Lact. delbrueckii* on jejunum innate immunity are rarely reported. This study aimed to investigate the effects of *Lact. delbrueckii* on serum profiles, jejunum inflammatory cytokines and innate immune-related genes expression in mouse. A total of 32 Kunming male mice were randomly divided into four groups and each group were orally administrated with saline normal, living *Lact. delbrueckii*, heat-killed *Lact. delbrueckii* and Spent Culture Supernatant (SCS). Serum profiles (IL-2, IgG) and jejunum immune-related gene expression were detected including inflammatory cytokines (INF- γ , TNF- α , IL-17, IL-13) peneth cell-sepcific gene (cryptdin 1, 4, 5, lysozyme C, CRS-1C, CRS-4C and sPLA2) and goblet cell-specific gene (mucin-2 and mucin-4). Compared with control group, orally administrated SCS significantly decreased ($p < 0.05$) ADG of mice. Orally administrated living and heat-killed *Lact. delbrueckii* group significantly increased ($p < 0.05$) serum IL-2 level and while SCS remarkably decreased ($p < 0.05$) serum IgG level. Meanwhile, oral administration of living *Lact. delbrueckii* significantly enhanced ($p < 0.05$) jejunum inflammatory cytokines (INF- γ , TNF- α , IL-17) expression. SCS significantly increased ($p < 0.05$) peneth cell-sepcific gene (cryptdin 1, 4, 5 and sPLA2) gene expression. In conclusion, oral administration of *Lact. delbrueckii* affects serum profiles and jejunum innate immune-related gene expression and thereby modulates jejunum innate immunity.

Key words: *Lactobacillus delbrueckii*, jejunum, peneth cell, innate immune, CRS-1C

INTRODUCTION

Lactobacillus delbrueckii (*Lact. delbrueckii*) is a gram-positive bacteria belonging to lactic acid bacteria and conventionally has been used to ferment milk because of its great production of lactic acid that give yogurt its flavor and textural properties (Moller *et al.*, 2007). Recently, many compelling studies have reported *Lact. delbrueckii* has potential probiotic function (Fernandez *et al.*, 2005). Abedi *et al.* (2013) have reported that *Lact. delbrueckii* exhibits remarked anti-bacterial and anti-adhesion properties against *Escherichia coli* (*E. coli*) *in vitro* (Abedi *et al.*, 2013) suggesting that *Lact. delbrueckii* may be able to prevent *E. coli* or other pathogens infection in the gut. Indeed, Dos Santos *et al.* (2011) have demonstrated this point that

Lact. delbrueckii UFV-H2b20 protects mice from death caused by *Listeria monocyto* genes (*L. monocytogenes*) infection (Dos Santos *et al.*, 2011). Besides anti-bacterial and anti-adhesion properties, numerous well-designed experiments have exemplified that *Lact. delbrueckii* exhibits versatile beneficial roles inmodulating intestinal microbiota (Mori *et al.*, 2011) enhancing intestinal barrier function (Yu *et al.*, 2012) and regulating intestinal immune response through modulation of Nuclear Factor kappa B (NF- κ B) and Mitogen-Activated Protein Kinases (MAPKs) signaling pathway (Hegazy and El-Bedewy, 2010; Thomas and Versalovic, 2010). Additionally, *Lact. delbrueckii* enable to facilitate activation of intestinal immune system, increase number of intestinal T cell and exhibits anti-inflammation properties (Picchiatti *et al.*, 2009; Del Carmen *et al.*, 2014).

Interestingly, orally administrated *Lact. delbrueckii* but also exhibits anti-aging ability by slowing the aging of the T cell and elevates antimicrobial peptide human Beta-Defensin-2 (hBD2) expression (Moro-Garcia *et al.*, 2013). Moreover, *Lact. delbrueckii* has been used to treat children with celiac disease (Di Cagno *et al.*, 2009) type 2 diabetes (Honda *et al.*, 2012) and Inflammation Bowel Disease (IBD) (Hegazy and El-Bedewy, 2010; Sengul *et al.*, 2011) and experimental autoimmune encephalomyelitis (Lavasani *et al.*, 2010). Considering versatile beneficial roles of *Lact. delbrueckii* it is plausible to speculate a possible role of *Lact. delbrueckii* in modulating jejunum innate immunity. However, directly evidence in favor of this hypothesis is missing.

The mucosa of the gastrointestinal tract is in directly contact with intestinal lumen contents including bacterial and antigens. Being an important harbor for microbes, the orchestrated intestinal innate immunity that protects host against pathogenic bacterial (Oswald, 2006; Ganz and Szabo, 2013; Vivier and Malissen, 2005) has been highlighted. Under normal or abnormal conditions, Pattern Recognition Receptors (PRRs) like Toll-Like Receptors (TLRs) recognize Pathogen Associated Molecular Pattern (PAMP) and then trigger TLR-MyD88 signaling pathway, leading secretion of inflammatory cytokines which perform crucial roles in regulating intestinal homeostasis (Duan *et al.*, 2014; Menendez *et al.*, 2013). Paneth cells that populate the crypts through the small intestine exert pivotal roles in intestinal innate immunity because of specific secreted antimicrobial peptides including α -defensins, lysozyme C and Cryptdin-Related Sequence (CRS) peptides (Selsted and Ouellette, 2005; Ouellette, 2004) which exhibit broad spectrum microbicides against gram negative and gram positive microbes by membrane-disruptive and inducing rapid microbial cell K^+ efflux (Shanahan *et al.*, 2010; Santaolalla *et al.*, 2011). In addition, mucins from goblet cell are indispensable components of intestinal mucus layer that separate host from lumen pathogens (Boonzaier *et al.*, 2013; Johansson *et al.*, 2008). Many investigations have shown that mucins from goblet cells play important roles in intestinal innate immunity by enhancing gut barrier function (Zarepour *et al.*, 2013) thus protecting host against pathogens invasion. Although, literatures covering probiotics on intestinal innate immunity have been largely documented into data, the effect of *Lact. delbrueckii* on paneth cell and goblet cell-specific molecules and inflammatory cytokines expression in jejunum remains rarely reported. For these reasons, researchers conducted the experiment to study the effect of *L. delbrueckii* on jejunum innate immunity including serum IgG and IL-2 concentration, expression of inflammation gene, paneth cell-specific gene and goblet cell-specific gene in jejunum in mouse.

MATERIALS AND METHODS

Preparation and culture of *Lact. delbrueckii*:

Lact. delbrueckii was obtained from laboratory of College of Animal Science and Technology, Hunan Agriculture University. For stimulation experiment, the bacteria were anaerobically cultured for 48 h at 37°C in de Man, Rogosa and Sharp broth (MRS broth) in an anaerobic condition prior to use. *Lact. delbrueckii* lysate were harvested by centrifugation (4,000x, 10 min, 4°C) for spent culture supernatant SCS and stored at 4°C for further use. The bacterial cells in the stationary growth phase were harvested by sterile saline suspension, centrifuged (4000xg, 10 min, 4°C) washed 3 times with sterile saline and subsequently adjusted to a final concentration of 1×10^9 CFU in sterile saline. Heat-killed *Lact. delbrueckii* was obtained by maintaining cells at 56°C for 60 min for inactivation and then reaffirmed whether live *L. delbrueckii* existed by MRS plate.

Experimental design: A total of 32 Kunming male mice were obtained from the animal Laboratory Animal Center of Central South University (Hunan, China). The mice (20±2 g) were randomly assigned to four treatment groups with 8 replicates per treatment. The mice were housed in a friendly and environmentally controlled pathogen-free colony and had access to standard rodent food and water *ad libitum*. After 3 days of adaptive feeding, the four groups of mice were orally administrated with saline normal, live *Lact. delbrueckii* (1×10^9 CFU) heat-killed *Lact. delbrueckii* (1×10^9 CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse. All of the animals had free access to diets and drinking water. After 7 days, all animals were sacrificed for sample collection. All animal care, handling and surgical techniques followed protocols approved by the Hunan Agricultural University Animal Care and Use Committee before study initiation.

Average Daily Gain (ADG): All mice were weighed individually on 0, 7 days of experiment to measure body weight. The ADG (g/mouse/day) during the experimental stages was calculated as the difference between the initial and final weight.

Serum Immunoglobulin G (IgG) and Interleukin-2 (IL-2) determination:

The concentration of serum Immunoglobulin G (IgG) and Interleukin-2 (IL-2) were measured measured using ELISA kit in accordance with the manufacturer's instructions (Cusabio biotech Co., Ltd. China).

Total RNA extraction and cDNA synthesis: Total RNA was isolated from liquid nitrogen frozen jejunum tissues with TRIZOL reagent (Invitrogen, USA) and then treated

with DNase I (Invitrogen, USA) according to the manufacturer's instructions. For each sample, the RNA quality was checked by 1% agarose gel electrophoresis, stained with 10 µg mL⁻¹ ethidium bromide. Synthesis of the first strand cDNA was performed with oligo (dT) 20 and Superscript II reverse transcriptase (Invitrogen, USA).

Quantification mRNA by real-time PCR analysis: Primers used in this study were presented in previous study (Ren *et al.*, 2014). β-actin was used as an house keeping gene to normalize target gene transcript levels. Real-time PCR was performed according the previous study (Ren *et al.*, 2013). Relative level of gene expression was normalized and expressed as a ratio of target gene expression to the control group.

Statistical analysis: The results were expressed as mean±Standard Error of the Mean (SEM). All statistical evaluation was performed by using SPSS 17.0 Software. Group comparisons for statistical difference were performed using the one-ANOVA's Duncan (D)-test. Differences (p<0.05) were considered significant.

RESULTS

Effect of oral administration of *Lact. delbrueckii* on mouse ADG: Although, the growth promotion of Lactobacillus have been extensively reported, it remains exist variable results with respect to body weight. Interestingly, Bai *et al.* (2013) have reported that dietary supplementation of probiotics can augment ADG of broiler chickens but the research from Fajardo *et al.* (2012) indicate that probiotic-supplemented exhibited no significant effect on Body Weight Gain (BWG) in the present experiment, orally administrated living and heat-killed *Lact. delbrueckii* exhibited no difference (p>0.05) on mouse ADG. However, SCS significantly decreased (p<0.05) mouse ADG (Table 1).

Table 1: Effect of *Lact. delbrueckii* on Average Daily Gain (ADG) in mice

| Catalogue | Initial weight | Final weight | ADG (g day ⁻¹) |
|-----------|----------------|-------------------------|----------------------------|
| Control | 24.90±0.45 | 31.47±1.00 ^a | 0.72±0.06 ^b |
| Live | 25.00±0.79 | 31.05±1.24 ^a | 0.67±0.07 ^b |
| Heat-kill | 24.98±0.51 | 32.14±1.12 ^a | 0.78±0.74 ^a |
| SCS | 24.98±0.59 | 29.18±0.91 ^b | 0.47±0.04 ^c |

The values are expressed as means±MES, n = 8 and the values with different superscripts are significant (p<0.05) while values with same superscripts are not significant different (p>0.05). All groups of mice were administrated orally respectively with saline normal, live *Lact. delbrueckii* (1×10⁹ CFU) heat-killed *Lact. delbrueckii* (1×10⁹ CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse

Effect of oral administration of *Lact. delbrueckii* on serum IgG and IL-2 concentration: Immunoglobulin G (IgG) and Interleukin-2 (IL-2) indispensable components in immunologic response, play important roles in preventing pathogens from invading. In the study, compared with control group, administration of living and heat-killed *Lact. delbrueckii* exhibited no remarkable difference (p>0.05) on serum IgG level but SCS remarkably decreased (p<0.05) serum IgG level (Fig. 1). Meanwhile, oral administration of living and heat-killed *Lact. delbrueckii* significantly elevated (p<0.05) level of serum IL-2 while there was no significant difference (p>0.05) between SCS and control group (Fig. 2).

Effect of oral administration of *Lact. delbrueckii* on cytokines expression in jejunum: Intestinal inflammatory cytokines that are secreted by immune cells, e.g., lymphocyte, dendritic cell play central roles in intestinal inflammation immunologic response and the expression of cytokines in jejunum is affected by various factors including commensal microbiota (Wang *et al.*, 2013). Thus, researchers also detected inflammatory cytokines (INF-γ, TNF-α, IL-13, IL-17) expression in jejunum by RT-PCR. In the study, compared with control group, living *Lact. delbrueckii* significantly augmented (p<0.05) the mRNA abundance of TNF-α, INF-γ and IL-17. However, heat-killed *Lact. delbrueckii* and SCS exhibited no




Fig. 1: Effect of *Lact. delbrueckii* on serum IgG concentration in mice. The values are expressed as means±MES, n = 8 and *p<0.05. All groups of mice were administrated orally, respectively with saline normal, living *Lact. delbrueckii* (1×10⁹ CFU) heat-killed *Lact. delbrueckii* (1×10⁹ CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse

significant ($p > 0.05$) effects on these gene expression excepting heat-killed *Lact. delbrueckii* remarked increased ($p < 0.05$) mRNA but but abundance of IL-17 (Fig. 3).

Effect of oral administration of *Lact. delbrueckii* on paneth cell and goblet cell function in jejunum: To investigate the effect of *Lact. delbrueckii* on jejunum innate immunity, researchers examined jejunum paneth




Fig. 2: Effect of *Lact. delbrueckii* on serum IL-2 concentration in mice. The values are expressed as means±MES, n = 8 and * $p < 0.05$. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10^9 CFU) heat-killed *Lact. delbrueckii* (1×10^9 CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse




Fig. 3: Effect of *Lact. delbrueckii* on cytokines gene expression in jejunum in mice. The values are expressed as means±MES, n = 8 and * $p < 0.05$. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10^9 CFU) heat-killed *Lact. delbrueckii* (1×10^9 CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse

cell-specific molecules (cryptdin-1, 4, 5, lysozyme C, sPLA2 but and CRS-1C, CRS-4C) and goblet cell-specific molecules (mucin-2 and mucin-4). In the study, compared with control group, SCS remarkably increased ($p < 0.05$) mRNA abundance of cryptdin-1, 4 and 5 and sPLA2 (Fig. 4 and 5). Moreover, apart from cryptdin-5 in heat-




Fig. 4: Effect of *Lact. delbrueckii* on criptdins expression in jejunum in mice. The values are expressed as means±MES, n = 8 and * $p < 0.05$. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10^9 CFU) heat-killed *Lact. delbrueckii* (1×10^9 CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse




Fig. 5: Effect of *Lact. delbrueckii* on sPLA2 and lysozyme C expression in jejunum in mice. The values are expressed as means±MES, n = 8 and * $p < 0.05$. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10^9 CFU) heat-killed *Lact. delbrueckii* (1×10^9 CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse




Fig. 6: Effect of *Lact. delbrueckii* on CRS-1C and CRS-4C expression in jejunum in mice. The values are expressed as means±MES, n = 8 and *p<0.05. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10⁹ CFU) heat-killed *Lact. delbrueckii* (1×10⁹ CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse




Fig. 7: Effect of *Lact. delbrueckii* on mucin 2 and 4 expression in jejunum in mice. The values are expressed as means±MES, n = 8 and *p<0.05. All groups of mice were administrated orally, respectively with saline normal, live *Lact. delbrueckii* (1×10⁹ CFU) heat-killed *Lact. delbrueckii* (1×10⁹ CFU) or Spent Culture Supernatant (SCS) once daily 0.5 mL per mouse

killed *Lact. delbrueckii* group, the mRNA abundance of cryptdin-1, 4, 5 and sPLA2 in SCS group were significant higher (p<0.05) than that in living and heat-killed *Lact. delbrueckii* group. However, all treatments exhibited no significant (p>0.05) effects on lysozyme C, CRS-1C and CRS-4 C expression (Fig. 5 and 6). In addition, the abundance of mucin 2 mRNA in living and heat-killed *Lact. delbrueckii* group was significantly lower (p<0.05) than that in SCS group (Fig. 7) but all treatment exhibited no effect (p>0.05) on mucin 4 (Fig. 7) expression.

Lact. delbrueckii that has been widely used to ferment milk is an important probiotic bacterial and recently has been used to treat some diseases because of its immunomodulation. In recent years, growing focuses have been located at immunomodulation of *Lact. delbrueckii*. Although, increasing numbers of research literatures have shown that *Lact. delbrueckii* enable to enhance intestinal barrier function, balance intestinal immune response and regulate intestinal homeostasis (Valdovinos-Diaz, 2013; Persborn *et al.*, 2013; Yang and Sheu, 2012; Howarth and Wang, 2013) the impact of *Lact. delbrueckii* on jejunum innate immunity remains obscurity and needs to be investigated. In the study, oral administration of *Lact. delbrueckii* increased serum IL-2 concentration and stimulated inflammatory cytokines expression and affected paneth cell-specific molecules and goblet cell-specific molecules expression in jejunum in mice.

Although, the growth promotion effects of Lactobacillus have been extensively reported, it remains exist variable results. Interestingly, Bai *et al.* (2013) have reported that dietary supplementation of probiotics augment ADG of broiler chickens but the research from Fajardo *et al.* (2012) indicates that probiotic-supplemented have no difference effect on Body Weight Gain (BWG) (Fajardo *et al.*, 2012). In the current research, living and heat-killed *Lact. delbrueckii* exhibited no significant effect on mice ADG, corresponding with Giang *et al.* (2010) report that supplementation of lactic acid bacterial has no effect on performance of weaning piglets. However, there existed other researchs suggesting that probiotics may be improve animal growth performance since supplementation with probiotic bacterial enable to enhance intestinal function and modulate intestinal microbial composition which are important for nutrients absorption (Wang *et al.*, 2012). In addition, growth performance effects of probiotics link to strains and dosage of probiotics. Mountzouris *et al.* (2010) have reported that probiotic inclusion levels exhibit a significant effect on broiler growth performance. According to these factors in the study, the differences with other research may be link to the effect of *Lact. delbrueckii* on intestinal bacterial composition and also may be related to strains or dosage of *Lact. delbrueckii*.

The mucosal membrane of the gastrointestinal tract is the first barrier against foreign antigens such as natural

toxins, pathogenic microflora (Oswald, 2006). Thus, intestinal innate immunity is orchestrated to protect host against pathogens invasion, guaranteeing to sustain intestinal homeostasis (Santaolalla *et al.*, 2011). Intestinal inflammatory cytokines that are secreted by activated T lymphocytes, macrophages but and dendritic cell, plays critical roles in intestinal innate immunity (Wang *et al.*, 2013; Heneghan *et al.*, 2014). Many studies have shown that supplementation with *Lact. delbrueckii* modulate intestinal innate immunity (Tohno *et al.*, 2011) and remarkably stimulate intestinal cytokines production and balance Th1 and Th2 immune response in health or pathological model (Hegazy and El-Bedewy, 2010; Rocha *et al.*, 2014; Elmadfa *et al.*, 2010) through the NF- κ B and MAPK signaling pathways (Thomas and Versalovic, 2010). In this study, administration with living *Lact. delbrueckii* elevate jejunal TNF- α , INF- γ and IL-17 expression, suggesting that orally administered *Lact. delbrueckii* may be able to induce jejunal Th1-immune response and jejunal inflammatory response. Similarly, many compelling studies also have demonstrated that *Lact. delbrueckii* mediates Th1 cytokines production and stimulate inflammatory response *in vivo* and *in vitro* (Castanheira *et al.*, 2007; Neumann *et al.*, 2009). Moreover, elevated serum IL-2 concentration in living and heat-killed *Lact. delbrueckii* further confirms this point because IL-2 signaling is involved in proliferation of T lymphocytes cell and Th1 development (Fujimura *et al.*, 2013). These results may be link to TLRs-MyD88 mediate various signaling pathway because TLRs-blocking antibodies reduce cytokines production induced by *Lact. delbrueckii* (Cai *et al.*, 2010). However, some studies also have reported that administration with *Lact. delbrueckii* significantly inhibits intestinal pro-inflammatory or inflammatory cytokines production, especially in pathological models such as Dextran Sulfate (DSS) induced colitis (Hegazy and El-Bedewy, 2010; Santos Rocha *et al.*, 2012). These different results may be linked to different experimental model and experimental conditions. However, to elucidate this point, further studies need to be carried out.

Moreover, antimicrobial peptides from paneth cells and mucins from goblet cells play pivotal roles in intestinal innate immunity. Many studies have shown that paneth cells secrete specific antimicrobial peptides into intestinal lumen (Selsted and Ouellette, 2005; Ouellette, 2004; Zhao and Lu, 2014) and exhibits broad spectrum microbicides by membrane-disruptive (Santaolalla *et al.*, 2011; Boonzaier *et al.*, 2013) therefore protecting host against microbe invasion. In

addition, mucins from goblet cell are important components of intestinal mucus layer and play important roles in intestinal innate immunity by enhancing gut barrier function (Johansson *et al.*, 2008; Zarepour *et al.*, 2013). Many well-designed experiments have shown that probiotics affect intestinal α -defensins, CRS-4C and mucins expression in human and mouse (Underwood *et al.*, 2012; Schlee *et al.*, 2008; Mondel *et al.*, 2009). Indeed, in the current study, orally administered SCS changed jejunal cryptdin-1, 4 and 5 and sPLA2 expression. However, living and heat-killed *Lact. delbrueckii* exhibited no significant effects on antimicrobial peptide and mucins expression which were different with Moro-Garcia *et al.* (2013) research that *Lact. delbrueckii* could increase human beta defensin-2 expression. Meanwhile, Wehkamp *et al.* (2004) also have demonstrated that probiotic, *Escherichia coli* Nissle 1917, up-regulates human beta defensin-2 through activation of NF- κ B and AP-1. As a result, researchers speculate that the different results may be ascribed to bioactive metabolites of *Lact. delbrueckii* such as hydrogen peroxide and lactic acid largely produced by *Lact. delbrueckii* (Batdorj *et al.*, 2007; Strus *et al.*, 2009). In addition, gut microbial composition also affects intestinal antimicrobial peptide and mucins expression (Salzman *et al.*, 2007). Thus, immunomodulation of *Lact. delbrueckii* on jejunal paneth cells and goblet cell function may link to the effect of *Lact. delbrueckii* on intestinal microbial composition (Hemarajata and Versalovic, 2013). Thus, to elucidate this point many researches should be carried out.

Intriguingly, in this study, different treatment with living, heat-killed *Lact. delbrueckii* and SCS exhibited different effects on jejunal inflammatory cytokines and paneth cell-specific molecules expression. Although, living *Lact. delbrueckii* induced remarkable inflammatory response, heat-killed *Lact. delbrueckii* and SCS exhibited no significant effects on inflammatory cytokines expression. These interesting results may link to the effects of *Lact. delbrueckii* on intestinal microbial composition and the bacteria-mediated fecal metabolites such as Short-Chain Fatty Acids (SCFAs) (Mori *et al.*, 2011) which play important roles in regulating differentiation of intestinal T cell population (Furusawa *et al.*, 2013). Additionally, SCS significantly increased paneth cell-specific antimicrobial peptide expression but other treatment failed to up-regulate these genes expression. With regard to these intriguing results, researchers speculate the possible reason is that bioactive metabolites such as hydrogen peroxide and lactic

acid largely produced by *Lact. delbrueckii* (Strus *et al.*, 2009). However, many researches should be conducted to elucidate this point.

CONCLUSION

Researchers found that oral administration of *Lact. delbrueckii* affects serum IL-2 levels, stimulate jejunum inflammatory cytokines expression and affect jejunum paneth cell-specific molecules and goblet cell-specific molecules expression in mice, suggesting that *Lact. delbrueckii* may be able to regulate jejunum innate immunity.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (No. 31272458, 31372322, 31272463) National Basic Research Program of China (2013CB127301) and Strategical Scientific and Technological Research projects of emerging industries in Hunan Province (2011GK4061).

REFERENCES

- Abedi, D., S. Feizizadeh, V. Akbari and A. Jafarian-Dehkordi, 2013. *In vitro* anti-bacterial and anti-adherence effects of *Lactobacillus delbrueckii* subsp. *bulgaricus* on *Escherichia coli*. Res. Pharm. Sci., 8: 260-208.
- Bai, S.P., A.M. Wu, X.M. Ding, Y. Lei, J. Bai, K.Y. Zhang and J.S. Chio, 2013. Effects of probiotic-supplemented diets on growth performance and intestinal immune characteristics of broiler chickens. Poultry Sci., 92: 663-670.
- Batdorj, B., V. Trinetta, M. Dalgalarondo, H. Prevost and X. Dousset *et al.*, 2007. Isolation, taxonomic identification and hydrogen peroxide production by *Lactobacillus delbrueckii* subsp. *lactis* T31, isolated from Mongolian yoghurt: Inhibitory activity on food-borne pathogens. J. Applied Microbiol., 103: 584-593.
- Boonzaier, J., E.L. Van der Merwe, N.C. Bennett and S.H. Kotze, 2013. A comparative histochemical study of the distribution of mucins in the gastrointestinal tracts of three insectivorous mammals. Acta Histochemica, 115: 549-556.
- Cai, S., B.H. Bay, Y.K. Lee, J. Lu and R. Mahendran, 2010. Live and lyophilized *Lactobacillus* species elicit differential immunomodulatory effects on immune cells. FEMS Microbiol. Lett., 302: 189-196.
- Castanheira, L.G., J.M. Castro, O.A. Martins-Filho, J.R. Nicoli, L.Q. Vieira and L.C. Afonso, 2007. *Lactobacillus delbrueckii* as a potential skin adjuvant for induction of type 1 immune responses. Front Biosci., 12: 1300-1307.
- Del Carmen, S., A.D.M. de LeBlanc, R. Martin, F. Chain, P. Langella, L.G. Bermudez-Humaran and J.G. LeBlanc, 2014. Genetically engineered immunomodulatory *Streptococcus thermophilus* strains producing antioxidant enzymes exhibit enhanced anti-inflammatory activities. Applied Environ. Microbiol., 80: 869-877.
- Di Cagno, R., C.G. Rizzello, F. Gagliardi, P. Ricciuti and M. Ndagijimana *et al.*, 2009. Different fecal microbiotas and volatile organic compounds in treated and untreated children with celiac disease. Applied Environ. Microbiol., 75: 3963-3971.
- Dos Santos, L.M., M.M. Santos, H.P. de Souza Silva, R.M.E. Arantes, J.R. Nicoli and L.Q. Vieira, 2011. Monoassociation with probiotic *Lactobacillus delbrueckii* UFV-H2b20 stimulates the immune system and protects germfree mice against *Listeria monocytogenes* infection. Med. Microbiol. Immunol., 200: 29-38.
- Duan, D., S. Zhang, X. Li, H. Guo and M. Chen *et al.*, 2014. Activation of the TLR/MyD88/NF- κ B signal pathway contributes to changes in IL-4 and IL-12 production in piglet lymphocytes infected with porcine circovirus type 2 *in vitro*. PLoS One, Vol. 9. 10.1371/journal.pone.0097653.
- Elmadfa, I., P. Klein and A.L. Meyer, 2010. Immune-stimulating effects of lactic acid bacteria *in vivo* and *in vitro*. Proc. Nutr. Soc., 69: 416-420.
- Fajardo, P., L. Pastrana, J. Mendez, I. Rodriguez, C. Fucinos and N.P. Guerra, 2012. Effects of feeding of two potentially probiotic preparations from lactic acid bacteria on the performance and faecal microflora of broiler chickens. Sci. World J. 10.1100/2012/562635.
- Fernandez, M.F., S. Boris and C. Barbes, 2005. Safety evaluation of *Lactobacillus delbrueckii* subsp. *lactis* UO 004, a probiotic bacterium. Res. Microbiol., 156: 154-160.
- Fujimura, K., A. Oyamada, Y. Iwamoto, Y. Yoshikai and H. Yamada, 2013. CD4 T cell-intrinsic IL-2 signaling differentially affects Th1 and Th17 development. J. Leukoc Biol., 94: 271-279.
- Furusawa, Y., Y. Obata, S. Fukuda, T.A. Endo and G. Nakato *et al.*, 2013. Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. Nature, 504: 446-450.

- Ganz, M. and G. Szabo, 2013. Immune and inflammatory pathways in NASH. *Hepato. Int.*, 7: 771-781.
- Giang, H.H., T.Q. Viet, B. Ogle and J.E. Lindberg, 2010. Effects of different probiotic complexes of lactic acid bacteria on growth performance and gut environment of weaned piglets. *Livest. Sci.*, 133: 182-184.
- Hegazy, S.K. and M.M. El-Bedewy, 2010. Effect of probiotics on pro-inflammatory cytokines and NF- κ B activation in ulcerative colitis. *World J. Gastroenterol.*, 16: 4145-4151.
- Hemarajata, P. and J. Versalovic, 2013. Effects of probiotics on gut microbiota: Mechanisms of intestinal: Immunomodulation and neuromodulation. *Therap. Adv. Gastroenterol.*, 6: 39-51.
- Heneghan, A.F., J.F. Pierre, A. Gosain and K.A. Kudsk, 2014. IL-25 improves luminal innate immunity and barrier function during parenteral nutrition. *Ann. Surg.*, 259: 394-400.
- Honda, K., M. Moto, N. Uchida, F. He and N. Hashizume, 2012. Anti-diabetic effects of lactic acid bacteria in normal and type 2 diabetic mice. *J. Clin. Biochem. Nutr.*, 51: 96-101.
- Howarth, G.S. and H. Wang, 2013. Role of endogenous microbiota, probiotics and their biological products in human health. *Nutrients*, 5: 58-81.
- Johansson, M.E., M. Phillipson, J. Petersson, A. Velcich, L. Holm and G.C. Hansson, 2008. The inner of the two Muc2 mucin-dependent mucus layers in colon is devoid of bacteria. *Proc. Natl. Acad. Sci. USA.*, 105: 15064-15069.
- Lavasani, S., B. Dzhambazov, M. Nouri, F. Fak and S. Buske *et al.*, 2010. A novel probiotic mixture exerts a therapeutic effect on experimental autoimmune encephalomyelitis mediated by IL-10 producing regulatory T cells. *PLoS One*, Vol. 5. 10.1371/journal.pone.0009009.
- Menendez, A., B.P. Willing, M. Montero, M. Wlodarska and C. So *et al.*, 2013. Bacterial stimulation of the TLR-MyD88 pathway modulates the homeostatic expression of ileal paneth cell α -defensins. *J. Innate Immun.*, 5: 39-49.
- Moller, C., W. Bockelmann, A. Ammann and K.J. Heller, 2007. Production of yoghurt with mild taste by a *Lactobacillus delbrueckii* subsp. *Bulgaricus* mutant with altered proteolytic properties. *Biotechnol. J.*, 2: 469-479.
- Mondel, M., B.O. Schroeder, K. Zimmermann, H. Huber and S. Nuding *et al.*, 2009. Probiotic *E. coli* treatment mediates antimicrobial human β -defensin synthesis and fecal excretion in humans. *Mucosal Immunol.*, 2: 166-172.
- Mori, K., T. Ito, H. Miyamoto, M. Ozawa, S. Wada and Y. Kumagai *et al.*, 2011. Oral administration of multispecies microbial supplements to sows influences the composition of gut microbiota and fecal organic acids in their post-weaned piglets. *J. Biosci. Bioeng.*, 112: 145-150.
- Moro-Garcia, M.A., R. Alonso-Arias, M. Baltadjieva, C.F. Benitez and M.A.F. Barrial *et al.*, 2013. Oral supplementation with *Lactobacillus delbrueckii* subsp. *bulgaricus* 8481 enhances systemic immunity in elderly subjects. *Age*, 35: 1311-1326.
- Mountzouris, K.C., P. Tsitsrikos, I. Palamidi, A. Arvaniti, M. Mohnl, G. Schatzmayr and K. Fegeros, 2010. Effects of probiotic inclusion levels in broiler nutrition on growth performance, nutrient digestibility, plasma immunoglobulins and cecal microflora composition. *Poult. Sci.*, 89: 58-67.
- Neumann, E., M.G. Ramos, L.M. Santos, A.C.P. Rodrigues and E.C. Vieira *et al.*, 2009. *Lactobacillus delbrueckii* UFV-H2b20 induces type 1 cytokine production by mouse cells *in vitro* and *in vivo*. *Braz. J. Med. Biol. Res.*, 42: 358-367.
- Oswald, I.P., 2006. Role of intestinal epithelial cells in the innate immune defence of the pig intestine. *Vet. Res.*, 37: 359-368.
- Ouellette, A.J., 2004. Defensin-mediated innate immunity in the small intestine. *Best Pract. Res. Clin. Gastroenterol.*, 18: 405-419.
- Persborn, M., J. Gerritsen, C. Wallon, A. Carlsson, L.M.A. Akkermans and J.D. Soderholm, 2013. The effects of probiotics on barrier function and mucosal pouch microbiota during maintenance treatment for severe pouchitis in patients with ulcerative colitis. *Aliment. Pharmacol. Ther.*, 38: 772-783.
- Picchietti, S., A.M. Fausto, E. Randelli, O. Carnevali and A.R. Taddei *et al.*, 2009. Early treatment with *Lactobacillus delbrueckii* strain induces an increase in intestinal T-cells and granulocytes and modulates immune-related genes of larval *Dicentrarchus labrax* (L.). *Fish Shellfish Immunol.*, 26: 368-376.
- Ren, W., S. Chen, J. Yin, J. Duan and T. Li *et al.*, 2014. Dietary arginine supplementation of mice alters the microbial population and activates intestinal innate immunity. *J. Nutr.*, 144: 988-995.
- Ren, W., S. Liu, S. Chen, F. Zhang and N. Li *et al.*, 2013. Dietary L-glutamine supplementation increases *Pasteurella multocida* burden and the expression of its major virulence factors in mice. *Amino Acids*, 45: 947-955.
- Rocha, C.S., A.C. Gomes-Santos, T.G. Moreira, M. de Azevedo and T.D. Luerce *et al.*, 2014. Local and systemic immune mechanisms underlying the anti-colitis effects of the dairy bacterium *Lactobacillus delbrueckii*. *PLoS One*, Vol. 9. 10.1371/journal.pone.0085923.

- Salzman, N.H., M.A. Underwood and C.L. Bevins, 2007. Paneth cells, defensins and the commensal microbiota: A hypothesis on intimate interplay at the intestinal mucosa. *Semin. Immunol.*, 19: 70-83.
- Santaolalla, R., M. Fukata and M.T. Abreu, 2011. Innate immunity in the small intestine. *Curr. Opin. Gastroenterol.*, 27: 125-131.
- Santos Rocha, C., O. Lakhdari, H.M. Blottiere, S. Blugeon and H. Sokol *et al.*, 2012. Anti-inflammatory properties of dairy lactobacilli. *Inflamm. Bowel Dis.*, 18: 657-666.
- Schlee, M., J. Harder, B. Koten, E.F. Stange, J. Wehkamp and K. Fellermann, 2008. Probiotic lactobacilli and VSL#3 induce enterocyte α -defensin 2. *Clin. Exp. Immunol.*, 151: 528-535.
- Selsted, M.E. and A.J. Ouellette, 2005. Mammalian defensins in the antimicrobial immune response. *Nat. Immunol.*, 6: 551-557.
- Sengul, N., S. Isik, B. Aslim, G. Ucar and A.E. Demirbag, 2011. The effect of exopolysaccharide-producing probiotic strains on gut oxidative damage in experimental colitis. *Digestive Dis. Sci.*, 56: 707-714.
- Shanahan, M.T., A. Vidrich, Y. Shirafuji, C.L. Dubois and A. Henschen-Edman, 2010. Elevated expression of paneth cell CRS4C in ileitis-prone SAMP1/YITFC mice regional distribution, subcellular localization and mechanism of action. *J. Biol. Chem.*, 285: 7493-7504.
- Strus, M., A. Janczyk, A. Gonet-Surowka, M. Brzywczy-Wloch, G. Stochel, P. Kochan and P.B. Heczko, 2009. Effect of hydrogen peroxide of bacterial origin on apoptosis and necrosis of gut mucosa epithelial cells as a possible pathomechanism of inflammatory bowel disease and cancer. *J. Physiol. Pharmacol.*, 60: 55-60.
- Thomas, C.M. and J. Versalovic, 2010. Probiotics-host communication: Modulation of signaling pathways in the intestine. *Gut. Microbes.*, 1: 148-163.
- Tohno, M., T. Shimosato, H. Aso and H. Kitazawa, 2011. Immunobiotic *Lactobacillus* strains augment NLRP3 expression in newborn and adult porcine gut-associated lymphoid tissues. *Vet. Immunol. Immunopathol.*, 144: 410-416.
- Underwood, M.A., A. Kananurak, C.F. Coursodon, C.K. Adkins-Reick and H. Chu *et al.*, 2012. *Bifidobacterium bifidum* in a rat model of necrotizing enterocolitis: Antimicrobial peptide and protein responses. *Pediatr. Res.*, 71: 546-551.
- Valdovinos-Diaz, M.A., 2013. Intestinal microbiota in digestive disorders. Probiotics, prebiotics and symbiotics. *Rev. Gastroenterol. Mex.*, 78: 25-27.
- Vivier, E. and B. Malissen, 2005. Innate and adaptive immunity: Specificities and signaling hierarchies revisited. *Nat. Immunol.*, 6: 17-21.
- Wang, H., J. Brown, S. Gao, S. Liang and R. Jotwani *et al.*, 2013. The role of JAK-3 in regulating TLR-mediated inflammatory cytokine production in innate immune cells. *J. Immunol.*, 191: 1164-1174.
- Wang, J.Q., F.G. Yin, C. Zhu, H. Yu, S.J. Niven, C.F.M. De Lange and J. Gong, 2012. Evaluation of probiotic bacteria for their effects on the growth performance and intestinal microbiota of newly-weaned pigs fed fermented high-moisture maize. *Livest. Sci.*, 145: 79-86.
- Wehkamp, J., J. Harder, K. Wehkamp, B.W.V. Meissner and M. Schlee *et al.*, 2004. NF- κ B- and AP-1-mediated induction of human β defensin-2 in intestinal epithelial cells by *Escherichia coli* Nissle 1917: A novel effect of a probiotic bacterium. *Infect. Immunity*, 72: 5750-5758.
- Yang, Y.J. and B.S. Sheu, 2012. Probiotics-containing yogurts suppress *Helicobacter pylori* load and modify immune response and intestinal microbiota in the *Helicobacter pylori*-infected children. *Helicobacter*, 17: 297-304.
- Yu, Q., L. Zhu, Z. Wang, P. Li and Q. Yang, 2012. *Lactobacillus delbrueckii* ssp. lactis R4 prevents *Salmonella typhimurium* SL1344-induced damage to tight junctions and adherens junctions. *J. Microbiol.*, 50: 613-617.
- Zarepour, M., K. Bhullar, M. Montero, C. Ma and T. Huang *et al.*, 2013. The mucin muc2 limits pathogen burdens and epithelial barrier dysfunction during *Salmonella enterica* serovar typhimurium colitis. *Infect. Immun.*, 81: 3672-3683.
- Zhao, L. and W. Lu, 2014. Defensins in innate immunity. *Curr. Opin. Hematol.*, 21: 37-42.