



Ootaxonomy and Egg Shell Morphology of Phthirapteran Species Infesting *Coturnix Coturnix* (Linnaeus, 1758) (Phthiraptera: Amblycera: Ischnocera)

¹Aftab Ahmad and ²Neelima Gupta

¹Northern Regional Centre, ZSI, Kaulagarh Road, Dehradun Uttarakhand

²Doctor Harisingh Gour Sagar University (a Central University), Sagar, M. P. India

Key words: *Coturnix coturnix*, egg structure, ischnocera, lice, phthiraptera

Corresponding Author:

Neelima Gupta

Doctor Harisingh Gour Sagar University (A Central University), Sagar Madhya Pradesh, India

Page No.: 189-193

Volume: 20, Issue10, 2021

ISSN: 1680-5593

Journal of Animal and Veterinary Advances

Copy Right: Medwell Publications

Abstract: The morphological features of eggs of three phthirapteran species (*Menacanthus abdominalis* Piaget, 1880, *Cuclotogaster cinereus* Nitzsch, 1866 and *Goniodes astrocephalus* Burn, 1838) of common quail lice, *Coturnix coturnix*, Linnaeus, 1758 were studied through SEM. Differences in shape, size, sculpturing of eggs of each species exhibits difference in microtopography (mention key differences).

INTRODUCTION

Phthirapteran ecto parasites exhibit considerable variation with respect to the egg laying sites, oviposition patterns and egg shell morphology. The eggs of phthirapteran species permanently attach to the feathers/hairs of the hosts and exhibit remarkable and quite polymorphic chorionic structure. Markings present on the eggs of phthirapteran species can act as a useful indicator for the identification of species.

A survey of literature reveals that specific studies on the egg morphology have rarely been made. Certain workers^[1-12] have provided information on the egg shell morphology of phthirapteran species.^[13-14] pointed out the role of egg morphology as a guide to louse taxonomy and further advocated the use of SEM for this purpose.^[20-21] Pointed out the role of egg shell morphology as a guide to louse taxonomy and further suggested the use of SEM of eggs for this purpose.

During the present studies, an attempt has been made to furnish information about the egg shell morphology of three phthirapteran species viz. Infesting common quail, with the help of Scanning Electron Microscopy.

MATERIALS AND METHODS

The nits of each species were obtained from the infested host. The intact eggs were dehydrated in different grades of alcohol. For SEM studies, eggs were fixed in 2.5% glutaraldehyde and passed through 0.2 M phosphate buffer, dehydrated, mounted on double sided glued tape on aluminum stubs, coated with gold palladium alloy in Neo Coater 100-240V and examined under SEM (Neo JCM-6000) at Centre of Excellence laboratory of Mahatma Jyotiba Phule Rohilkhand University, Bareilly, Uttar Pradesh. The samples were then observed under SEM at varying magnifications and selected areas were photographed. Some specimens were treated with Osmium tetra oxide (2%) for achieving better results.

RESULTS AND DISCUSSIONS

***Cuclotogaster cinereus* Nitzsch, 1866:** The egg of *C. cinereus* is miniature rice grain like elongated structure (0.45-0.47 mm. in length, and 0.10-0.12mm. in width)




Fig. 1a-c: (a) Entire egg shell of *Cuclotogaster cinereus*, (b) Enlarged view of opercular end of the egg of *C. cinereus* showing the micropyles and (c) Enlarged view of the posterior end of the egg of *C. cinereus* showing the stigma

fig.1a. Anterior end of egg is capped with almost dome shaped operculum (0.05-0.08mm. in diameter) fig.2. 12-14 micropyles (0.01-0.02mm. in diameter) are arranged near the opercular rim of the egg. Each micropyle appears as a typical button shaped structure. The rear end of the egg shell bears a prominent bee hive like stigma/hydropyle fig.3a. The entire egg chorion of the egg is smooth and does not show any marking or sculpturing/ornamentations.

Goniodus astrocephalus Burn, 1838: The egg of *G.astrocephalus* is ovoid in shape (0.45-0.48 mm. in length, and 0.15-0.18mm. in width) fig.1. The egg chorion remains smooth (devoid any kind of sculpturing/markings and apophyses). The opercular disc is nearly conical in shape (0.04-0.08 mm. in diameter) and shows feebly defined ridges fig.2a. The opercular disc also bears 13-15 button shaped micropyles which are irregular along the ring (0.01-0.02mm. in diameter). A thick rod like polar thread structure arises from anterior end of the operculum which curves downward towards the operculum thus making a hook like structure. The tip of the polar thread is an oblique disc like structure (0.012 mm. in diameter).

The posterior end of the egg exhibits presence of a bee hive like structure called stigma fig.3. The stigma appears to be composed of a hollow chamber.

Menacanthus abdominalis Piaget, 1880: The egg of *M. abdominalis* is ovoid in shape (0.72-0.86mm. in length and 0.23-0.27mm. in width) fig.1. The egg chorion is smooth devoid of sculptures/ornamentations. The operculum is hat shaped structure and lacks polar thread structure (0.11-0.15 mm. in diameter) fig.2. The opercular disc bears faint hexagonal marks. The egg mouth bears 38-40 small filaments like apophyses which are arranged in two rows. Apophyses belonging to the outer row are divided while the inner ones are undivided.

The entire basal portion of the egg remains heavily covered by cementing materials used by lice to glue the egg on the feathers fig.3. Hence, stigma/hydropyle remain concealed in the cementing material. Adult lice in many cases lack significant intergeneric morphological differences and are thus difficult to classify¹⁴. At the same time, the eggs of different species exhibited fascinating differences when examined through SEM. The phthirapteran eggs exhibit certain distinctive




Fig. 2a-c: (a) Entire egg shell of *Goniodes astrocephalus*, (b) Enlarged anterior end of the egg of *G. astrocephalus* showing the micropyles and (c) Enlarged view of the posterior end of the egg of *G. astrocephalus* showing stigma




Fig. 3a-c: (a) Entire egg shell of *Menacanthus abdominalis*, (b) Enlarged anterior end of the egg of *M. abdominalis* showing the micropyles and (c) Enlarged view of the posterior end of the egg of *M. abdominalis* showing the stigma

characteristics on or within chorionic shell. A scrutiny of literature also indicates a number of fascinating adaptive differences in the form of sculpturing/ ornamentations on the chorion. The occurrence of polar thread, filament like process (apophyses) on the egg shell, markings/ ornamentations on the egg shells, opercular disc sculpturing/ ornamentations on the opercular disc and the arrangements of the micropyles on the opercular discs are the main points relating to diversity of the eggs¹⁴. Has categorically pointed out that egg morphology can be used as a guide to louse taxonomy and further advocated the use of SEM of eggs for identification of eggs to genera and where possible to species level. Furthermore, many external features of the eggs are difficult to resolve by the light microscopic studies. However, SEM proved to be an ideal instrument for this purpose revealing the minutest details.

Examined the egg shell morphology of three species of *Lipeurus* (*L. caponis* with granular protuberances; *L. heterographus* having hexagonal pattern; *L. tropicalis* having shell pitted with faint hexagonal markings). Observed that the egg shell of selected species of genus *Menacanthus* differed in location, number and nature of apophyses present on the egg shells. The egg shells of selected species of *Brueelia* differed in the number of micropyles, presence of polar thread.

Review of literature reveals that in case of amblyceran species the differences in the egg morphology of the phthiraptera species appear to be more distinct i.e. poultry shaft louse. Likewise, the differences between chorionic sculpturing of another amblyceran species also have been noted.

However, the differences in the structure of the ischnoceran species are less marked. Two species of genus *Gonicocotes* (*G. gallinane* and *G. jirufti*); three species of genus *Brueelia* (*B. cyclothorax*, *B. amandavae*, *B. saliemii*) and three species of the Genus *Lipeurus* (*L. tropicalis*, *L. caponis* and *L. heterographus*) and one species of the genus *Rallicola* (*R. unguiculatus*) have been studied from this point of view^{3,6}.

ACKNOWLEDGMENTS

I wish to express my sincere gratitude to Dr. Dhriti Banerjee Scientist 'G' & Director, Zoological Survey of India, Kolkata for her encouragement and for providing the facilities for the execution of this work. We also express our sincere thanks to Dr. Gaurav Sharma, Scientist E & Officer-In-Charge, Northern Regional Centre, Zoological Survey of India Kaulagarh Road, Dehradun Uttarakhand for his useful support and encouragement. I also express my deep heartfelt indebtedness to Dr. Anil Mohapatra Scientist E & Officer-In-Charge, Estuarine Biology Regional Centre, Zoological Survey of India, Gopalpur for his valuable suggestions and guidance.

CONCLUSION

The present report furnishes further information on the nature of egg shell of two ischnoceran species, and one amblyceran species *Menacanthus abdominalis* Piaget, 1880 infesting common quail, *Coturnix coturnix* Linnaeus, 1758) with the help of Scanning Electron Microscopy.

The peculiar hexagonal marking/sculpturing on the operculum and filament like apophyses occur along the egg mouth on the eggs of *M. abdominalis* that can be easily differentiated from *C. cinereus* and *G. astrocephalus*. Moreover, a hook like polar thread arises from the opercular disc of the eggs of *G. astrocephalus* and are not found on the eggs of *C. cinereus* on *M. abdominalis*. The arrangement of micropyles on the operculum can also be easily differentiated in the species i.e. micropyles arrangement near the opercular rim in a row present in the egg of *C. cinereus* while irregular distribution of the micropyles on the operculum of the eggs of *G. astrocephalus*. The micropyles are absent on the operculum of *M. abdominalis*.

REFERENCES

1. Agarwal, G.P., A. Ahmad, G. Arya, N. Bansal and A.K. Saxena, 2011. The egg of *laemobothrion maximum* (Amblycera: Phthiraptera). (Amblycera: Phthiraptera) 1: 37-40.
2. Ahmad, A., V. Khan, S. Badola, G. Arya, N. Bansal and A.K. Saxena, 2010. Population characteristics and the nature of egg shells of two Phthirapteran species parasitizing Indian cattle egrets. *J. Insect Sci.*, 10: PP; 1-7.
3. Aftab, A. 2017. The eggshell morphology of *Rallicola unguiculatus* Piaget, 1880 (Ischnocera: Phthiraptera), *J. Parasit. Dis.*, 41: 562-564.
4. Ahmad, A. 2018. Egg laying pattern and the egg morphology of an ischnoceran louse, *Gonicocotes jirufti* (Ansari, 1947) parasitizing black partridges, *Francolinus francolinus* (Phthiraptera: Ischnocera), *Journal of Appl. Nat. Sci.*, 10: 838-840.
5. Castro, D.C., Cicchino, A. Conrado, D. Villalobos and Cristina, 1991. a comparative study of the external chorionic architecture of the eggs of some neotropical species of the genus *hoplopleura enderlein*, 1904 (phthiraptera, anoplura). *Rev. Bras. Entomol.*, 35: pp; 663-669.
6. Gupta, N., V. Khan, S. Kumar, S. Saxena, A. Rashmi and A.K. Saxena, 2009. Eggshell morphology of selected Indian bird lice (Phthiraptera: Amblycera and Ischnocera). *Entomol. News*, 120: 327-336.

7. Kumar, A., A. Kumar, S. Kumar, S.K. Singh and A.K. Saxena, 2003. Egg structure of five phthirapteran species infesting sheep and goats. *J. Parasitic Applied Anim. Biol.*, 12: 25-34.
8. Rajput, S., N. Gupta, V. Khan, A.K. Saxena and V.D. Joshi, 2010. Microtopography of the eggshell of *menacanthus eurysternus* (phthiraptera: amblycera). *J. Appl. Nat. Sci.*, 2: pp; 111-113.
9. Saxena, A.K., S.K. Singh, A. Kumar and Surman 2000. SEM Studies on the microtopography of eggs four pigeon lice (phthiraptera, insecta). *Riv. Parassitol.*, 3: pp; 351-358.
10. Saxena, A. K., G. Arya, N. Bansal, 2012. Egg Laying Site and Oviposition Pattern of Two Phthirapteran Species Parasitizing Red Whiskered Bulbul (*Pycnonotus jocosus*). *Turkiye Parazitol Derg.*, 36: 166-168.
11. Tyagi, S., N.Gupta, V.D. Joshi, A. Rashmi, G. Arya and A.K. Saxena, 2009. The eggshell morphology of *Heterodoxus spiniger*, infesting dog, *Canis familiaris* (Boopidae, Mallophaga). *Appl. Nat. Sci.*, 1: PP; 71-73.
12. Zawadzka, M., W. Jankowska, S.M. Biliński, 1997. Egg shells of mallophagans and anoplurans (Insecta: Phthiraptera): morphogenesis of specialized regions and the relation to F-actin cytoskeleton of follicular cells *Tissue Cell* 29: 665-673.
13. Balter, R.S., 1968. Lice egg morphology as a guide to taxonomy. *Med. Biol. Illus.*, 18: pp; 94-95.
14. Balter, R.S., 1968. The microtopography of avian lice eggs. <https://www.semanticscholar.org/paper/The-microtopography-of-avian-lice-eggs.-Rs/f41e44e3bf23434d0ca94db21b6c81d43af98cdf>