

A Method of the Best Approximation by Fractal Function

¹O. Thae Gun, ¹Kang Yong Suk and ²Rim Myong Gil

¹Faculty of Mathematics, Kim Il Sung University, Pyonyang, North Korea

²Faculty of Mathematics, Kim Hyen Jik Education University,
Democratic People's Republic of Korea, Pyongyang, Korea

Abstract: We present a method constructing a function which is the best approximation for given data and satisfies the given self-similar condition. For this, we construct a space F of local self-similar fractal functions and show its properties. Next we present a computational scheme constructing the best fractal approximation in this space and estimate an error of the constructed fractal approximation. Our best fractal approximation is a fixed point of some fractal interpolation function.

Key words: Fractal interpolation, fractal approximation, iterated function system, fractal function space, construct

INTRODUCTION

Fractal approximation has been applied to model the objects which have fractal characteristics in nature. Fractal functions whose graphs are fractal sets have been widely used in approximation theory, signal processing, interpolation theory, computer graphics and so on. Hence, constructions of fractal functions and fractal approximation have been studied in many papers (Barnsley, 1986, 1988; Barnsley *et al.*, 1989; Bouboulis and Dalla, 2007a, b; Bouboulis and Mavroforakis, 2011; Bouboulis *et al.*, 2006; Chand and Kapoor, 2006; Dalla, 2002; Feng *et al.*, 2012; Malysz, 2006; Massopust, 1997; Metzler and Yun, 2010; Navascues and Sebastian, 2004; Lonardi and Sommaruga, 1999; Kang *et al.*, 2014; Yun *et al.*, 2014; Hefei *et al.* 1999).

Constructions of fractal functions by fractal interpolation have been introduced by many researchers. A construction of one variable fractal interpolation functions by the Iterated Function System (IFS) with a data set on \mathbb{R} was studied by Barnsley (1986, 1988), Navascues and Sebastian (2004) where the constructed fractal functions were self-similar ones. The construction was generalized by Barnsley *et al.* (1989), Bouboulis and Dalla (2007a) and Yun *et al.* (2014) which constructed local self-similar fractal functions. Constructions of Bivariate Fractal Interpolation Functions (BFIFs) have been studied by Bouboulis and Mavroforakis (2011), Bouboulis *et al.* (2006), Chand and Kapoor (2006), Dalla (2002), Feng *et al.* (2012), Malysz (2006), Massopust (1997), Navascues and Sebastian (2004) and Yun *et al.* (2014). A construction of BFIFs by fractal interpolation on \mathbb{R} was presented by Bouboulis and Dalla (2007b), Yun *et al.* (2014) and self-affine fractal interpolation functions were constructed by IFS with a data set on a

triangular domain by Metzler and Yun (2010). Constructions of self-similar BFIFs by Dalla (2002), Feng *et al.* (2012) and Malysz (2006) and self-affine BFIFs by Massopust (1997) by IFS with a data set on a rectangular grid were introduced. Bouboulis and Mavroforakis (2011) local self-similar BFIFs were constructed by the Recurrent Iterated Function System (RIFS) on a rectangular grid.

A construction of local self-similar fractal interpolation functions in \mathbb{R}^n was studied by Bouboulis and Dalla (2007b). To construct fractal interpolation we need a data set $\{(x_i, y_i), i = 0, 1, \dots, n\}$ and a set of scale parameters $\{s_i, i = 1, \dots, n\}$. The fractal property of the graph of the interpolation function is determined by those data. Let a division of the interval and scale parameters be given, that is a fractal property of the function be given. If the number of experimental data is more than the number of the interval division, then we can not construct the fractal interpolation for the data using fractal interpolation theory.

So, we assume that a division of the interval and scale parameters be given (that is a fractal property of the function) and study the problem constructing the best fractal approximation for the data set $\{(\bar{x}_i, \bar{z}_i), i = 0, 1, \dots, m\}$ where, $m > n$ (n is the number of the interval division). Lonardi and Sommaruga (1999) and Hefei *et al.* (1999) constructions of the best approximation of functions by the fractal functions were presented, respectively. But the continuity of the approximation was not guaranteed then. The best fractal approximation of a continuous function in L^2 space was introduced by Bouboulis and Mavroforakis (2011) a space of differentiable fractal interpolation functions was constructed and it was proved that the constructed space is the reproducing Kernel Hilbert space.

MATERIALS AND METHODS

A space of local self-similar fractal functions and a space of contractive operators: In this study, we construct a space of local self-similar fractal functions and a space of contraction operators which are isomorphic to each other. Let:

$$\Delta = \{x_i \in \mathbb{R} : i = 0, 1, \dots, n\}, a = x_0 < x_1 < \dots < x_n = b,$$

$$I = [a, b], I_i = [x_{i-1}, x_i]$$

$$\{s_i : |s_i| < 1, i = 1, 2, \dots, n\}$$

be given. Let $1 < q \leq n$ ($q \in \mathbb{N}$), $J_k = [x_{s(k)}, x_{e(k)}]$, $x_{s(k)}, x_{e(k)} \in \{x_0, x_1, \dots, x_n\}$ and $e(k) - s(k) \geq 2$, $k = 1, \dots, q$. I_i is called a region and J_k a domain. We define a mapping $\gamma: \{1, \dots, n\} \rightarrow \{1, \dots, q\}$ which means that we relate every region to a domain. For each $i \in \{1, \dots, n\}$, denote $k = \gamma(i)$. For $i \in \{1, \dots, n\}$ define a mapping $u_{i,k}: J_k \rightarrow I_i$ by:

$$u_{i,k}(x) = a_i x + b_i \tag{1}$$

which satisfies:

$$u_{i,k}(x_{s(k)}) = x_{i-1}, u_{i,k}(x_{e(k)}) = x_i$$

Let $f \in C(I)$ be a continuous function satisfying:

$$f(x) = s_i f(u_{i,k}^{-1}(x)) + p_{i,k}(u_{i,k}^{-1}(x)), x \in I_i \tag{2}$$

where, functions $p_{i,k}: J_k \rightarrow I_i$, $i = 1, \dots, n$ are defined by $p_{i,k}(x) = c_i x + d$ and satisfy the following conditions:

$$s_i f(x_{s(k)}) + p_{i,k}(x_{s(k)}) = f(x_{i-1}) \tag{3}$$

$$s_i f(x_{e(k)}) + p_{i,k}(x_{e(k)}) = f(x_i) \tag{4}$$

Define a space of functions satisfying Eq. 3 and 4 by F . The graph of $f \in F$ has a local self-similarity and we get $f(x) = 0 \in F$ which corresponds to $c_i = 0, d_i = 0, i \in \{1, \dots, n\}$.

Lemma 1: F is a linear subspace of dimension $n+1$ of $C(I)$.

Proof: For $f, \tilde{f} \in F$ and $\lambda \in \mathbb{R}$, we have:

$$f(x) = s_i f(u_{i,k}^{-1}(x)) + p_{i,k}(u_{i,k}^{-1}(x)), \tilde{f}(x) = s_i \tilde{f}(u_{i,k}^{-1}(x)) + \tilde{p}_{i,k}(u_{i,k}^{-1}(x))$$

Hence:

$$(f + \tilde{f})(x) := s_i (f + \tilde{f})(u_{i,k}^{-1}(x)) + (p_{i,k} + \tilde{p}_{i,k})(u_{i,k}^{-1}(x))$$

$$(\lambda f)(x) := s_i (\lambda f)(u_{i,k}^{-1}(x)) + (\lambda p_{i,k})(u_{i,k}^{-1}(x))$$

Thus, $f + \tilde{f} \in F$ and $\lambda f \in F$. Because for $f \in F$, $(f(x_0), f(x_1), \dots, f(x_n)) \in \mathbb{R}^{n+1}$ is uniquely determined, a mapping $\Psi: F \rightarrow \mathbb{R}^{n+1}$ is defined by:

$$\Psi(f) = (f(x_0), f(x_1), \dots, f(x_n)) \tag{5}$$

And for $(y_0, y_1, \dots, y_n) \in \mathbb{R}^{n+1}$, there exists a unique $f \in F$ such that $f(x_i) = y_i, i = 0, \dots, n$. In fact, the existence and uniqueness of f are ensured by the existence and uniqueness of the recurrent fractal interpolation function (Barnsley *et al.*, 1989). This shows that the mapping $\Psi: F \rightarrow \mathbb{R}^{n+1}$ is a bijection. We can easily check that the mapping Ψ is linear. Hence, F and \mathbb{R}^{n+1} are isomorphic. A basis of F is:

$$\Psi^{-1}(e_i), e_i = (0, \dots, 1, 0, \dots, 0) \quad i = 1, \dots, n+1 \tag{6}$$

The space F is a Banach space with the norm $\|\cdot\|_\infty$. For a $y = (y_0, y_1, \dots, y_n) \in \mathbb{R}^{n+1}$, define a function space F_y by:

$$F_y = \{f \in C(I) : f(x_i) = y_i, i = 1, \dots, n\}$$

Then, $(F_y, \|\cdot\|_\infty)$ is a complete space. For $f \in F$ define a function $T_y f: I \rightarrow \mathbb{R}$ by:

$$(T_y f)(x) := s_i f(u_{i,k}^{-1}(x)) + p_{i,k}^{T_y}(u_{i,k}^{-1}(x)) \quad x \in I_i \tag{7}$$

where, $p_{i,k}^{T_y}(x) = c_i^{T_y} x + d_{i,k}^{T_y}$ satisfies the following conditions:

$$(T_y f)(x_{i-1}) = y_{i-1}, (T_y f)(x_i) = y_i$$

i.e.,

$$s_i y_{s(k)} + p_{i,k}^{T_y}(x_{s(k)}) = y_{i-1}, s_i y_{e(k)} + p_{i,k}^{T_y}(x_{e(k)}) = y_i$$

By these $c_{i,k}^{T_y}$ conditions $d_{i,k}^{T_y}$ and are uniquely given by:

$$c_{i,k}^{T_y} = \frac{(y_i - y_{i-1}) - s_i (y_{e(k)} - y_{s(k)})}{x_{e(k)} - x_{s(k)}} \tag{8}$$

$$d_{i,k}^{T_y} = \frac{y_{i-1} x_{e(k)} - y_i x_{s(k)} + s_i (x_{s(k)} y_{e(k)} - y_{s(k)} x_{e(k)})}{x_{e(k)} - x_{s(k)}} \tag{9}$$

Therefore, for a $y = (y_0, y_1, \dots, y_n) \in \mathbb{R}^{n+1}$, we get a unique operator T_y . And because $T_y f$ is continuous by Eq. 7, $T_y f \in F_y$. Thus, the operator $T_y: F_y \rightarrow F_y$ is defined by Eq. 7. It is easy to verify that the operator T_y is a contraction with respect to $\|\cdot\|_\infty$. According to the fixed-point theorem in a complete space, there exists a unique $f_{T_y} \in F_y$ such that:

$$(f_{T_y}(x_0), f_{T_y}(x_1), \dots, f_{T_y}(x_n)) = (y_0, y_1, \dots, y_n)$$

Let τ be the set of such operators. Define a mapping $\Phi: R^{n+1} \rightarrow \tau$ by $\Phi(y) = T_y \in \tau$. Then, the mapping Φ is a bijection.

Lemma 2: The τ is a linear space of dimension $n+1$.

Proof: For $T_{y_1}, T_{y_2} \in \tau$ and $\lambda \in R$, define T_{y_1}, T_{y_2} on $F_{y_1+y_2}$ by:

$$f \in F_{y_1+y_2} (T_{y_1} + T_{y_2})(f)(x) := s_1 \cdot f(u_1^{-1}(x)) + (p_1^{T_{y_1}} + p_1^{T_{y_2}})(u_1^{-1}(x)), x \in I_1$$

and λT_{y_1} on $E_{\lambda y_1}$ by:

$$\begin{aligned} \tilde{f} \in F_{\lambda y_1} (\lambda T_{y_1})(\tilde{f})(x) &:= s_1 \cdot \tilde{f}(u_1^{-1}(x)) + \\ \lambda p_1^{T_{y_1}}(u_1^{-1}(x)) \quad x \in I_1 \quad i = 1, \dots, n \end{aligned}$$

Note that we omit a subscript k after this because the domain and region are all fixed. It is clear that $(T_{y_1} + T_{y_2})f \in F_{y_1+y_2}$ and $(\lambda T_{y_1})\tilde{f} \in F_{\lambda y_1}$. Therefore, $T_{y_1} + T_{y_2} \in F$ and $\lambda T_{y_1} \in \tau$, i.e., the linear operations are defined in the set T . It is easy to prove that the set T is a linear space with respect to the linear operations. The mapping $\Phi: R^{n+1} \rightarrow T$ is linear. In fact, because for $y_1 = (y_{1,0}, y_{1,1}, \dots, y_{1,n})$ and $y_2 = (y_{2,0}, y_{2,1}, \dots, y_{2,n}) \in R^{n+1}$, $\lambda \in R$ by Eq. 8 and 9:

$$\begin{aligned} p_1^{T_{y_1+y_2}}(x) &= c_1^{T_{y_1+y_2}}x + d_1^{T_{y_1+y_2}} = (c_1^{T_{y_1}} + c_1^{T_{y_2}})x + (d_1^{T_{y_1}} + d_1^{T_{y_2}}) \\ &= c_1^{T_{y_1}}x + d_1^{T_{y_1}} + c_1^{T_{y_2}}x + d_1^{T_{y_2}} = p_1^{T_{y_1}}(x) + p_1^{T_{y_2}}(x) \\ p_1^{T_{\lambda y_1}}(x) &= c_1^{T_{\lambda y_1}}x + d_1^{T_{\lambda y_1}} = \lambda(c_1^{T_{y_1}}x + d_1^{T_{y_1}}) = \lambda p_1^{T_{y_1}}(x) \end{aligned}$$

and we get:

$$\begin{aligned} (\Phi(y_1 + y_2))(f)(x) &= (\Phi(y_1) + \Phi(y_2))(f)(x), \\ (\Phi(\lambda y_1))(f)(x) &= \lambda(\Phi(y_1))(f)(x) \end{aligned}$$

Hence, τ and R^{n+1} are isomorphic which means that the dimension of τ is $n+1$. By the isomorphic relation $(0, 0, \dots, 0) \in R^{n+1}$ corresponded to the operator T defined by:

$$(Tf)(x) = s_1 \cdot f(u_1^{-1}(x)), \quad x \in I_1$$

whose fixed point is $f_T(x) \equiv 0$.

Theorem 1: Let F and T be the linear spaces constructed above. Then, they are isomorphic.

Proof: This follows from Lemmas 1 and 2. Denote the isomorphism of F - T by $\tilde{\Psi}$. Note that for $f \in F$, the fixed point of T with $\tilde{\Psi}(f) = T$ is f .

RESULTS AND DISCUSSION

Construction of LSFA of a data set: In this study, we prove that there exists the least squares fractal approximation f in F of a data set and present an algorithm for finding f by calculating approximately the contraction operator T in τ corresponding to f . Let P be a data set given by:

$$P = \{(\bar{x}_i, z_i) : i = 0, 1, \dots, m\} \tag{10}$$

$$(\bar{x}_0 < \bar{x}_1 < \dots < \bar{x}_m, \bar{x}_0 = x_0, \bar{x}_m = x_n)$$

where, $m > n$. An f^* is called the Least Squares Fractal Approximation (LSFA) if f^* is a solution of the following question:

$$\min_{f \in F} \sum_{i=0}^m (f(\bar{x}_i) - \bar{z}_i)^2 \tag{11}$$

First, we consider the existence and uniqueness of LSFA.

Theorem 2: If $\{x_0, x_1, \dots, x_n\} \subset \{\bar{x}_0, \dots, \bar{x}_m\}$ then there exist a unique solution $f^* \in F$ of Eq. 11 and a unique $T_p \in \tau$ whose fixed point is f^* .

Proof: Define an operator $B_m: F \rightarrow R^{n+1}$ by:

$$f \in F, B_m f = (f(\bar{x}_0), f(\bar{x}_1), \dots, f(\bar{x}_m))$$

and denote $B_m F$ by D . Then B_m is a linear operator and D is a linear subspace of R^{m+1} . Equation 11 is represented by:

$$\min_{f \in F} \|B_m f - \hat{z}\|_E^2$$

where, $\hat{z} = \{\bar{z}_0, \bar{z}_1, \dots, \bar{z}_m\} \in R^{m+1}$ and $\|\cdot\|_E$ is the Euclidean norm. Therefore Eq. 11 is equivalent to the Eq. 13:

$$\min_{z \in D} \|z - \hat{z}\|_E^2 \tag{13}$$

Because $(R^{m+1}, \|\cdot\|_E)$ is a Hilbert space and D is a subspace of R^{m+1} , there exists a unique solution z^* of Eq. 13. If $B_m f = 0$, then from the hypothesis of the theorem:

$$(f(x_0), f(x_1), \dots, f(x_n)) = (0, 0, \dots, 0)$$

and $f(x) = 0$ by the construction of F . Therefore, B_m is an injection and there exists a unique $f^* \in F$ such that $B_m f^* = z^*$, i.e., there exists a unique $\tau^* = \Psi^{-1}(f^*) \in \tau$. From Theorem 2, Eq. 11 is equivalent to the following equation:

$$\min_{\tau \in \tau} \|B_m f_\tau - \hat{z}\|_E^2$$

Now, we consider a construction of the LSFA. Let Ψ be the linear mapping defined by Eq. 5 and denote:

$$v_i = \Psi^{-1}(e_i), e_i = (0, \dots, 1, 0, \dots, 0), i = 1, \dots, n+1$$

Then, $\{v_i\}_{i=1}^{n+1}$ is a basis of F and there exist unique $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ such that:

$$f^* = \sum_{k=0}^n \lambda_k v_k$$

For $f, g \in F$, define $\langle f, g \rangle \in \mathbb{R}$ by $\langle f, g \rangle = \sum_{k=0}^m f(\bar{x}_k) \cdot g(\bar{x}_k)$. We get a normal equation:

$$A\alpha = b$$

$$A = (a_{ij}), a_{ij} = \langle v_i, v_j \rangle, b = (b_i), b_i = \sum_{k=0}^m z_k v_i(\bar{x}_k) \quad (14)$$

to find $f^* \in F$. Since, $v_i, i = 1, \dots, n$ are fractal functions in Eq. 14, it needs enormous operations. Therefore, we consider an algorithm for calculating the approximation of contraction operator T_F . We calculate approximately f^* as the fixed point of T_F . Now, for $p_0 = (\bar{x}_0, \bar{x}_1, \dots, \bar{x}_m)$, let us denote $X_m = p_0 \times \mathbb{R}^{m+1}$. Define an operator T_m on X_m by:

$$z = (z_0, z_1, \dots, z_m) \in \mathbb{R}^{m+1}, T_m(p_0, z) = (p_0, \bar{z}),$$

$$\bar{z} = (\bar{z}_0, \bar{z}_1, \dots, \bar{z}_m)$$

where, $\bar{z}_i, i = 1, 2, \dots, m$ are defined as follows: for \bar{x}_i , there exist $l \in \{1, \dots, n\}$ and $k \in \{0, 1, \dots, m-1\}$ such that $\bar{x}_i \in I_l$ and $\bar{x}_k \leq u_1^{-1}(\bar{x}_i) \leq \bar{x}_{k+1}$. Then:

$$\bar{z}_i = s_l(z_k + z_{k+1}) / 2 + c_l u_1^{-1}(\bar{x}_i) + d_l$$

The operator T_m is given by s_i, c_i and $d_i, i = 1, \dots, n$ where $\|s_i\| < 1, i = 1, \dots, n$ and $c_i, d_i, i = 1, 2, \dots, n$ are calculated by Eq. 8 and 9 and represented by y_0, y_1, \dots, y_n . Let us denote $(T_m(p_0, z))_2 = \bar{z}$. We find a T_m^* such that:

$$\|(T_m(p_0, \hat{z}))_2 - \hat{z}\|_E = \|\bar{z} - \hat{z}\|_E \rightarrow \min \quad (15)$$

This problem is a minimization problem of a multi-variable function with unknown y_0, y_1, \dots, y_n . We find




Fig. 1: Relation between T and T_m

y_0, y_1, \dots, y_n from this problem. Next we find the RB operator T_m^* using the method constructing the fractal interpolation and its fixed point, that is the fractal interpolation with $\{(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)\}$ and scale parameters s_1, s_2, \dots, s_n is our best fractal approximation.

Estimation for errors of the approximation: In this study, we consider a relation between T_m and T and estimate an error between the approximation solution f_{τ^*} and given data. For $(p_0, y), (p_0, g) \in X_m$ and $\lambda \in \mathbb{R}$, define $(p_0, y) + (p_0, g), \lambda(p_0, y)$ and $\|(p_0, y)\|$ as follows:

$$(p_0, y) + (p_0, g) := (p_0, y + g), \lambda(p_0, y) := (p_0, \lambda y)$$

$$\|(p_0, y)\| := \max_{0 \leq i \leq m} |y_i|$$

Lemma 3: T_m is a contraction operator on X_m .

Proof: For $(p_0, y), (p_0, g) \in X_m$, we get:

$$|\bar{y}_i - \bar{g}_i| = \frac{s_i}{2} |y_k + y_{k+1} - g_k - g_{k+1}|$$

$$\|T_m(p_0, y) - T_m(p_0, g)\| = \|(p_0, \bar{y}) - (p_0, \bar{g})\| = \|(p_0, \bar{y} - \bar{g})\| = \max_i |\bar{y}_i - \bar{g}_i|$$

$$\leq \frac{s_1}{2} (|y_k - g_k| + |y_{k+1} - g_{k+1}|)$$

$$\leq s_1 \max_i |y_i - g_i| = s_1 \|(p_0, y - g)\|$$

where, $c = \max\{|s_i|, 1, \dots, |s_n|\} < 1$. Therefore, T_m is a contraction operator with contraction constant c . Because X_m is equivalent to \mathbb{R}^{m+1} , we identify X_m with \mathbb{R}^{m+1} and get a diagram that shows the relation between T and T_m (Fig. 1).

Lemma 4: Let $\bar{x}_i - \bar{x}_{i-1} = \bar{x}_m - \bar{x}_0 / m$, for $i = 1, 2, \dots, m$. Let T and T_m be defined by the same $s_i, c_i, d_i, i = 1, 2, \dots, n$. Then for $g \in F$, we have:

$$\|T_m B_m g - B_m T g\|_{X_m} \rightarrow 0 \quad (16)$$

Proof: By the definitions of B_m and T_m , we get:

$$B_m g = (g(\bar{x}_0), g(\bar{x}_1), \dots, g(\bar{x}_m)), T_m B_m g = (\bar{g}_0, \bar{g}_1, \dots, \bar{g}_m)$$

$$\begin{aligned} \bar{g}_i &= s_i(g_k + g_{k+1})/2 + p_i(u_1^{-1}(\bar{x}_i)) \\ \bar{x}_i \in I_1, \bar{x}_k \leq u_1^{-1}(\bar{x}_i) &\leq \bar{x}_{k+1} \end{aligned}$$

And:

$$\begin{aligned} Tg(x) &= s_i g(u_1^{-1}(x)) + p_i(u_1^{-1}(x)), x \in I_1 \\ B_m Tg &= (Tg(\bar{x}_0), Tg(\bar{x}_1), \dots, Tg(\bar{x}_m)) \end{aligned}$$

Therefore, we have:

$$\begin{aligned} |(T_m B_m g)_i - (B_m Tg)_i| &= |s_i(g_k + g_{k+1})/2 + \\ p_i(u_1^{-1}(\bar{x}_i)) - s_i g(u_1^{-1}(\bar{x}_i)) - p_i(u_1^{-1}(\bar{x}_i))| &= \\ |s_i((g_k + g_{k+1})/2 - g(u_1^{-1}(\bar{x}_i)))| &\rightarrow 0 \quad (m \rightarrow \infty) \end{aligned}$$

which gives in Eq. 16. If T and T_m are defined by the same s_i, c_i and d_i, i = 1, 2, ..., n, then since contraction constants of T and T_m are given by s_i, i = 1, 2, ... n, the elements of T and T_m have the same contraction constant c.

Theorem 3: Let T and T_m be defined by the same s_i, c_i and d_i, i = 1, 2, ... n. If for $\bar{x}_0, \bar{x}_1, \dots, \bar{x}_m, u_1^{-1}(\bar{x}_i) \in \{\bar{x}_0, \bar{x}_1, \dots, \bar{x}_m\}, \bar{x}_i \in I_1$, then we get $B_m f_T = T_m B_m f_T$ and:

$$\|B_m f_T - \hat{z}\| \leq \frac{1}{1-c} \|T_m \hat{z} - \hat{z}\| \quad (17)$$

Proof: Since, f_T is the fixed point of T, we have:

$$f_T(x) = T f_T(x) = s_i \cdot f_T(u_1^{-1}(x)) + p_i(u_1^{-1}(x)), x \in I_1$$

and by the definitions of B_m and T_m, we get:

$$(B_m f_T(x))_i = s_i \cdot f_T(u_1^{-1}(\bar{x}_i)) + p_i(u_1^{-1}(\bar{x}_i))$$

Since, $B_m f_T = (f_T(\bar{x}_0), \dots, f_T(\bar{x}_m))$, we have:

$$T_m B_m f_T = T_m (f_T(\bar{x}_0), \dots, f_T(\bar{x}_m))$$

And:

$$(T_m B_m f_T)_i = s_i \cdot f_T(u_1^{-1}(\bar{x}_i)) + p_i(u_1^{-1}(\bar{x}_i))$$

Hence, we have $B_m f_T = T_m B_m f_T$ and:

$$\begin{aligned} \|B_m f_T - \hat{z}\| &\leq \|B_m f_T - T_m \hat{z}\| + \|T_m \hat{z} - \hat{z}\| \\ &= \|T_m B_m f_T - T_m \hat{z}\| + \|T_m \hat{z} - \hat{z}\| \\ &\leq c \|B_m f_T - \hat{z}\| + \|T_m \hat{z} - \hat{z}\| \end{aligned}$$

Thus:

$$\|B_m f_T - \hat{z}\| \leq \frac{1}{1-c} \|T_m \hat{z} - \hat{z}\|$$

Lemma 5: Barnsley (1986), let X be a Banach space and T a contraction operator on X with the contraction constant c. Let f_T be the fixed point of T. If for $f \in X, \|f - T f\| < \epsilon$ then:

$$\|f_T - f\| < \epsilon / (1 - c)$$

Denote $\epsilon_1 = \|T_m^* \hat{z} - \hat{z}\|$ and $\epsilon_2 = \|T_m^* B_m f_T - B_m f_T\| = \|T_m^* B_m f_T - B_m T^* f_T\|$.

Theorem 4: Let f_{T*} be the fixed point of T* defined by the solution (y^{*}₀, ..., y^{*}_n) of Eq. 15. Then, we have

$$\|B_m f_{T^*} - \hat{z}\| \leq (\epsilon_1 + \epsilon_2) / (1 - c) \quad (19)$$

where, c is the contraction constant of the contraction operator T^{*}_m. Especially, $\epsilon_2 = 0$ under the conditions of Theorem 3.

Proof: We can easily see that:

$$\|B_m f_{T^*} - \hat{z}\| \leq \|B_m f_{T^*} - f_{T_m^*}\| + \|f_{T_m^*} - \hat{z}\| \quad (20)$$

From Lemma 4, we have $\|T_m^* B_m f_{T^*} - B_m f_{T^*}\| \rightarrow 0$. Let us denote $p = T_m^* B_m f_{T^*} - B_m f_{T^*}, p = (p_1, \dots, p_m)$. For $\bar{x}_i \in I_1$, there exists a $k \in \{0, 1, \dots, m\}$ such that $\bar{x}_k \leq u_1^{-1}(\bar{x}_i) \leq \bar{x}_{k+1}$. Then, we get:

$$\begin{aligned} p_i &= s_i(f_{T^*}(\bar{x}_k) + f_{T^*}(\bar{x}_{k+1})) / 2 + p_i(u_1^{-1}(\bar{x}_i)) - s_i f_{T^*}(u_1^{-1}(\bar{x}_i)) - \\ p_i(u_1^{-1}(\bar{x}_i)) &= s_i((f_{T^*}(\bar{x}_k) + f_{T^*}(\bar{x}_{k+1})) / 2 - f_{T^*}(u_1^{-1}(\bar{x}_i))) \end{aligned}$$

From Lemma 5, we have:

$$\|B_m f_{T^*} - f_{T_m^*}\| \leq \epsilon_2 / (1 - c), \|f_{T_m^*} - \hat{z}\| \leq \epsilon_1 / (1 - c)$$

where, $c = \max \{|s_i|, i = 1, \dots, n\}$. By Eq. 20 and 21, we get Eq. 19.

Examples of calculation

Example 1: Let P be a data set given by: $P = \{(x_i, z_i) \in \mathbb{R}^2 : i = 0, 1, \dots, 10\} = \{(0, 3.6), (0.1, 5.1), (0.2, 5.6), (0.3, 6.3), (0.4, 6.0), (0.5, 5.4), (0.6, 5.6), (0.7, 5.0), (0.8, 4.2), (0.9, 3.2), (1, 1.7)\}$. Let $\{x_0, x_1, x_2, x_3, x_4\} = \{0, 0.2, 0.5, 0.7, 1\}$, $\Delta = \{(0, y_0), (0.2, y_1), (0.5, y_2), (0.7, y_3), (1, y_4)\}$, $S = \{s_1, s_2, s_3, s_4\} = \{1/3, 2/5, 1/7\}$. Then, $I_1 = [0, 0.2], I_2 = [0.2, 0.5], I_3 = [0.5, 0.7]$ and $I_4 = [0.7, 1]$. Let $J_1 = [0, 1], J_2 = [0, 1], J_3 = [0, 1], J_4 = [0, 1]$. By Eq. 1 and 2, we have $u_1(x) = 0.2x, u_2(x) = 0.3x + 0.2, u_3(x) = 0.2x + 0.5, u_4 = 0.3x + 0.7$ and by Eq. 8 and 9 $c_p, d_p = I = 1, \dots, 4$ given by $c = \{0.666667y_0 + y_1 - 0.333333y_4, 0.4y_0 - y_1 + y_2 - 0.4y_4, 0.166667y_0 - y_2 + y_3 - 0.166667y_4, 0.142857y_0 - y_3 + 0.857143y_4\}$, $d = \{0.666667y_0 - 0.4y_0 + y_1, -0.166667y_0 + y_2, -0.142857y_0 + y_3\}$. Then, we have $(y_0, y_1, y_2, y_3, y_4) = (3.16738, 4.97274, 5.05272, 4.84987, 1.66452)$ from the Eq. 15. Hence, we get $c = (2.30631, 0.681127, 0.0476312, -2.97066), d = (2.11159, 3.70579, 4.52482, 4.39739)$. The attractor of IFS $\{\mathbb{R}^2, w_1, w_2, w_3, w_4\}$:




Fig. 2: LSFA of a data set. The points are one of the data set




Fig. 4: A data set of the coastline




Fig. 3: The coastline




Fig. 5: The fractal approximation of the coastline

$$w_i \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} u_i(x) \\ s_i y + c_i x + d_i \end{pmatrix}, i = 1, 2, 3, 4$$

is the graph of the found least squares fractal approximation (Fig. 2).

Example 2: Calculate LSFA of a coastline with a data set taken from the coastline in Fig. 3. We take the following data set: $P = \{(2.0, 5.82), (2.1, 5.86), (2.2, 5.9), (2.3, 5.92), (2.4, 5.92), (2.5, 5.92), (2.6, 6.0), (2.7, 6.22), (2.8, 6.29), (2.9, 6.31), (3.0, 6.29), (3.1, 6.27), (3.2, 6.39), (3.3, 6.63), (3.4, 7.15), (3.5, 7.29), (3.6, 7.47), (3.7, 7.49), (3.8, 7.55), (3.9, 7.59), (4.0, 7.61), (4.1, 7.54), (4.2, 7.42), (4.3, 7.26), (4.4, 7.28), (4.5, 7.36), (4.6, 7.36), (4.7, 7.40), (4.8, 7.40), (4.9, 7.36), (5.0, 7.32), (5.1, 7.42), (5.2, 7.38), (5.3, 7.38), (5.4, 7.28), (5.5, 7.01), (5.6, 6.75), (5.7, 6.54), (5.8, 6.23), (5.9, 5.52), (6.0, 5.48), (6.1, 5.44), (6.2, 5.36), (6.3, 5.46), (6.4, 5.34), (6.5, 5.18), (6.6, 5.24), (6.7, 5.28), (6.8, 5.16), (6.9, 5.14), (7.0, 5.12), (7.1, 5.16), (7.2, 5.12), (7.3, 5.10), (7.4, 5.06), (7.5, 5.1), (7.6, 5.48), (7.7, 5.82), (7.8, 5.98), (7.9, 5.98), (8.0, 5.84), (8.1, 5.70), (8.2, 5.62), (8.3, 5.58), (8.4, 5.4), (8.5, 5.28), (8.6, 5.20), (8.7, 5.24), (8.8, 5.2), (8.9, 5.14), (9.0, 5.12), (9.1, 5.20), (9.2, 5.50), (9.3, 5.50), (9.4, 5.54), (9.5, 5.56), (9.6, 5.42), (9.7, 5.40), (9.8, 5.6), (9.9, 5.64), (10.0, 5.68), (10.1, 5.64), (10.2, 5.64), (10.3, 5.60), (10.4, 5.62), (10.5, 5.70),$

$(10.6, 5.74), (10.7, 5.74), (10.8, 5.78), (10.9, 5.86), (11.0, 5.88), (11.1, 5.88), (11.2, 5.70), (11.3, 5.76), (11.4, 5.84), (11.5, 5.78), (11.6, 5.74), (11.7, 5.74), (11.8, 5.68), (11.9, 5.58), (12.0, 5.52)\}$
 Then, we have $I = [2, 12], \Delta = \{x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}\}, I_1 = [2, 3], I_2 = [3, 4], I_3 = [4, 5], I_4 = [5, 6], I_5 = [6, 7], I_6 = [7, 8], I_7 = [8, 9], I_8 = [9, 10], I_9 = [10, 11], I_{10} = [11, 12], \{y_0, y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8, y_9, y_{10}\}, (5.71008, 6.22253, 7.4376, 7.58819, 5.95495, 4.91328, 5.74059, 5.12894, 5.59619, 5.83781, 5.6046), c = (0.0522994, 0.122562, 0.0161142, -0.162269, -0.103113, 0.0837858, -0.0601096, 0.0477794, 0.0252167, -0.0222661), d = (5.03448, 5.4064, 6.83436, 7.34172, 5.59017, 4.1747, 5.2898, 4.46238, 4.97475, 5.31133). The attractor of IFS $\{R^2: w_1, w_2, w_3, w_4\}$:$

$$w_i \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} u_i(x) \\ s_i y + c_i x + d_i \end{pmatrix}, i = 1, 2, \dots, 10$$

is the graph of the found least squares fractal approximation (Fig. 4 and 5).

CONCLUSION

We construct a space of fractal interpolation functions with a given division of the interval and scale

parameters and find a function satisfying some approximation condition for data $\{(\bar{x}_i, \bar{y}_i), i=0, 1, \dots, m\}$ with $m>n$ in this space. We call it a local self-similar fractal approximation. The values of the function at nodes of division $\{y_i, i = 0, 1, \dots, n\}$ are unknown unlike interpolation function.

REFERENCES

- Barnsley, M.F., 1986. Fractal functions and interpolation Constructive Approximation, 2: 303-329.
- Barnsley, M.F., 1988. Fractals Everywhere. Academic Press, New York, USA., ISBN-10: 0-12-079061-0.
- Barnsley, M.F., J.H. Elton and D.P. Hardin, 1989. Recurrent iterated function systems. Constructive Approximation, 5: 3-31.
- Bouboulis, P. and L. Dalla, 2007a. A general construction of fractal interpolation functions on grids of n . Eur. J. Appl. Math., 18: 449-476.
- Bouboulis, P. and L. Dalla, 2007b. Fractal interpolation surfaces derived from fractal interpolation functions. J. Math. Anal. Appl., 336: 919-936.
- Bouboulis, P. and M. Mavroforakis, 2011. Reproducing kernel Hilbert spaces and fractal interpolation. J. Comp. Appl. Math., 235: 3425-3434.
- Bouboulis, P., L. Dalla and V. Drakopoulos, 2006. Construction of recurrent bivariate fractal interpolation surfaces and computation of their box-counting dimension. J. Approximation Theory, 141: 99-117.
- Chand, A.K.B. and G.P. Kapoor, 2006. Generalized cubic spline fractal interpolation functions. SIAM. J. Numer. Anal., 44: 655-676.
- Dalla, L., 2002. Bivariate fractal interpolation functions on grids. Fractals, 10: 53-58.
- Feng, Z., Y. Feng and Z. Yuan, 2012. Fractal interpolation surfaces with function vertical scaling factors. Appl. Math. Lett., 25: 1896-1900.
- Hefei, Z., T. Ran, Z. Siyong and W. Yue, 1999. Wavelet-based fractal function approximation. J. Syst. Eng. Electron., 10: 60-66.
- Kang, Y.S., C.H. Yun and D.H. Kim, 2014. A construction of the best fractal approximation. Electron. J. Math. Anal. Appl., 2: 144-151.
- Lonardi, S. and P. Sommaruga, 1999. Fractal image approximation and orthogonal bases. Signal Process. Image Commun., 14: 413-423.
- Massopust, P.R., 1997. Fractal functions and their applications. Chaos Solitons Fractals, 8: 171-190.
- Malysz, R., 2006. The Minkowski dimension of the bivariate fractal interpolation surfaces. Chaos Solitons Fractals, 27: 1147-1156.
- Metzler, W. and C.H. Yun, 2010. Construction of fractal interpolation surfaces on rectangular grids. Intl. J. Bifurcation Chaos, 20: 4079-4086.
- Navascues, M.A. and M.V. Sebastian, 2004. Generalization of Hermite functions by fractal interpolation. J. Approximation Theory, 131: 19-29.
- Yun, C.H., C.O. Hyong and H.C. Choi, 2014. Construction of fractal surfaces by recurrent fractal interpolation curves. Chaos Solitons Fractals, 66: 136-143.