

## QSAOR Iterative Method for the Solution of Time-Fractional Diffusion Equation

<sup>1</sup>Baharuddin, <sup>2</sup>Andang Sunarto and <sup>3</sup>Juhriyansyah Dalle

<sup>1</sup>State University of Medan, Medan, Indonesia

<sup>2</sup>State Institute for Islamic Studies of Bengkulu, Bengkulu, Indonesia

<sup>3</sup>University of Lambung Mangkurat, Banjarmasin, Indonesia

**Abstract:** The main purpose of this study is to investigate the performance of the Quarter-Sweep AOR (QSAOR) iterative method by using Caputo fractional derivative operator and implicit finite difference scheme to solve time-fractional diffusion equation. To solve the problems, a linear system will be constructed via discretization of the one-dimensional linear time-fractional diffusion equations by using the Caputo's fractional derivative. Then the generated linear system has been solved using the proposed QSAOR iterative method. In the addition, the formulation and application of the QSAOR method to solve the problems are also presented. Two numerical examples and comparison with FSAOR and HSAOR methods are given to show the effectiveness of QSAOR method.

**Key words:** QSAOR, Caputo, implicit, time-fractional, diffusion

### INTRODUCTION

Many science and engineering problems can be modeled mathematically as differential equations (Oldham and Spanier, 1974; Podlubny, 1999a). Since, the rapid growth of computer technology, the numerical techniques are used to solve a large size problem. Then, there are some numerical methods proposed to solve the Time Fractional Diffusion Equations (TFDE) such as Crank-Nicolson finite difference method (Sweilam *et al.*, 2012), two implicit finite method (Ma, 2014) and high-order finite element methods (Jiang and Ma, 2011). In this research, one dimensional time-fractional diffusion equation will be solved numerically and represented as follows:

$$\frac{\partial^\alpha U(x, t)}{\partial x^\alpha} = a(x) \frac{\partial^2 U(x, t)}{\partial x^2} + b(x) \frac{\partial U(x, t)}{\partial x} + c(x)U(x, t) \quad (1)$$

With boundary conditions  $u(0, t) = g_0(t)$ ,  $U(l, t) = g_1(t)$  and the initial condition  $U(x, 0) = f(x)$ . To get the approximate solution of the Time Fractional Diffusion Equation (TFDE) problem the problem needs to be discretized to form an approximation equation. Based on the implicit finite difference scheme and Caputo fractional derivative operator, the approximation equations can be derived to construct a linear system at each time level. For to solving linear systems, many researchers have also discussed several concepts of the iterative methods such as Young (1971), Hackbusch (1995) and Saad (1996). In addition to these iterative methods, Abdullah (1991) initiated Half-Sweep iteration which is one of the

most known and widely used iterative techniques to solve any linear systems. Differently from the Half-Sweep iteration approach, Othman and Abdullah (2000) have expanded this approach to initiate the Modified Explicit Group (MEG) method based on the quarter-sweep approach. In this study, we propose the QSAOR iterative method for solving time-fractional fractional diffusion equations based on the Caputo's implicit finite difference approximation equation. To demonstrate the capability of the QSAOR method, we also implement the Full-Sweep AOR (FSAOR) and Half-Sweep AOR (HSAOR) iterative methods being used as a control method.

Before discretizing Problem (1), let us consider some definitions that can be applied for fractional derivative theories in order to construct the approximation equation of Problem (1).

**Definition 1:** The Riemann-Liouville fractional integral operator,  $J^\alpha$  of order- $\alpha$  is defined as Podlubny (1999b), Kilbas *et al.* (2006) and Zhang (2009):

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \alpha > 0, x > 0 \quad (2)$$

**Definition 2:** The Caputo's fractional partial derivative operator,  $D^\alpha$  of order- $\alpha$  is defined as Kilbas *et al.* (2006) and Caputo (1967):

$$D^\alpha f(x) = \frac{1}{\Gamma(m-\alpha)} \int_0^x \frac{f^{(m)}(t)}{(x-t)^{\alpha-m+1}} dt, \alpha > 0 \quad (3)$$

with  $m-1 < \alpha \leq m$ ,  $m \in \mathbb{N}$ ,  $X > 0$ . From Definitions 1 and 2,  $\Gamma(\alpha)$  is known as a gamma function which is given by:

$$\Gamma(\alpha) = \int_0^{\infty} x^{\alpha-1} e^{-x} dx$$

For obtain the numerical solution of Time-Fractional Diffusion Equation (TFDE's) in Eq. 1 we get numerical approximation equations by using the Caputo's derivative definition with Dirichlet boundary conditions and consider the non-local fractional derivative operator. This approximation equation can be categorized as unconditionally stable scheme. On strength of Problem (1), the solution domain of the problem has been restricted to the finite space domain  $0 \leq x \leq \gamma$  with  $0 < \alpha < 1$  whereas the parameter  $\alpha$  refers to the fractional order of space derivative. In order to solve Problem (Eq. 1), let us consider boundary conditions of Problem (Eq. 1) be given as:

$$U(0, t) = g_0(t), U(l, t) = g_l(t)$$

and the initial condition:

$$U(x, 0) = f(x)$$

where,  $g_0(t)$ ,  $g_l(t)$  and  $f(x)$  are given functions. Based on a discretize approximation to the time fractional derivative in Eq. 1, we consider Caputo's fractional partial derivative of order  $\alpha$ , defined by Hadjidimos (1978) and Saad (1996):

$$\frac{\partial^\alpha u(x, t)}{\partial t^\alpha} = \frac{1}{\Gamma(n-1)} \int_0^t \frac{\partial u(x-s)}{\partial t} (t-s)^{-\alpha} ds, t > 0, 0 < \alpha < 1 \tag{4}$$

**Quarter-sweep implicit finite difference approximation:**  
By using Eq. 4, the formulation of Caputo's fractional partial derivative of the first order approximation method is given as:

$$D_t^\alpha U_{i,n} \equiv \sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) \tag{5}$$

and we have the following expressions:

$$\sigma_{\alpha,k} = \frac{1}{\Gamma(1-\alpha)(1-\alpha)k^\alpha}$$

And:

$$\omega_j^{(\alpha)} = j^{1-\alpha} - (j-1)^{1-\alpha}$$

Previously to discretize Problem (1), let the solution domain of the problem be partitioned uniformly. To do this we consider some positive integers  $m$  and  $n$  in which the grid sizes in space and time directions for the finite difference algorithm are defined as  $h = \Delta x = \gamma/m$  and  $k = \Delta t = T/n$ , respectively. Based on these grid sizes, we develop the uniformly grid network of the solution domain where the grid points in the space interval  $[0, \gamma]$  are shown as the numbers  $x_i = ih$ ,  $i = 0, 1, 2, \dots, m$  and the grid points in the time interval  $[0, T]$  are labeled  $t_j = jk$ ,  $j = 0, 1, 2, \dots, n$ . Then, the values of the function  $U(x, t)$  at the grid points are denoted as  $U_{i,j} = U(x_i, t_j)$ . According to Eq. 5 and using the implicit finite difference discretization scheme, the quarter-sweep implicit finite difference approximation equation of Problem in Eq. 1 to the grid point centered at  $(x_i, t_j) = (ih, jk)$  is given as:

$$\sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) = \alpha_i \frac{1}{16h^2} (U_{i-4,n} - 2U_{i,n} + U_{i+4,n}) + b_i \frac{1}{8h} (U_{i+4,n} - U_{i-4,n}) + c_i U_{i,n} \tag{6}$$

For  $i = 4, 8, \dots, m-4$ . Based on Eq. 6 this approximation equation is known as the fully implicit finite difference approximation equation which is consistent first order accuracy in time and second order in space. Particularly, the approximation Eq. 6 can be rewritten based on the specified time level. For instance, we have for  $n \geq 2$ :

$$\sigma_{\alpha,k} \sum_{j=1}^n \omega_j^{(\alpha)} (U_{i,n-j+1} - U_{i,n-j}) = p_i U_{i-4,n} + q_i U_{i,n} + r_i U_{i+4,n} \tag{7}$$

Where:

$$p_i = \frac{\alpha_i}{16h^2} = \frac{b_i}{8h}$$

$$q_i = c_i = \frac{\alpha_i}{8h^2}$$

$$r_i = \frac{\alpha_i}{16h^2} + \frac{b_i}{8h}$$

Also, we get for  $n = 1$ :

$$-p_i U_{i-4,1} + q_i^* U_{i,1} - r_i U_{i+4,1} = f_{i,1} \quad i = 4, 8, \dots, m-4 \tag{8}$$

Where:

$$\omega_j^{(\alpha)} = 1$$

$$q_i^* = \sigma_{\alpha,k} - q_i$$

$$f_{i,1} = \sigma_{\alpha,k} - U_{i,1}$$



Table 1: Comparison between No. of iterations (K), the execution time (sec) and maximum errors for the iterative methods using examples 1 at  $\alpha = 0.25, 0.50, 0.75$

M	Methods	$\alpha = 0.25$			$\alpha = 0.50$			$\alpha = 0.75$		
		K	Time	Max. error	K	Time	Max. error	K	Time	Max. error
128	FSAOR	657	1.93	9.95e-05	615	1.83	9.84e-05	613	1.79	1.29e-04
	HSAOR	517	1.78	9.95e-05	510	1.54	9.84e-05	599	1.39	1.29e-04
	QSAOR	234	0.91	9.9e-05	396	1.02	9.84e-05	337	1.03	1.29e-04
256	FSAOR	870	6.38	9.95e-05	659	2.99	9.84e-05	630	1.72	1.29e-04
	HSAOR	857	3.02	9.95e-05	425	2.93	9.84e-05	413	1.50	1.29e-04
	QSAOR	255	1.11	9.95e-05	215	1.36	9.84e-05	212	1.33	1.29e-04
512	FSSOR	4940	31.35	9.96e-05	3080	22.62	9.84e-05	1301	16.67	1.29e-04
	FSAOR	870	6.54	9.05e-05	859	6.37	9.84e-05	830	6.45	1.29e-04
	QSAOR	452	2.31	9.95e-05	417	2.23	9.84e-05	448	2.27	1.29e-04
1024	FSSOR	19033	45.72	9.97e-05	12232	101.21	9.80e-05	5911	105.27	1.30e-04
	HSAOR	4940	33.28	9.96e-05	3080	42.68	9.80e-05	1301	17.87	1.29e-04
	QSAOR	2673	20.97	9.96e-05	1664	13.26	9.84e-05	698	5.95	1.29e-04
2048	FSSOR	70547	1142.09	1.00e-04	45700	1342.00	9.89e-05	22474	1002.85	1.30e-04
	HSAOR	19033	512.59	1.00e-04	12232	338.14	9.85e-05	5911	157.02	1.30e-04
	QSAOR	10332	158.37	9.96e-05	6633	100.86	9.85e-05	3200	49.16	1.30e-04

Table 2: Comparison between number of iterations (K), the execution time (sec.) and maximum errors for the iterative methods using examples 2 at  $\alpha = 0.25, 0.50, 0.75$

M	Methods	$\alpha = 0.25$			$\alpha = 0.50$			$\alpha = 0.75$		
		K	Time	Max. error	K	Time	Max. error	K	Time	Max. error
128	FSAOR	370	6.19	1.94e-02	260	5.86	8.29e-02	257	5.80	1.37e-01
	HSAOR	199	5.11	1.95e-02	154	5.04	8.30e-02	146	5.02	1.37e-01
	QSAOR	72	2.79	1.95e-02	61	2.05	8.30e-02	48	2.25	1.37e-01
256	FSAOR	1761	19.30	1.95e-02	809	11.82	8.29e-02	291	7.39	1.37e-01
	HSAOR	545	6.25	1.95e-02	328	5.70	8.29e-02	163	5.29	1.37e-01
	QSAOR	166	3.02	1.95e-02	116	2.39	8.30e-02	72	2.40	1.37e-01
512	FSSOR	6746	125.09	1.94e-02	3240	63.93	8.29e-02	1471	30.79	1.37e-01
	FSAOR	2246	19.45	1.95e-02	1402	14.27	8.29e-02	625	9.09	1.37e-01
	QSAOR	849	7.63	1.95e-02	629	6.97	8.30e-02	279	4.77	1.37e-01
1024	FSSOR	25054	866.80	1.94e-02	12126	435.44	8.29e-02	5644	208.20	1.37e-01
	HSAOR	8478	117.00	1.94e-02	5313	76.75	8.29e-02	2461	38.54	1.37e-01
	QSAOR	3301	28.86	1.95e-02	2425	23.05	8.29e-02	1099	13.21	1.37e-01
2048	FSSOR	91984	6092.40	1.94e-02	44563	1428.60	8.29e-02	20921	1609.51	1.37e-01
	HSAOR	40541	1109.00	1.94e-02	19650	557.72	8.29e-02	9198	268.45	1.37e-01
	QSAOR	18209	272.76	1.94e-02	8832	139.41	8.29e-02	4126	68.64	1.37e-01

$$U(x, t) = \sum_{n=0}^{m-1} \frac{\partial^n U(x, 0) t^n}{\partial t^n n!} + \sum_{n=1}^{\infty} \sum_{i=0}^{m-1} \frac{\partial^{mn+i} U(x, 0) t^{n\alpha+i}}{\partial t^{mn+i} \Gamma(n\alpha+i+1)} \quad U(x, 0) = x^2 \tag{18}$$

To  $U(x, t)$  for  $0 < \alpha < 1$  it can be shown that the analytical solution of Problem (16) is given as:

$$U(x, t) = x^2 + 2k \frac{t^\alpha}{\Gamma(\alpha + 1)}$$

**Examples 2:** Let us consider the following time fractional initial boundary value problem be defined as Ali *et al.* (2013):

$$\frac{\partial^\alpha U(x, t)}{\partial t^\alpha} = \frac{1}{2} x^2 \frac{\partial^2 U(x, t)}{\partial x^2}, \quad 0 < \alpha \leq 1, \quad 0 \leq x \leq \gamma, \quad t > 0 \tag{16}$$

where the boundary conditions are given in fractional terms:

$$U(0, t) = 0, \quad U(1, t) = e^t \tag{17}$$

and the initial condition:

From Problem (19) as taking  $\alpha = 1$ , it can be shown that Eq. 19 can also be reduced to the standard diffusion equation:

$$\frac{\partial U(x, t)}{\partial t} = \frac{1}{2} x^2 \frac{\partial^2 U(x, t)}{\partial x^2}, \quad 0 \leq x \leq \gamma, \quad t > 0 \tag{19}$$

Then, the analytical solution of Problem (17) is obtained as follows:

$$U(x, t) = x^2 e^t$$

Now by applying the series:

$$U(x, t) = \sum_{n=0}^{m-1} \frac{\partial^n U(x, 0) t^n}{\partial t^n n!} + \sum_{n=1}^{\infty} \sum_{i=0}^{m-1} \frac{\partial^{mn+i} U(x, 0) t^{n\alpha+i}}{\partial t^{mn+i} \Gamma(n\alpha+i+1)}$$

to  $U(x, t)$  for  $0 < \alpha < 1$  it can be shown that the analytical solution of Problem (17) is stated as (Table 1 and 2):

$$U(x, t) = x^2 \left[ 1 + \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots \right]$$

All results of numerical experiments for Problems (12) and (15), obtained from implementation of FSAOR, HSAOR and QSAOR iterative methods are recorded in Table 1 and 2 at different values of mesh sizes,  $M = 128, 256, 512, 1024$  and  $2048$ .

### CONCLUSION

As a conclusion for the numerical solution of the time-fractional diffusion problems this study deals with the implementation of QSAOR iterative method to solve a linear system generated by the Quarter-Sweep Caputo's implicit approximation equations. Through numerical experiments results from Table 1 and 2, clearly it demonstrates that two promising improvements in the number of iteration ( $K$ ) and execution time with implementing a QSAOR iterative method have been shown as compared to the FSAOR and HSAOR methods. Overall, the numerical results showed that the quarter-sweep iteration concept in association with the AOR iterative method is superior and it has reduced the computational complexity significantly. Therefore further investigation of two-step (Evans and Sahimi, 1988; Ruggiero and Galligani, 1990; Sahimi *et al.*, 1993) iterative methods should be considered for future works.

### REFERENCES

Abdullah, A.R., 1991. The four point Explicit Decoupled Group (EDG) method: A fast poisson solver. *Int. J. Comput. Math.*, 38: 61-70.

Ali, S.E., B. Ozgur and E. Korkmaz, 2013. Analysis of fractional partial differential equations by Taylor series expansion. *Boundary Value Prob.*, 24: 68-81.

Caputo, M., 1967. Linear models of dissipation whose  $Q$  is almost frequency independent-II. *Geophys. J. Intl.*, 13: 529-539.

Evans, D.J. and M.S. Sahimi, 1988. The Alternating Group Explicit iterative method (AGE) to solve parabolic and hyperbolic partial differential equations. *Annu. Rev. Numer. Fluid Mech. Heat Transfer*, 2: 283-389.

Hackbusch, W., 1995. *Iterative Solution of Large Sparse Systems of Equations*. Springer, New York, USA., ISBN:9783540940647, Pages: 429.

Hadjidimos, A., 1978. Accelerated over-relaxation method. *Math. Comput.*, 32: 149-157.

Jiang, Y. and J. Ma, 2011. High-order finite element methods for time: Fractional partial differential equations. *J. Comput. Appl. Math.*, 235: 3285-3290.

Kilbas, A.A., H.M. Srivastava and J.J. Trujillo, 2006. *Theory and Applications of Fractional Differential Equations*. Elsevier, New York, USA., ISBN: 0-444-51832-0, Pages: 502.

Ma, Y., 2014. Two implicit finite difference methods for time-fractional diffusion equation with source term. *J. Appl. Math. Bioinf.*, 2: 125-145.

Oldham, K.B. and J. Spanier, 1974. *The Fractional Calculus*. Academic Press, New York, USA.,.

Othman, M. and A.R. Abdullah, 2000. An efficient four points modified explicit group poisson solver. *Int. J. Comput. Math.*, 76: 203-217.

Podlubny, I., 1999a. *Fractional Differential Calculus*. Academic Press, New York, USA.,.

Podlubny, I., 1999b. *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations to Methods of their Solution and Some of their Applications*. Academic Press, New York, USA., ISBN:9780125588409, Pages: 340.

Ruggiero, V. and E. Galligani, 1990. An iterative method for large sparse systems on a vector computer. *Comput. Mathematics Appl.*, 20: 25-28.

Saad, Y., 1996. *Iterative Method for Sparse Linear Systems*. International Thomas Publishing, Boston, Massachusetts.,

Sahimi, M.S., A. Ahmad and A.A. Bakar, 1993. The Iterative Alternating Decomposition Explicit (IADE) method to solve the heat conduction equation. *Intl. J. Comput. Math.*, 47: 219-229.

Sunarto, A., J. Sulaiman and A. Saudi, 2014a. Half-sweep accelerated over-relaxations iterative method for the solution time-fractional diffusion equations. *Proceedings of the 22nd National Symposium on Mathematical Science*, November 24-26, 2014, Grand Bluewave Hotel Shah Alam, Shah-Alam, Malaysia, pp: 109-115.

Sunarto, A., J. Sulaiman and A. Saudi, 2014b. Implicit finite difference solution for time-fractional diffusion equations using AOR method. *J. Phys. Conf. Ser.*, 495: 2024-2031.

Sweilam, N.H., M.M. Khader and A.M.S. Mahdy, 2012. Crank-Nicolson finite difference method for solving time-fractional diffusion equation. *J. fractional calculus appl.*, 2: 1-9.

Tian, H., 2003. Accelerated over-relaxation method for rank deficient linear systems. *Appl. Math. Comput.*, 14: 485-499.

Young, D.M., 1971. *Iterative Solution of Large Linear Systems*. Academic Press, London, England, Pages: 570.

Zhang, Y., 2009. A finite difference method for fractional partial differential equation. *Appl. Math. Comput.*, 215: 524-529.