

The Dynamics of the Fixed Points to Modified Jerk Map

Samah Abdulhadi Al-Hashemi, Zainab Abdulmunim Sharba and Najlaa Adnan

Department of Computer Science, College of Science for Women, University of Babylon, Hillah, Iraq

Abstract: Recently, Jerk equation which is the third-order explicit autonomous differential equation is noticed to be a motivating sub-class of dynamical systems. Where many of regular and chaotic motion features can be reveal from these systems. In this study, a simplified version Jerk map is presented. Different properties of dynamical behavior is acquired by replacing three dimensional systems to two dimensional one. Where a new parameter is added with the same properties. Moreover, we study the fixed point of modified Jerk map with the form $M_{j_{ab}} = (y-ax+by^2) = (xy)$ and the general properties of them, so, we find the contracting and expanding area of this map. Also, we determine fixed point attracting repelling or saddle.

Key words: Modified Jerk map, fixed point, attracting-expanding area, dimensional, properties, saddle

INTRODUCTION

Non-linear dynamical happens and covered a large area in engineering, physics, biology and many other scientific disciplines. There is a large amount of interest in the chaos literature in found out of chaos in natural and physical system. Poincare way the first to notice the possibility of chaos according to which a deterministic system to show periodic behavior that depends on the initial condition. There by rendering long term could be expected impossible.

Three dimensional system which is simple and chaotic have been tried to find by Gottlieb (1996), Hoover (1995) Linz (1997) Patidar and Sud (2005) and Posch *et al.* (1986). By Sprott (1994) found 19 clear chaotic models (one of them is conservative and the lastings are dissipative, mentioned as Models A-S with three-dimensional vector fields that involve of five terms including two non-linearities or of six terms with non-linearity one quadratic.

Hoover (1995) stated that the special case of nose that has been created by Sprott (1994) is simply a conservative system (Model A) which is a known Hoover thermostat dynamical system. This system shows a time reversible Hamiltonian chaotic (Linz, 1997). Apart from this the others models from B-S are unknown. Moreover, Gottlieb indicated that 'Jerk function' is an noticeable third-order form $y = J(Y, \dot{Y}, \ddot{Y})$ produces from reorder the Sprott's Model A. This function contains the third derivative of x . According to Gottlieb (1996) study, exciting question has been annoyance "what is the simplest Jerk function that gives chaos?"

Linz (1997) shows that three models can be reduce to a Jerk form named original Rossler Model, Lorenz

Model and Sprott's Model R. Moreover, the difficulty of Rossler and Lorenz Models are higher and inappropriate candidate for the Gottlieb's simplest Jerk function.

In Patidar and Sud (2005), the global dynamic of some member of special family of dynamic system has been explored by Patidar. The following form represents this dynamic system under consideration:

$$\dot{Y} + AY + \ddot{Y} = G(X)$$

Where, the system parameter represented by A and the non-linear function that have three arguments which are one nonlinearity, one systems parameter and a constant term represented by $G(X)$.

In this research, we simplified the Jerk map by replacing three dimensional systems to two dimensional systems and include the new parameter, so, we get the difference (new various) properties of dynamical behavior from Jerk map also there exist the same properties of it. Also, we show that the modified Jerk map is diffeomorphism map and it has two fixed point, we determine the type of fixed points.

MATERIALS AND METHODS

Preliminaries: "For any map G define from \mathbb{R}^2 to \mathbb{R}^2 , we say G is C^k if its continuous for all $k \in \mathbb{Z}^+$ and its mixed K -th partial derivatives exist and we say that G is diffeomorphism map if it is onto, one to one, C^k and its inverse is C^k . Let U be a subset of \mathbb{R}^2 and v_0 be any element in \mathbb{R}^2 . Consider $G: V \rightarrow \mathbb{R}^2$ be a map. Furthermore, assume that the first partials of the coordinate maps f and g of G exist at s_0 . The differential of g at s_0 is the linear map $DG(s_0)$ defined on \mathbb{R}^2 by:

$$DG(s_0) = \begin{pmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \\ \frac{\partial g}{\partial x} & \frac{\partial g}{\partial y} \end{pmatrix}$$

for all the s_0 in R^2 . The determinate of $DG(s_0)$ is called the Jerk of G at s_0 and it is denoted by $JG(s_0) = \det DG(s_0)$, so, G is said to be area-contracting at v_0 if $|\det DG(v_0)| < 1$, G is said to be area-expanding at v_0 if $|\det DG(v_0)| > 1$.

For any $n \times n$ matrix B , the scalar λ is called aneigenvalue of $n \times n$ matrix B if there exists a non zero vector $Z \in R^n$ such that $BZ = \lambda Z$ such an Z is called eigenvector of B corresponding to eigen value λ . Any pair for which $f(pq) = p.g(pq) = q$ is called a fixed point of the two dimensional dynamical system it is repelling fixed point if λ_1 and λ_2 are >1 in absolute value and it is an attracting fixed point if λ_1 and λ_2 is <1 in absolute value $B \in GL(2, Z)$ with $\det(B) = \pm 1$ is called hyperbolic matrix if $|\lambda_i|$ wher λ_i are the eigenvalues of B (Denny, 1992).

RESULTS AND DISCUSSION

General properties of modified jerk map

Proposition (3-1): Let $MJ: R^2 \rightarrow R^2$ be modified Jerk map then MJ has tow fixed points:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

Proof: By definition of fixed point:

$$MJ_{ab} = \begin{pmatrix} y \\ -ax + by^2 \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$$

Then $x = y$ and $-Ay + By^2 - y = 0$, $by^2 - (a+1)y = 0$, $y[by - (a+1)] = 0$ then $y = 0$ or $y = a+1/b$ and $x = a+1/b$. Therefore:

$$P_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ and } P_2 = \begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

Proposition (3-2): Let $MJ: R^2 \rightarrow R^2$ be modified Jerk map then the Jacobin of $MJ_{a,b}$ is a:

Proof:

$$DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -a & 2by \end{pmatrix} \text{ so, } DMJ_{a,b} \begin{pmatrix} x \\ y \end{pmatrix} = \det \begin{pmatrix} 0 & 1 \\ -a & 2by \end{pmatrix} = a$$

Proposition (3-3): The eigen value of:

$$DMJ_{a,b} \begin{pmatrix} x \\ y \end{pmatrix} \text{ are } \lambda_{1,2} = by \mp \sqrt{(by)^2 - a} \text{ at } p_2, \forall \begin{pmatrix} x \\ y \end{pmatrix} \in R^2$$

Proof: If λ is eigen value of DMJ_{ab} then must be satisfied the following equation:

$$\begin{vmatrix} -\lambda & 1 \\ -a & 2by - \lambda \end{vmatrix} = 0$$

Then:

$$\begin{aligned} (-\lambda)(2by - \lambda) + a &= 0 \\ \lambda^2 - 2by\lambda + a &= 0 \end{aligned}$$

And the solution of characteristic equations are:

$$\begin{aligned} \lambda_{1,2} &= \frac{2by \mp \sqrt{(2by)^2 - 4a}}{2} \\ \lambda_{1,2} &= by \mp \sqrt{(by)^2 - a} \end{aligned}$$

Remark:

- It is clear that if $|by| > \sqrt{a}$ then the eigen value of $DMJ_{a,b}$ is real
- The eigen value of $DMJ_{a,b}$ at fixed point P_1 are $\lambda_{1,2}^0 = \mp \sqrt{-a}$ and $\lambda_{1,2} = a+1 \mp \sqrt{(a+1)^2 - a}$ for the fixed point P_2

Proposition (3-4): Let $MJ: R^2 \rightarrow R^2$ be modified Jerk map if $a \neq 0$ then $Mj_{a,b}$ is diffeomorphism.

Proof: To prove is one to one. Let:

$$\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \in R$$

Such that:

$$MJ_{a,b} \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}$$

Then:

$$\begin{pmatrix} y_1 \\ -ax_1 + by_1^2 \end{pmatrix} = \begin{pmatrix} y_2 \\ -ax_2 + by_2^2 \end{pmatrix}$$

Then:

$$y_1 = y_2 \text{ and } -ax_1 + by_1^2 = -ax_2 + by_2^2$$

Hence, $-ax_1 = -ax_2$ and then $x_1 = x_2$, $Mj_{a,b}$ is C^∞ :

$$MJ_{a,b} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ -ax + by^2 \end{pmatrix}$$

then all first partial derivative exist and continuous, note that:

$$\frac{\partial^n f(x,y)}{\partial x^n} = 0 \forall n \in \mathbb{N}$$

And:

$$\frac{\partial^n f(x,y)}{\partial y^n} = 0 \forall n \geq 2$$

$$\frac{\partial^n g(x,y)}{\partial x^n} = 0 \forall n \geq 3$$

We find all its $M_{j_{a,b}}$ exist kth partial derivative exist and continuous for all k. $M_{j_{a,b}}$ is onto:

$$\text{Let } \begin{pmatrix} v \\ w \end{pmatrix} \in \mathbb{R}^2$$

Such that:

$$y = v \text{ and } -Ax + By^2 = w$$

So:

$$x = \frac{w - bv^2}{-a}$$

Then, there exist:

$$\begin{pmatrix} \frac{bv^2 - w}{a} \\ v \end{pmatrix} \in \mathbb{R}^2$$

Such that:

$$MJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} v \\ w \end{pmatrix}$$

Then, MJ_{ab} is onto, MJ_{ab} has an invers.

Remark: If $a = 0$ then:

$$MJ_{ab} = \begin{pmatrix} y \\ by^2 \end{pmatrix}$$

So:

$$\text{Ker}(MJ_{ab}) = \begin{pmatrix} x \\ y \end{pmatrix} : x,y \in \mathbb{R}$$

Then, MJ_{ab} is not one to one hence, MJ_{ab} is not diffeomorphism.

Proposition (3-5):

$$DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix}$$

is non hyperbolic if $\mp \frac{\sqrt{a}}{b}$ also it is non hyperbolic matrix if $a = -1$.

Proof: Let $y = \mp \frac{\sqrt{a}}{b}$ then $b^2y^2 = a$ and $b^2y^2 - 2by\sqrt{a} + a = (by)^2 - \sqrt{a}$
Hence:

$$x = \left| by \mp \sqrt{(by)^2 - a} \right| = 1$$

So, $\lambda_1 = \lambda_2 = 1$, $\lambda_{1,2}^0 = \mp \sqrt{-a}$ the eigenvalue at p_1 .
If $a = -1$ then $\lambda_{1,2}^0 = \mp 1$.

Proposition 3-6: For all:

$$\begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2, y \neq \frac{\sqrt{a}}{b}$$

And:

$$|by| > \sqrt{a}. DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix}$$

is hyperbolic matrix if and only if and only if $|a| = 1$.

Proof: Since, $DMJ_{ab} \in GL(2, \mathbb{Z})$ and:

$$\det \left(DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} \right) = |a|$$

then $a = 1$ or $a = -1$.

Conversely: Let $a = 1$, since:

$$\det \left(DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} \right) = a$$

we have $\lambda_1 \lambda_2 = a = 1$. Then, $\lambda_1 = 1/\lambda_2$ if $\lambda_1 > 1$ then $\lambda_2 < 1$ or $\lambda_1 < 1$ then $\lambda_2 > 1$. So, DMJ_{ab} is hyperbolic matrix.

Proposition (3-7): If $|a| < 1$ then DMJ_{ab} is area-contracting and its area expanding if $|a| > 1$.

Proof: If $|a| < 1$ then by proposition (Jorobian):

$$\left| \det \left(DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} \right) \right| < 1$$

that is DMJ_{ab} is area -contracting map and if $|a| > 1$ then:

$$\left| \det \left(DMJ_{ab} \begin{pmatrix} x \\ y \end{pmatrix} \right) \right| > 1$$

This implies that MJ_{ab} is an area-expending.

Fixed point properties of modified Jerk map

Proposition (4-1): If $|a| > 1$ then the fixed point:

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

of modified Jerk map is repelling where $a > 1$. The fixed point:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

of modified Jerk map is repelling where $a < -1$.

Proof (1): Since:

$$+1 > a, \sqrt{(a+1)^2 - a} > \sqrt{(a-2)^2}$$

Then:

$$a+1 \mp \sqrt{(a+1)^2 - a} > a \mp \sqrt{(a-2)^2}$$

By proposition (3-1 and 3-3). Thus, $|\lambda_1| = |\lambda_2| > 1$. Then:

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

is repelling fixed point.

Proof (2): Since, $a < -1$ then, we have $\sqrt{-a} > 1$ by proposition (2-1 and 2-3) $|\lambda_1^0| = |\lambda_2^0| > 1$. Hence:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

is repelling fixed point.

Proposition (4-2): If $|a| < 1$ then, MJ_{ab} has attracting fixed point:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

and saddle fixed point:

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix} \text{ where } a > 1$$

$MJ_{a,1D^{467b}}$ has attracting fixed point:

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix} \text{ where } a < 1$$

Proof (1): Since, $-1 < a$ then, we have $\sqrt{-a} < 1$ and $|\lambda_1^0| = |\lambda_2^0| < 1$ by proposition (3-1) (3-3). Hence:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

is attracting fixed point.

Now to show that p_2 is a saddle fixed point by proposition 2-3, we have two eigenvalue:

$$\lambda_1 = a+1+\sqrt{(a+1)^2 - a}, \lambda_2 = a+1-\sqrt{(a+1)^2 - a} \quad (1)$$

Since:

$$\sqrt{a^2+a+1} < \sqrt{(a+1)^2} = |a+1| = -(a+1) \quad (2)$$

Hence:

$$a+1\sqrt{a^2+a+1} < 1 \quad (3)$$

So:

$$|a+1+\sqrt{a^2+a+1}| = |\lambda_1| < 1 \quad (4)$$

For λ_2 since, $-1 < a < 0$, we have:

$$\sqrt{a^2+a+1} > \sqrt{a^2} = |a| = -a \quad (5)$$

By adding (a+1) for both side, we get:

$$a+1-\sqrt{(a+1)^2 - a} > 1 \text{ so, } |\lambda_2| > 1 \quad (6)$$

From Eq. 3-5 and by proposition (3-1), we obtain:

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

is a saddle fixed point.

Proof 2: Since:

$$\sqrt{a^2+a+1} < \sqrt{(a+1)^2} = |a+1| = -(a+1)$$

hence by adding (a+1) for both side, we get:

$$a+1+\sqrt{a^2+a+1} < 1 \tag{6}$$

$$\sqrt{a^2+a+1} > \sqrt{a^2} = a \text{ so, } -\sqrt{a^2+a+1} < -a$$

By adding (a+1) for both side, we get:

$$a+1-\sqrt{(a+1)^2-a} < 1 \tag{7}$$

From Eq. 6 and 7 and by proposition (3-1), we obtain

$$\begin{pmatrix} \frac{a+1}{b} \\ \frac{a+1}{b} \end{pmatrix}$$

is attracting fixed point.

CONCLUSION

In this study, a simplified version Jerk map is presented. Different properties of dynamical behavior is acquired by replacing three dimensional systems to two dimensional one. Where a new parameter is added with the same properties.

For this class of dynamical system the improved Jerk map becomes a representative. The main mathematical

properties of the modified Jerk map is formulated. Two fixed point is gotten P₁ and P₂ If b≠0. The eignvalues of the modified Jerk map is λ_{1,2}⁰ = ∓√-a at P₁ and λ_{1,2} = a+1±√(a+1)²-a at P₂, if |a| <1 then DMJ_{a,b} is an area contraction map and if |a|>1 then DMJ_{a,b} is an area expanding map. Moreover, the improve Jerk map is shown to be a diffeomorphism map.

REFERENCES

Denny, G., 1992. Encounters with Chaos. McGraw Hill, New York, USA.,
 Gottlieb, H.P.W., 1996. Question#38: What is the simplest jerk function that gives chaos?. Am. J. Phys., 64: 525-525.
 Hoover, W.G., 1995. Remark on some simple chaotic flows. Phys. Rev. E., 51: 759-760.
 Linz, S.J., 1997. Nonlinear dynamical models and jerky motion. Am. J. Phys., 65: 523-526.
 Patidar, V. and K.K. Sud, 2005. Bifurcation and chaos in simple jerk dynamical systems. Pramana. J. Phys., 64: 75-93.
 Posch, H.A., W.G. Hoover and F.J. Vesely, 1986. Canonical dynamics of the Nose oscillator: Stability, order and chaos. Phys. Rev. A., 33: 4253-4265.
 Sprott, J.C., 1994. Some simple chaotic flows. Phys. Rev. E., 50: R647-R650.