

## Comparison of Nutrients Uptake in Different Varieties of Rice in Pakistan

<sup>1</sup>Hamid ur Rehman, <sup>1</sup>Fahad Akhtar, <sup>2</sup>Qazi Inam ul Haq, <sup>1</sup>Safi Ullah, <sup>1</sup>Azhar Abbas,  
<sup>1</sup>Usman Shoukat, <sup>5</sup>Faridah Kormin, <sup>2</sup>Tauseef Ahmad, <sup>3</sup>Suleman Khan and <sup>4</sup>Muhammad Khan  
<sup>1</sup>Department of Biochemistry, <sup>2</sup>Department of Microbiology,  
<sup>3</sup>Department of Genetics, <sup>4</sup>Centre for Human Genetics,  
Hazara University Mansehra, 21300 Mansehra, KPK, Pakistan  
<sup>5</sup>Faculty of Science, Technology and Human Development,  
University Tun Hussein Onn Malaysia, Parit Raja,  
Batu Pahat, 86400 Johor, Malaysia

**Abstract:** The objective of this research was to determine the uptake of different elements (K, Mg, Ca, Na, Fe, Co, Mn, Pb, Cu, Zn and Ni) which are used as nutrients by the rice crop. Different types of rice seeds KSK-282, KSK-134, BAS-515, IR6, BAS-2000, KSK-133 and super-BAS were collected from National Agricultural Research Centre (NARC) Islamabad Pakistan and grown in different plastic pots containing soil under the same condition. Ten days old immature plants of different varieties of rice were dried and their roots were separated from the whole plants. The dried roots were ground into fine powder followed by acid digestion (HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub> and HClO<sub>4</sub>) solution in a ratio of (5:1:0.1) individually. After digestion the solutions were filtered and the filtrates were diluted by adding distilled water. The diluted solutions of all the above mentioned varieties of rice were analyzed by Atomic Absorption Spectrophotometer (AAS) for nutrients (K, Mg, Ca, Na, Fe, Co, Mn, Pb, Cu, Zn and Ni) concentration. The different values obtained from AAS had shown that some rice varieties had taken up nutrients (elements) in large amount (e.g., KSK-134 had taken up maximum Fe while KSK-133 had taken up minimum) as compared to other varieties. Similarly, several varieties absorbed comparatively minimum concentration of elements. The results obtained in this research work clearly indicated that the absorption of these elements from the soil was dependent on the rice variety (e.g., BAS-2000 absorbed maximum Mg while minimum Super-BAS). Interestingly most of the rice varieties had absorbed the beneficial elements in large quantity as compared to other toxic elements.

**Key words:** Rice varieties, nutrients uptake, essential elements, minimum, comparatively, spectrophotometer

### INTRODUCTION

Rice (*Oryza sativa*) is the main source of carbohydrate as well as the source of staple nourishment of the 50% of the total world population (Ohtsubo *et al.*, 2005). Just like oats, barley and wheat, rice yields grains for nourishment and belongs to grass family Gramineae (Halliwell, 1997). In addition to small quantity of other micronutrients, rice also gives nutritionally important quantity of zinc, riboflavin, niacin and thiamin (Brown, 2007). According to Brown (2007) and Kennedy *et al.* (2002), rice has approximately 40 thousand varieties all over the world. Minor amount of lipids are present in white rice prevailed by lengthy chain double or triple bond containing fats for example, linolenic and linoleic acids chain. In correlation, rice flour is rich in these double or triple bond containing fats and represent approximately

15/100 on dry weight basis (Wilson *et al.*, 2000). These unsaturated fatty acids could be degraded when store for extended period, lead in deterioration of flavor, loss of taste and also affect the quality of eating (Zhou *et al.*, 2003). In non-glutinous brown rice, lipid substances and unsaturated fats synthesis are influenced by day to day mean temperature at the time of ripening process (Taira and Maeshige, 1979).

Rice is most significant crop and is the main source of nourishment for half of the world (Zhu and Wu, 2008). More than 80% population in Asia alone use rice as their staple food. Similarly in Asia more than 90% of the world's rice is produced and consumed (Khush, 2005).

Rice in Pakistan both fragrant and non-fragrant is cultivated on 2.62 mill ha (100 km<sup>2</sup>) approximately which covers 12% of the total area under cultivation (Hussain *et al.*, 2011). Total 6.5 million tons of rice was

**Table 1: Methodology flow chart for the first phase of experiments**

Samples collection and seeds preparation	Sprouting of seeds	Soil preparation and sowing of seeds
Seeds were collected from National Agricultural Research Centre (NARC)	The selected and disinfected twelve to fifteen seeds of seven varieties of rice were soaked with H <sub>2</sub> O	Seven different plastic pots of size 15 inches cubic volume were taken and were labeled
↓	↓	↓
Seeds were sterilized	In seven labeled different sterile disposable Petri plates (20×100 mm) of 25-30 mL callus induction medium	The soils used for sowing of these were collected from same place Barakaho Islamabad Pakistan
↓	↓	↓
71% ethanol for 65 sec	↓	The soil was dried crushed and passed through 2 mm sieve to remove the stones
↓	The labeled seven different cultures were kept in a room temperature at 30 ± 1°C for four days until sprouting was appeared	↓
Immersion in 52% (v/v) commercial bleach (5.20% NaClO <sub>2</sub> )		The sand was mixed to this soft soil in 1:3 ratios
↓		↓
On a magnetic stirrer, nonstop stirring for 30-40 sec		The sand was mixed in order to show good growth by the rice plants
↓		↓
After sterilization, the seeds were washed 3-4 times with distilled water		The seven different sprouted rice seeds were sown at rate of 30/pot in seven special seedling raising labeled plastic pots
		↓
		Tube well water was given until all the soil in pot was completely soaked
		↓
		The next water was given after every 2 days
		↓
		The sown seeds were kept at room temperature for 2 days

produced in the years 2005-2006 with the increase of 2.12 tons/ha. To meet every day metabolic requirements, individuals need twenty three element components that micro and macro nutrients are included and are obtained from foods dietary. Being the staple nourishment for more than a large portion of the world population, rice meets 20% of protein and 27% of caloric supply in developed countries (Tollens, 2007). In Asia, where 95% of the world's rice yield is used, 40-80% of the caloric ingestion originates from rice and its items. Even though the grains of rice are deficient in essential nutritional substances, 50% of these elements are found in the bran (i.e., seed coat, nucellus, seed coat, aleurone layer and pericarp) and 10% in the embryo and process of milling gives us 28% of the grain in polished rice by removing the bran and embryo. The large use of rice grain is in the form of milled (white) rice. The process of polishing (milling) removes both embryo and bran of rice. The milling process leaves only 28% of elements in the white rice (Hunt *et al.*, 2002). Internationally the malnutrition of the micronutrients produces learning disabilities in children, raises the rate of mortality and morbidity and diminishes the work efficiencies which decline the nation economic development (Welch and Graham, 2004). Almost 66% of the children death is mainly linked with deficiencies of these micronutrients (Caballero, 2003).

Mineral components most regularly deficient in human body eating regimens are Zn and Fe, at the same time, different components for example Selenium (Se), Iodine (I), Copper (Cu), Magnesium (Mg) and Calcium

(Ca) can be not enough in the diet of a few populations (Welch and Graham, 2004). Around three billion individuals of the world suffered from the effects of micronutrient insufficiencies, especially of Fe and Zn and the said proportion is as yet rising (Caballero, 2003; Mason and Garcia, 1993; Nestel *et al.*, 2006; Welch *et al.*, 1997). Billions of individuals are at danger of Zinc in-sufficiency and it is positioned as the 5th driving danger variable for diseases for example, loose motions and pneumonia in youngsters Virk. These insufficiencies are produced by continual eating habitual of the same diet (over-reliance on a single staple sustenance), circumstances of nourishment inadequacy where populaces don't have enough to eat, low eating of main sources of elements such as fruits, vegetables and products of fish and animals (FAO, 2001). The majority of those harrowed with micronutrient malnourishment are subject to staple products like rice. The approaches on food to accomplish the requirement of micronutrients to human beings have worldwide strong support (Halliwell, 1997; FAO/WHO, 1996) (Table 1).

## MATERIALS AND METHODS

The seven varieties of rice seeds which were collected are KSK-282, KSK-134, BAS-515, IR6, BAS-2000, KSK-133 and super-BAS. The chemicals that were used are Tap H<sub>2</sub>O, 71% Ethanol, Nitric acid (HNO<sub>3</sub>), Sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) and Perchloric acid (HClO<sub>4</sub>). All the chemicals were of Merck Company. The equipment's that were used are sieve (2 mm), Incubator (Shemadso), Petri plates (20×100 mm), flasks (50 and 80 mL), hot plate, filter

**Table 2: Methodology flow chart for the second phase of experiments**

Green house	Drying	Elemental analysis
Plastic pots were shifted to green house and were kept under in plastic sheets	The grown plants the roots of plants were washed by tube well H <sub>2</sub> O and were kept on new labeled and cleaned study sheets	100 ppm of stock solution of the K, Mg, Ca, Na, Fe, Co, Mn, Cu, Zn, Pb and Ni were prepared by dissolving required amount of salts in distilled water
Plastic pots were kept for 60 days at	The roots of plants were washed by tube well H <sub>2</sub> O and were kept on new labeled and cleaned paper sheets	HNO <sub>3</sub> , H <sub>2</sub> SO <sub>4</sub> and HClO <sub>4</sub> were taken in three room temperature sterilized different 50 mL flasks
After the completion of growth duration the plastic sheet was removed	The roots were washed 3-4 time through sterilized distilled water	Were mixed in ratio of 5:1:0.1, respectively
The roots were washed 3-4 time through plants for 8 days at room temperature sterilized distilled water	Plants were kept for 8 days at room temperature	From the mixed acids solution 6.5 mL solutions were kept were taken in sterilized seven different 50 mL flasks
	The roots were separated (i.e., Removed) from plants and were kept in different sterile disposable Petri plates (20×100 mm) containing 25-30 mL callus induction medium	Flasks were labeled as KSK-282, KSK-134, BAS-515, IR6, BAS-2000, KSK-133 and Super-BAS
	Petri plates were shifted to incubator for further drying	The powder form roots rice were poured in the respective labeled flasks
	The roots were incubated for 3 days at 37°C	Flask was kept on hot plate and boiled
	The separated dried roots of each variety of rice were weighed 0.5g by analytical balance separately	During boiling process colored fumes were emerged from flask
	The labeled dried roots were milled into powder with electrical grinder	Sample was boiled for 50 min and colorless fumes were emerged from flask this indicates that digestion is completed
	Stored in airtight bottles before analysis	
	Determination of mentioned macro and micro nutrients were done by Perchloric-acid digestion procedure	Few drops of distilled H <sub>2</sub> O were added and were allowed to cool at room temperature
		Make up the volume up to 50 mL by adding distilled H <sub>2</sub> O in every flask
		The solution of digested roots, three acids solution and distilled H <sub>2</sub> O were filtered through whatmann filter paper No. 42 and was collected in plastic bottles
		The concentration of macro and micro nutrients (K, Mg, Ca, Na, Fe, Co, Mn, Cu, Cr, Pb, Zn and Ni) were determined by atomic absorption Spectrophotometer

paper (Whatmann No. 42), funnel and Atomic Absorption Spectrophotometer (Shimadzu aa-6300) (Table 2).

**Erythematic formula:** The AAS values were converted by using the following equation:

$$OD = W \times \text{Dilution factor}$$

Where:

- OD = Optical density (that is given by AAS)
- W = Weight of dry plant
- Dilution factor = 50 mL
- W = The 0.5 g (weight of dried root of each rice type was 0.5 g)

## RESULTS AND DISCUSSION

Grain, vegetables and fruits provide high amount of minerals, vitamins and other natural nutrients. These substances function as antioxidant (Allen, 1974). Researchers have studied that people who eat lots of fruits and veggies have lower rates of cancer and heart disease (Miller, 2016).

After the analysis of roots of grown different varieties of immature rice by atomic absorption spectrophotometer, following different nutrients in different concentration were founded such as under (Table 3 and 4).

Rice is mostly consumed as staple food source from different soils with different composition. Investigation of heavy metals and trace elements in the rice is important

Table 3: Comparisons of Mg, Ca, Na, K, Cr and Ni Quantity up take by different varieties of immature rice plant roots

Rice varieties	Mg contents (mg/kg)	Ca contents (mg/kg)	Na contents (mg/kg)	K contents (mg/kg)	Cr contents (mg/kg)	Ni contents (mg/kg)
KSK-282	2700.20	2097.52	1326.10	457.624	1.712	7.44
KSK-134	3444.36	2179.20	1593.74	702.840	0.648	13.36
BAS-515	3364.32	989.84	1249.40	698.552	0.816	11.52
IR-6	3076.04	430.16	1108.39	942.056	5.064	18.64
BAS-2000	3720.50	946.72	1633.00	876.056	8.560	20.16
KSK-133	2233.11	1159.76	1660.40	546.152	9.496	14.80
Super-BAS	2117.93	2063.84	1808.60	554.976	12.128	10.96

Table 4: Comparisons of Mn, Fe, Zn, Cu and Pb Quantity were taken up by different varieties of immature rice plant roots

Rice varieties	Mn contents (mg/kg)	Fe contents (mg/kg)	Zn contents (mg/kg)	Cu contents (mg/kg)	Pb contents (mg/kg)
KSK-282	12.240	3024.48	79.9600	12.80	20.0
KSK-134	132.24	3632.48	29.9200	2.640	25.6
BAS-515	73.040	3014.32	41.3600	10.32	35.2
IR-6	83.600	3118.48	41.5840	10.08	17.6
BAS-2000	76.800	2972.48	55.4320	12.48	24.0
KSK-133	46.080	1586.64	50.4000	3.920	28.8
Super-BAS	104.96	1657.76	102.168	6.400	27.2

because soil might be contaminated. On December 16, 2002, UNGA introduced concept “Rice is Life” and also called ‘2004’ as the year of rice because rice is recognized as primary food source for more than half world’s population (FAO and M.H.I. FOODS, 2004).

Sixteen elements out of 109 are vital for the appropriate growth and maturity of the crop. Among these Mg and S are macronutrients, needed in larger quantity while, Fe, Cu, Zn, B, Mb, Mn and Cl are micronutrients, needed in smaller quantity for the vegetative and reproductive development of the plant (Imran and Gurmani, 2011). Seventeen genotype of the Kernel rice of three locations were studied for the absorption of soil micronutrients and also the influences of intrinsic soil variable on these elements were studied. Fe, Zn, Cu and Mg were significantly varied with respect to genotype and location; similarly, these elements were affected significantly by the environmental interactions. Furthermore, soil pH and Phosphorous concentration had significantly affected the Fe concentration while the concentration of Zn was dependent on the electrical conductivity of the soil and also on the Zn availability (Pandian *et al.*, 2011).

Six domestic types of red rice (Karad, Matali, Begmi, BhriguDhan, Sukara and Chohartu), selected from the district Chamba, Himachal Pradesh, India, analyzed for macro and micro nutrients and compared with HPR-2143, the variety of white rice. During the examination, Fe was found maximum in Matali and minimum in Chohartu, similarly, Zn contents were examined highly in Matali and minimally in HPR-2143. Likewise, BhriguDhan had the largest Mg and Mn concentration while Karad had the lowest contents of the Mn and Mn. At the same time, the concentration of Na was found to be maximum in Matali and minimum in Sukara while, K was reported to be in the highest concentration in Sukara and in the lowest

concentration in Begmi (Sharma *et al.*, 2012). Liao *et al.* (2013) compared the bioavailability and bioaccumulation of Cu, Pb, Cd and Cr in the grown in the Non-Flooded controlled irrigated (NFI) and Flooding Irrigated (FI) soils and concluded that NFI took higher metal concentration of Cd, Cu and Pb than FI while the concentration of Cr in the roots of NFI rice was lower than FI. They further stated that Cr bound to Fe and Mn oxides were more stable under NFI conditions, played important role in the bioavailability of Cr in paddy soil while Cu, Cd and Pb, bound to organic matters were more likely to be released and improved the bioaccumulation of the these metals in NFI’s rice.

Three nutrients sources, i.e., organic, integrated use of organic and inorganic and chemical fertilizer were used for the growth of medicinal rice, Njavara in karamana, studied their nutrients concentration by analyzing the dry root matter at weekly intervals. The concentrations of N, P, K, Fe, Mn and Zn in the rice were higher significantly by using conventional management practices than other established techniques. By the use of integrated nutritional sources, the concentrations of N, P, K, Fe, Mn and Zn were higher than organic and inorganic sources (Rani and Sukumari, 2013).

Emumejaye (2014) studied the concentration of trace elements and heavy metals including Cu, Zn, Fe, Ni, Be, Cd, Cr, Hg and Pb in the samples of rice purchased from the market of Delta State, Nigeria. He concluded that level of trace elements were below the permissible limit while Pb and Fe were detected to have high concentration in some samples.

## CONCLUSION

It is concluded from this research that the rice KSK-282 had taken up Ni in minimum concentration as compared to other selected rice varieties. KSK-134 had

absorbed Ca, Fe and Mn in maximum amount while Cr in minimum amount as compared to other rice types. BAS-515 had taken up the maximum quantity of Pd in contrast with other 6 varieties. While K nutrient was taken up comparatively maximum by IR-6 rice. Similarly IR-6 had absorbed Ca and Na in lowest concentration as compared to other rice plants. BAS-2000 had taken up Ni in highest amount from soil under similar conditions as compared to other varieties of rice. KSK-133 had up taking maximum quantity of iron (Fe) when evaluated this rice type to other varieties. Super-BAS had shown the uptake of Na, Cr and Zn in larger amount while Mg in smaller amount in contrast to other rice types.

### REFERENCES

- Allen, S.E., 1974. Chemical Analysis of Ecological Materials. Blackwell Scientific Publications, London, UK., Pages: 565.
- Brown, A.C., 2007. Understanding Food: Principles and Preparation. 3rd Edn., Thomson Learning Incorporation, USA., ISBN-13: 9780495107453, Pages: 696.
- Caballero, B., 2003. Fortification, supplementation and nutrient balance. *Eur. J. Clin. Nutr.*, 57: S76-S78.
- Emumejaye, K., 2014. Heavy and trace elements in some brands of rice consumed in Delta State, Nigeria. *IOSR. J. Appl. Phys.*, 6: 1-5.
- FAO, J. and M.H.I. FOODS, 2004. Food and agriculture organization of the United Nations. United Nations, Rome, Italy.
- FAO/WHO, 2001. Human vitamin and mineral requirements. Report of a Joint FAO/WHO Expert Consultation, Bangkok, Thailand.
- Halliwell, B., 1997. Antioxidants and human disease: A general introduction. *Nutr. Rev.*, 55: S44-S52.
- Hunt, J.R., L.K. Johnson and B.O. Juliano, 2002. Bioavailability of Zinc from cooked Philippine milled, undermilled and brown rice, as assessed in rats by using growth, bone Zinc and Zinc-65 retention. *J. Agric. Food Chem.*, 50: 5229-5235.
- Hussain, J., N. Ur-Rehman, A.L. Khan, H. Hussain and A. Al-Harrasi, *et al.*, 2011. Determination of macro and micronutrients and nutritional prospects of six vegetable species of Mardan, Pakistan. *Pak. J. Bot.*, 43: 2829-2833.
- Imran, M. and Z.A. Gurmani, 2011. Role of macro and micro nutrients in the plant growth and development. National Agricultural Research Centre, Islamabad, Pakistan.
- Kennedy, G., B. Burlingame and V.N. Nguyen, 2002. Nutritional Contribution of Rice: Impact of Biotechnology and Biodiversity in Rice Consuming Countries. International Rice Commission, Bangkok, Thailand.
- Khush, G.S., 2005. What it will take to feed 5.0 billion rice consumers in 2030. *Plant Mol. Biol.*, 59: 1-6.
- Liao, L., J. Xu, S. Peng, Z. Qiao and X. Gao, 2013. Uptake and bioaccumulation of heavy metals in rice plants as affect by water saving irrigation. *Adv. J. Food Sci. Tech.*, 5: 1244-1248.
- Mason, J.B. and M. Garcia, 1993. Micronutrient deficiency-the global situation. *SCN News*, 9: 11-16.
- Miller B.D., 2016. Cancer: We can Win the War against Cancer by Aggressively Pursuing Prevention. Oak Publication Sdn Bhd, Malaysia.
- Nestel, P., H.E. Bouic, J.V. Meenakshi and W. Pfeiffer, 2006. Biofortification of staple food crops. *J. Nutr.*, 136: 1064-1067.
- Ohtsubo, K., K. Suzuki, Y. Yasui and T. Kasumi, 2005. Bio-functional components in the processed pre-germinated brown rice by a twin-screw extruder. *J. Food Compos. Anal.*, 18: 303-316.
- Pandian, S.S., S. Robin, K.K. Vinod, S. Rajeswari and S. Manonmani *et al.*, 2011. Influence of intrinsic soil factors on genotype-by-environment interactions governing micronutrient content of milled rice grains. *Aust. J. Crop Sci.*, 5: 1737-1742.
- Rani, S. and P. Sukumari, 2013. Root growth, nutrient uptake and yield of medicinal rice njavara under different establishment techniques and nutrient sources. *Am. J. Plant Sci.*, 1: 1568-1570.
- Sharma, J.K., T.R. Sharma and S.K. Sharma, 2012. Comparative study on macro and micro minerals composition of selective red rice landraces from Chamba district of Himachal Pradesh-India. *J. Agric. Sci.*, 8: 378-380.
- Taira, H. and M. Maeshige, 1979. Influence of variety and crop year on lipid content and fatty acid composition of lowland non-glutinous brown rice. *Jpn. J. Crop Sci.*, 48: 220-228.
- Tollens, E., 2007. Markets and institutions for promoting rice for food security and poverty reduction in Sub-Sahara Africa. *Afr. Crop Sci. J.*, Vol. 15,

- WHO/FAO, 1996. Preparation and use of Food-Based Dietary Guidelines. Nutrition Program, World Health Organization, Geneva, Switzerland..
- Welch, R.M. and R.D. Graham, 2004. Breeding for micronutrients in staple food crops from a human nutrition perspective. *J. Exp. Bot.*, 55: 353-364.
- Welch, R.M., G.F. Combs Jr. and J.M. Duxbury, 1997. Toward a greener revolution. *Sci. Technol.*, 14: 50-58.
- Wilson, T.A., L.M. Ausman, C.W. Lawton, D.M. Hegsted and R.J. Nicolosi, 2000. Comparative cholesterol lowering properties of vegetable oils: Beyond fatty acids. *J. Am. Coll. Nutr.*, 19: 601-607.
- Zhou, Z., C. Blanchard, S. Helliwell and K. Robards, 2003. Fatty acid composition of three rice varieties following storage. *J. Cereal Sci.*, 37: 327-335.
- Zhu, Z. and R. Wu, 2008. Regeneration of transgenic rice plants using high salt for selection without the need for antibiotics or herbicides. *Plant Sci.*, 174: 519-523.