

Synthesis, Characterization and Structural Studies of Co(II), Ni(II), Cu(II) and Cd(II) Complexes of Hydrazone Schiff Base Ligands Pyrimidine Derivative

Taghreed M. Musa, Anaam M. Rasheed, Sinan M. AL-Bayati,
Wessal M. Khamis and Bayader F. Abbas

Department of Chemistry, College of Science, AL-Mustansiriya University, Baghdad, Iraq

Abstract: The tridentate Schiff base ligands containing donor ONO were prepared by the reaction of 2-Furaldehyde and urea or thiourea with ethyl acetoacetate to product pyrimidine derivatives (1a, 1b) which upon treatment with hydrazine afforded the hydrazone derivatives (2a, 2b) the reaction of (2a, ab) with 3-acetyl coumarin obtained novel Schiff base Ligands (L_1 , L_2) in good yield. The Ligands (L_1 , L_2) were characterized by elemental analysis (C.H.N.S), FT-IR, Uv spectroscopy, HNMR and mass spectral analysis. Complexes of Co(II), Ni(II), Cu(II) and Cd(II) with the novel Schiff base ligands have been prepared. The prepared complexes were identified and their structural geometries were suggested by using elemental analysis (C.H.N.S), FT-IR, UV-Vis spectroscopy and flame atomic absorption technique in addition magnetic susceptibility, conductivity measurements and thermal analysis TG-DSC. The results suggest that the Schiff base Ligands (L_1 , L_2) coordinate with their tridentate O, N, O donors derived from carbonylamide and azomethine nitrogen and carbonyl oxygen atom on the coumarin ring.

Key words: Pyrimidine derivative, coumarine derivative, Schiff base, transition metal complexes, spectral studies, susceptibility

INTRODUCTION

Pyrimidine derivatives represent very important heterocyclic compounds for their antimicrobial activity (Gulcan *et al.*, 2012) and photoluminescence properties (Onal *et al.*, 2011). The coordination chemistry of hydrazones system has received an increasing interest because of its play an important role in inorganic chemistry (Pouralimardan *et al.*, 2007; Fouda *et al.*, 2008) as they easily form stable complexes with most transition metal ions (Yaul *et al.*, 2009; Kriza *et al.*, 2010). Many derivatives of this system showed biological significance (Dhande *et al.*, 2007). The derivatives of coumarine nucleus have occupied a unique place in field of medicinal chemistry due to wide range of biological activities like

antimicrobial (Lamani *et al.*, 2009; Farshori *et al.*, 2011), anticancer (Bhattacharyya *et al.*, 2009), antibacterial activity and antifungal activity of coumarin (Darla *et al.*, 2015). Schiff bases and their transition metal complexes have also been widely reported to be biologically versatile compounds having antimicrobial (Kulkarni *et al.*, 2009) and antifungal (Mishra and Soni, 2008; Srivastava *et al.*, 2014). The present paper describes the preparation of new Schiff's base derivatives produced from the reaction of hydrazone derivatives and 3-acetyl coumarin to be used Ligands (L_1 , L_2) that provide three potential donor sites to form complexes with some transition metal ions. The Ligands (L_1 , L_2) and its complexes have been fully characterized and their structures were determined as outlined (Fig. 1).




Fig. 1: Referred to preparation equation of L_1 , L_2

MATERIALS AND METHODS

All the chemicals used were analytical annular and of highest purity available. Melting points were measured on a Gallan Kamp MFB-600 melting points apparatus. Fourier Transform Infrared (FT-IR) spectra were recorded on a Shimadzu spectrophotometer. Electronic spectra using varian UV-visible spectrophotometer. Elemental analysis for ligand and metal complexes were carried out using CHNS elemental analyzer and atomic absorption using Shimadzu Model 6809. Mass spectra were recorded by Electron Impact Mass Spectrometry (EIMS) using a Direct injection probe on a Shimadzu GCMS-QP5050 A spectrometer. Molar conductivity of the complexes were measured in DMF as solvent in 0.001M solution using a CON 510 bench conductivity meter with 2-ring stainless steel conductivity electrode (Cell constant, K = 1.0) ¹HNMR spectra were recorded on Burker DMX-5000 NMR (300-600 MHz) spectrophotometer with using CDCl₃ as a solvent in Jordan University. TG and DSC (differential scanning Colorimetry) thermos grams indifferent ranges were carried out at (RT) heating rate = 10°C/min. (Linseis STA university Abn Al-Hitham College.

Synthesis: The two ligands were prepared in a similar manner and here the synthesis of (L₁, L₂) are given:

Synthesis of pyrimidine derivative (1a), (1b): Ethyl-4-(furan-2-yl)-6-methyl-2-oxo-1, 2, 3, 4-tetrahydropyrimidine-5-carboxylate (1a). Ethyl-4-(furan-2-yl)-6-methyl-2-thioxo-1, 2, 3, 4-tetrahydropyrimidine-5-carboxylate (1b).

Ethyl acetoacetate (1.30 g, 0.01 mole) was added to the mixture of 2-furaldehyde (0.96 g, 0.01 mole) and urea (1.80 g, 0.03 mole) or thiourea (2.28 g, 0.01 mole), respectively was dissolved in absolute ethanol (30 mL). To this solution added (0.9 g) from Lanthanum Chloride heptahydrate and the mixture was refluxed for 7 h then

cooled at room temperature. The reaction mixture was poured to crushed ice (30 g) with stirring for 10 min. The precipitates (1a, 1b) were filtered and washed with cold water then recrystallized from ethanol.

Synthesis of hydrazone derivative (2a), (2b): 4-(furan-2-yl)-6-methyl-2-oxo-1, 2, 3, 4-tetrahydropyrimidine-5-Carbohydrazide. (2a). 4-(furan-2-yl)-6-methyl-2-thioxo-1, 2, 3, 4-tetrahydropyrimidine-5-carbohydrazide (2b).

To a solution of compound (1a) (2.5 g, 0.01 mole) or compound (1b) (2.66 g, 0.01 mole), respectively in absolute ethanol (20 mL), hydrazone hydrate (2.5 g, 0.05 mole) was added. The mixture was refluxed for 12 h. The solvent was removed and the solid products (2a, 2b) were collected and crystallized from ethanol.

Synthesis of Coumarin derivative (C): A mixture of saileyldehyde (1.06 g, 0.01 mole) and ethyl aceto acetate (1.3 g, 0.01 mole) in absolute ethanol (20 mL) with drops of tripropyl amine was added. The mixture was refluxed for 5 h. After cooling, the product was filtrated and recrystallized from ethanol.

Synthesis of hydrazone Schiff base Ligands (L₁, L₂): 4-(furan-2-yl)-6-methyl-2-oxo-N-[1-(2-oxo-2H-Chromen-3-yl) ethylidene] -1, 2, 3, 4-tetrahydropyrimidine-5-carbohydrazide (L₁). 4-(furan-2-yl)-6-methyl-N-[1-(2-oxo-2H-chromen-3-ethylidene)]-2-thioxo-1, 2, 3, 4-tetrahydropyrimidine-5-carbohydrazide (L₂). A mixture of compound (2a) (2.36 g, 0.01 mole) or compound (2b) (2.52 g, 0.01 mole), respectively and (1.88 g, 0.0 mole) of acetyl coumarin in absolute ethanol (25 mL) with three drops of glacial acetic acid was added. The mixture was refluxed for 4 h and cooled at room temperature. The solid products were collected by filtration and then recrystallized from ethanol. The physical properties of the prepared ligands are given in Table 1.

Table 1: Physical data for (1a, 1b, 2a, 2b, L₁, L₂) and their complexes

Comp. No.	Colors	M.P.°C	M.wt g/mol	Elementa analysis found (Calc.) (%)						M:L (EtOH)	Suggest formula
				C%	H%	N%	S%	M%			
1a	Pale yellow	147	250	---	---	---	---	---	---	C ₁₂ H ₁₄ N ₂ O ₄	
2a	Light yellow	196	236	---	---	---	---	---	---	C ₁₀ H ₁₂ N ₄ O ₃	
L ₁	Yellow	109	406	62.32 (62.06)	4.65 (13.79)	13.47	---	---	---	C ₂₁ H ₁₈ N ₄ O ₅ S	
1b	Light brown	160	266	---	---	---	---	---	---	C ₁₂ H ₁₄ N ₂ O ₃ S	
2b	Light grey	202	252	---	---	---	---	---	---	C ₁₀ H ₁₂ N ₄ O ₂ S	
L ₂	Dark green	212	422	59.93 (59.71)	4.09 (4.26)	13.52 (13.27)	7.75 (7.58)	---	---	C ₂₁ H ₁₈ N ₄ O ₄ S(L ₂)	
C	Yellowish white	119	188	---	---	---	---	---	---	C ₁₁ H ₈ O ₃	
CoL ₁	Brown	175	535.9	47.27 (47.02)	3.60 (3.35)	10.21 (10.44)	---	10.65 (10.99)	1:1	[Co(L ₁)Cl]Cl	
NiL ₁	Light brown	155	941.7	53.401	3.95	11.99	---	6.49	1:2	[Ni(L ₁) ₂]Cl ₂	
CuL ₁	Green	190	946.5	53.61 (53.24)	3.63 (3.80)	11.55 (11.83)	---	6.47 (6.70)	1:2	[Cu(L ₁) ₂]Cl ₂	
CdL ₁	Light yellow	180	589.4	42.51	3.30	9.40	---	19.32	1:1	[Cd(L ₁)Cl]Cl	
CoL ₂	Dark brown	245d'	551.9	45.48 (45.66)	3.55 (3.26)	10.36 (10.14)	5.42 (5.79)	10.39	1:1	[Co(L ₂)Cl]Cl	
CuL ₂	Brown	235d.	978.5	51.39 (51.50)	3.552 (3.67)	11.41 (11.44)	6.23 (6.48)	6.29	1:2	[Cu(L ₂) ₂]Cl ₂	

d* = decompose

Synthesis of metal complexes: A hot ethanolic solution of each of metal ion salts (1 m mole) [(0.238 g, of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$), (0.237 of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$), (0.17g of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$) and (0.219 g of $\text{CdCl}_2 \cdot 2\text{H}_2\text{O}$)], respectively was added to a hot ethanolic solution of Ligand (L_1) (0.406 g, 1 m mole) or of Ligand (L_2) (0.422 g, 1 m mole), respectively with stirring. The mixture was heated under reflux 2 h during this time the color solid was formed. The products were collected by filtration and then recrystallized from absolute ethanol. The physical data of the prepared complexes are shown in Table 1.

Study of complexes formation in solution: Complexes of Schiff base Ligands (L_1 , L_2) with metal ions have been studied in solution using ethanol as a solvent in order to evaluate the ratio of metal to ligand in complex using molar ratio method (Douglas *et al.*, 2011). A series of solution have been prepared having a constant concentration 10^{-3} M for each metal ion and ligand. The ratio have been determined from the relationship between the absorption of the absorbed light and molar ratio of $[\text{M}:\text{L}]$ at λ_{max} of maximum absorption.

RESULTS AND DISCUSSION

Reaction between 2-Furaldehyde, (Urea or thiourea) and ethylacetoacetate in absolute ethanol with lanthanum chloride heptahydrate afforded the pyrimidine derivative (1a, 1b) which were treated with hydrazine afforded the hydrazide derivative (2a, 2b). These compounds were treated with coumarin derivative to obtain Ligands (L_1 , L_2) in good yield. Compounds were formed through the following suggest mechanism (Fig. 2):

Elemental analysis: The analytical data of Ligands (L_1 , L_2) and it's metal complexes are given in Table 1 and are in a satisfactory agreement with the calculated values. The suggested molecular is formulated and characterized by subsequent spectral and molar ratio as well as magnetic moment.

$^1\text{HNMR}$ spectra: The $^1\text{HNMR}$ spectrum of the Ligand (L_1) in (300 MHz, CDCl_3) exhibited (Batterham, 1973) peaks at 1.2 ppm (1H, CH) and 2.3-2.5 ppm (2H, CH_2) due to CH_3 pyrimidine ring, 3.5 ppm (3H, methyl) (Yaul *et al.*, 2009),




Fig. 2: Referred to preparation steps of L_1 , L_2

4-4.2 ppm (2H, furanyl), 5.3 ppm (1H, furanyl) (Chohan and Kausar, 2000), 9.3 ppm (1H, iminol) (Yaul *et al.*, 2009), 6.3-7.9 ppm (6H, Ar-H), 8.6 ppm (2H, NH pyrimidine ring) (Yaul *et al.*, 2009; Silverstein *et al.*, 1981) (Fig. 1).

Mass spectra: The formation of compounds (1a,1b) and Ligand (L_2) were studied through their GC-Mass spectrometer. The proposed molecular formula of compounds was confirmed by comparing their molecular formula weight with the m/z values. In the mass spectra of the compounds, peaks were attributed to the molecular ions, m/z:249 for the compound (1a) $C_{12}H_{14}N_2O_4$; 266 for the compound (1b) $C_{12}H_{14}N_2O_3S$; 252 for the compound (2b) $C_{10}H_{12}N_4O_2S$; 188 for the compound (C) $C_{11}H_5O_3$ and m/z: 422 for the Ligand (L_2) which data excellent agreement with theoretical values (Srivastava *et al.*, 2014; Parikh, 1974; Porter, 1985). These results showed in Fig. 2-5.




Fig. 3: Referred to preparation steps of L_1, L_2




Fig. 4: 1H NMR spectra of free Ligand (L_1)




Fig. 5: GC-Mass spectrum of compound (1a)

Table 2: Characteristic infrared absorption frequencies in (cm⁻¹) of Ligands (L₁, L₂) and their metal complexes

Comp.No.	$\nu\text{C} = \text{O}_{\text{amide}}$	$\nu\text{C} = \text{N}_{\text{azomethine}}$	$\nu\text{C} = \text{O}_{\text{coumarin ring}}$	M-N	M-O	M-Cl
L ₁	1645	1610	1699	----	----	----
CoL ₁	1629	1589	1660	570	460	395
NiL ₁	1625	1540	1678	560	455	----
CuL ₁	1620	1589	1687	565	450	----
CdL ₁	1637	1558	1689	555	435	385
L ₂	1647	1600	1726	----	----	----
CoL ₂	1639	1585	1687	550	445	390
CuL ₂	1641	1527	1697	565	430	----

Table 3: Electronic spectra, conductance in (DMF) and Magnetic moment (B.M) for (L₁, L₂) and their metal complexes

Comp. NO.	Bands (cm ⁻¹)	Assignment	Molar Cond. (μs.cm ⁻¹)	μB.M	Suggested structure
L ₁	38461 28490	$\pi \rightarrow \pi^*$ $n \rightarrow \pi^*$	----	----	---
CoL ₁	15384 16949 25974	$^4A_2 \rightarrow ^4T_{1(F)}$ $^4A_2 \rightarrow ^4T_{1(P)}$ C.T	71	4.49	T.h
NiL ₁	12820 18348 29069	$^3A_2 \rightarrow ^3T_2g(F)$ $^3A_2g \rightarrow ^3T_2g(F)$ $^3A_2g \rightarrow ^3T_1g(P)$	135	2.92	O.h
CuL ₁	15267	$^2Eg \rightarrow ^2T_2g$	168	1.79	O.h
CdL ₁	47619 32258 12987	$\pi \rightarrow \pi^*$ $n \rightarrow \pi^*$ L-M(C.T)	78	0.00	T.h
L ₂	35087 29154	$\pi \rightarrow \pi^*$ $n \rightarrow \pi^*$	----	----	---
CoL ₂	15037 17921 25641	$^4A_2 \rightarrow ^4T_1(F)$ $^4A_2 \rightarrow ^4T_1(P)$ C.T	66	4.52	T.h
CuL ₂	14925	$^2Eg \rightarrow ^2T_2g$	159	1.83	O.h

Infrared spectroscopic study: The main stretching frequencies of the IR spectra of the Ligands (L₁, L₂) and their metal complexes are given in Table 2. The structure of compounds (1a, 1b) were confirmed by the presence of $\nu\text{C} = \text{O}$ ester stretching vibration at (1734, 1716) cm⁻¹ in addition to the bands at (1215, 1203), (2920, 2990) and (3170-3275) cm⁻¹ due to $\nu\text{C} = \text{O}$ ester, $\nu\text{C-H}$ and $\nu\text{N-H}$ for pyrimidine ring (Nakamoto, 2009). The reaction of compounds (1a, 1b) with hydrazine hydrate afforded the compounds (2a, 2b), the appearance of IR new absorption bands at (3215-3344) cm⁻¹ due to νNH_2 (Nakamoto, 2009). The structure of compound 3-acetyl Coumarin (C) was confirmed by the appearance of $\nu(\text{C} = \text{O}, \text{C-O})$ coumarin ring stretching frequency at (1735, 1047) cm⁻¹ and $\nu\text{C} = \text{O}$ ester coumarin stretching frequency at (1674) cm⁻¹ (Sonmez and Sekerci, 2002). Moreover, treatment of NH₂ in hydrazide derivatives (2a, 2b) with 3-acetyl coumarin to give Schiff base Ligands (L₁, L₂). These reaction were followed by disappearance absorption bands due to $\nu(\text{NH}_2)$ and characteristic bands at (1610,1600) cm⁻¹ for azomethine group $\nu(\text{C} = \text{N})$ are utilized confirm the structures of (L₁, L₂) (Abon-Melha and Faruk, 2008). The all metal complexes of the Ligands (L₁, L₂) behave as tridentate coordinating to the metal through the oxygen of carbonyl (amide and coumarin ring) and nitrogen of the azomethine groups, therefore, the bands due to

$\nu\text{C} = \text{O}$ and $\nu\text{C} = \text{N})$ were shifted to lower frequencies (Tharmaraj *et al.*, 2009), Table 2. These observation were further indicated by the appearance of ($\nu\text{M-O}$, $\nu\text{M-N}$ and $\nu\text{M-Cl}$), respectively (Tharmaraj *et al.*, 2009) (Table 2).

Electronic spectra, magnetic moment and conductance studies: The Uv spectrum of the Ligand (L₁) mostly showed two intense maxima bands at 260 nm and 351 nm which belong to $\pi \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$, respectively (Silverstein *et al.*, 1981), (Fig. 6) while the Uv spectrum of the Ligand (L₂) showed intense bands at 285 and 343 nm which belong to $\pi \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$, respectively (Silverstein *et al.*, 1981) (Fig. 7 and Table 3). The solution spectra of the Cobalt (II) complex for the Ligand (L₁) showed three band at (15384,16949) cm⁻¹ due to $^4A_2 \rightarrow ^4T_{1(F)}$, $^4A_2 \rightarrow ^4T_{1(P)}$ with charge transfer band at 25974 cm⁻¹ (Fig. 8), these bands correspond to those of tetrahedral geometry (Mohamed *et al.*, 2006), magnetic moment of the solid complex (4.49B.M) showed a high spin Co(II) complex (Nicholls, 1973). Conductivity measurement in DMF showed that the complex was conductive behavior (Table 3). The solution spectra of the Ni(II) complex exhibited a well-discernable band with a shoulder on the low energy side the bands generally observed in the region at (12820, 18348, 29069) cm⁻¹ are assigned to $^3A_2g(F) \rightarrow ^3T_2g(F)$, $^3A_2g(F) \rightarrow ^3T_1g(F)$ and $^3A_2g(F) \rightarrow ^3T_1g(P)$ transition, respectively (Singh *et al.*, 2010) (Fig. 9). The magnetic




Fig. 6: GC-Mass spectrum of compound (1b)




Fig. 7: GC-Mass spectra of compound (2b)




Fig. 8: GC-Mass spectrum of compound (C)

moment of the Nickel complex at room temperature was (2.92 B.M) showing an octahedral environment around the Ni(II) ion in the complex (Kriza *et al.*, 2010), table. Conductivity measurement in DMF showed an electrolyte behavior of the complex. The electronic absorption spectrum of Cu(II) complex for the Ligand (L_2) exhibits a

band at 15267 cm^{-1} that was assigned to the ${}^2E_g \rightarrow {}^2T_{2g}$ transition, suggesting octahedral geometry around the Cu(II) ion. The magnetic moment value of this complex was (1.79 B.M) which indicates the presence of one unpaired electron per Cu(II) ion (Lever, 1984). The conductivity measurement showed that the complex was




Fig. 9: UV-visible spectrum of (L_1)




Fig. 10: UV-visible spectrum of Ligand (L_2)

to be ionic, Table 3, Cd (II) complex show no absorption peak at range (380-1000) nm that indicates no d-d electronic transition happened (d^{10} -system) invisible region that is a good result for Cd (II) tetrahedral complex (Abid *et al.*, 2016). Conductivity measurement in DMF showed that the complex was to be ionic (Table 3). The solution spectrum of the prepared Co(II) complex for the Ligand (L_2) showed three bands at (15037, 17921) cm^{-1} due to ${}^4A_2 \rightarrow {}^4T_{1(F)}$, ${}^4A_2 \rightarrow {}^4T_{1(F)}$ with charge transfer band at (25641) cm^{-1} (Fig. 10) these bands correspond to those of tetrahedral geometry (Al-Hasani *et al.*, 2011). The value of the magnetic measurement (4.52 B.M) showed a high spin Co(II) complex (Earnshaw, 1968). Conductivity measurement in DMF showed that complex was due to be ionic (Table 3). The electronic spectrum of Cu(II) complex for Ligand (L_2) gave broad band at (14925) cm^{-1} which assigned to ${}^2E_g \rightarrow {}^2T_{2g}$ transition of the octahedral geometry (Al-Jibori *et al.*, 2013). The magnetic moment value was (1.83 BM) corresponding to one unpaired




Fig. 11: UV-visible spectrum of NiL_1 complex




Fig. 12: UV-visible spectrum of CoL_2 complex

electron (Rasheed *et al.*, 2014). Conductivity measurement in DMF showed a conductive behavior of this complex (Table 3).

Thermal degradation of ligand and complexes: The TG data suggest a mechanism for degradation of the complexes of $[Cu(L_1)_2]Cl_2[Co(L_2)Cl]Cl$, to 351°C . while the $[L_2]$ and Complex of $[Cu(L_2)_2]Cl_2$ to 600°C . The thermal degradation of $[L_2]$ display in Fig. 11. The thermo gram curve show three decomposition steps. The TGA peak observed at 259.59°C assigned to the loss of (C-O) furfural group fragment, (det.1.43 mg, 6.63%; calcu. = 1.40 mg). The other steps occurred at 437.5°C revealed the loss of (COO) of coumarin ring, (obs.2.31 mg, 11.20%; calcu. = 2.34 mg). The third step of decomposition of compound 592.9 is related to the loss (CH = CHO) furfural ring, (obs.2.46 mg, 11.74%; calcu. = 2.30 mg). The differences in the weight may be related to a sublimation process occurred at high temperature. The DSC curve recorded peaks at 84.3, 269.6 which refer to an exo thermic




Fig. 13: DSC thermogram of an argon atmosphere [L₂]




Fig. 14: TG thermogram of an argon atmosphere [Co(L₂)]Cl₂

decomposition process (Abid *et al.*, 2016) (Fig. 12). The thermal analysis clearly indicated that the decomposition of the complexes of [Co(L₂)Cl] is shown in Fig. 13-16 which indicated the complex is stable up in 134°C. The TGA peak observed at 134°C assigned to the loss of Cl coordination outer sphere (counter ions), (det = 0.51 mg, 5.56%; calcu. = 0.5749 mg. Figure 13 and 14 TG thermo

gram of an argon atmosphere of [Cu(L₂)₂]Cl₂, [Cu(L₁)₂]Cl₂, respectively (Al-Bayati *et al.*, 2018). The molecular model using Hyperchem. Program for L₁, Cd(L₁) complex, L₂ and CoL₂ complex were depicted, respectively in Fig. 18 and 19. According to these results the structure configuration of ligands and these metal complexes may be proposed in Fig. 16-21.




Fig. 15: TG thermogram of an argon atmosphere $[Cu(L_2)_2]Cl_2$




Fig. 16: TG thermogram of an argon atmosphere $[L_2]$




Fig. 17: DSC thermogram of $[Cu(L_1)_2]Cl_2$ in an argon atmosphere




Fig. 18: a, b) The proposed molecular structure of Ligand $[L_1]$




Fig. 19: a, b) The proposed molecular structure of Ligand $[L_2]$




Fig. 20: a, b) The proposed molecular structure of [Co(L₂)Cl]Cl₁




Fig. 21: a, b) The proposed molecular structure of [Cd(L₁)Cl]Cl₁

CONCLUSION

We have prepared two Schiff base Ligands (L₁, L₂) the ligands behave as tridentate ligand through (O, N, O) atoms. According to the results obtained from GC-Mas spectra and elemental analysis (C.H.N.) gave formula C₂₁H₁₈N₄O₅ (L₁) and C₂₁H₁₈N₄O₄S(L₂). Spectroscopic (FT-IR, HNMR) data showed the structure of the ligand contain azomethine and carbonyl groups. From conductivity measurements showed that all prepared complexes are ionic as well as magnetic moment, Uv-visible study and the molecular model using Hyper Chem for [CoL₁, CdL₁ and CoL₂] complexes have tetrahedral geometry and [NiL₁, CuL₁ and CuL₂] complexes have octahedral geometry.

ACKNOWLEDGMENT

Thanks should be present to our university (Mustansiriyah University) for all helps that submitted to accomplishment this research. The grate should be submitted to the dean of science college and head of chemistry department. Also the researchers are grateful to all colleagues whose help to achieve this research.

REFERENCES

- Abid, K., S. Al-Bayati and A. Rasheed, 2016. Synthesis, characterization, thermal study and biological evaluation of transition metal complexes supported by ONNNO-Pentadentate Schiff base ligand. *Am. J. Chem.*, 6: 1-7.
- Abon-Melha, K.S. and H. Faruk, 2008. Bimetallic complexes of Schiff base-[4-hydroxy coumarin-3-yl]-N, 5N-thiocarbohydrazone as potentially Dibasic pentdentate ligand Synthesis spectra and Antimicrobial properties. *J. Iran. Chem. Soc.*, 5: 122-124.
- Al-Bayati, S., A. Rasheed, E. Zuhair, K. Abid and W. Khamis, 2018. Synthesis and spectroscopic studies of new alkoxy Schiff base complexes based on coumarin moiety. *Am. J. Chem.*, 8: 1-7.
- Al-Hasani, R.A., S.M. Al-Byatti and S.M. Al-Azawi, 2011. Synthesis, structural and biological studies of /3-(1, 3-benzothiazol-2-yl)- /4H-spiro [indole2, 3-[1, 3] thiazolidine]- 2,/4 (1H) dion with Cr (III), Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) ions. *Eng. Technol. J.*, 29: 3067-3078.

- Al-Jibori, M.N., S.M. Al-Bayati and A.M. Rasheed, 2013. Coordination modes of a new ligand derived from pyrazoline with Cr(III), Mn(II), Co(II), Ni(II), Cu(II) and Zn(II) metal ions: Synthesis, identification and biological studies. *Chem. Mater. Res.*, 3: 1-10.
- Batterham, T.J., 1973. *NMR Spectra of Simple Heterocycles*. 99th Edn., Wiley, New York, USA., ISBN:9780471057024, Pages: 540.]
- Bhattacharyya, S.S., S. Paul, S.K. Mandal, A. Banerjee and N. Boujedaini *et al.*, 2009. A synthetic coumarin (4-methyl-7 hydroxy coumarin) has anti-cancer potentials against DMBA-induced skin cancer in mice. *Eur. J. Pharm.*, 614: 128-136.
- Chohan, Z.H. and S. Kausar, 2000. Synthesis, characterization and biological properties of tridentate NNO, NNS and NNN donor thiazole-derived furanyl, thiophenyl and pyrrolyl schiff bases and their Co(II), Cu(II), Ni(II) and Zn(II) metal chelates. *Met. Based Drugs*, 7: 17-22.
- Darla, M.M., B.S. Krishna, K.U. Rao, N.B. Reddy and M.K. Srivash *et al.*, 2015. Synthesis and bio-evaluation of novel 7-hydroxy coumarin derivatives via Knoevenagel reaction. *Res. Chem. Int.*, 41: 1115-1133.
- Dhande, V.V., V.B. Badwaik and A.S. Aswar, 2007. Hydrazone as complexing agent: Synthesis, structural characterization and biological studies of some complexes. *Russ. J. Inorg. Chem.*, 52: 1206-1210.
- Douglas, A.S., M.N. Donald, F.J. Holler, S.R. Crouch and S.C. Chen, 2011. *Introduction to Analytical Chemistry*. 1st Edn., Case Bound-Think, Inc., USA.,
- Earnshaw, A., 1968. *Introduction to Magnetochemistry*. Academic Press, Cambridge, Massachusetts, USA., Pages: 115.
- Farshori, N.N., M.R. Banday, A. Ahmad, A.U. Khan and A. Rauf, 2011. 7-Hydroxy-coumarin derivatives: Synthesis, characterization and preliminary antimicrobial activities. *Med. Chem. Res.*, 20: 535-541.
- Fouda, A.S., G.E. Badr and M.N. El-Haddad, 2008. The inhibition of C-steel corrosion in H₃PO₄ solution by some furfural hydrazone derivatives. *J. Korean Chem. Soc.*, 52: 124-132.
- Gulcan, M., M. Sonmez and I. Berber, 2012. Synthesis, characterization and antimicrobial activity of a new pyrimidine schiff base and its Cu(II), Ni(II), Co(II), Pt(II) and Pd(II) complexes. *Turk. J. Chem.*, 36: 189-200.
- Kriza, A., V.L. Ababei, N. Cioatera, I. Rau and N. Stanica, 2010. Synthesis and structural studies of complexes of Cu, Co, Ni and Zn with isonicotinic acid hydrazide and isonicotinic acid (1-naphthylmethylene) hydrazide. *J. Serb. Chem. Soc.*, 75: 229-242.
- Kulkarni, A.D., S.A. Patil and P.S. Badami, 2009. Electrochemical properties of some transition metal complexes: Synthesis, characterization and in-vitro antimicrobial studies of Co(II), Ni(II), Cu(II), Mn(II) and Fe(III) complexes. *Intl. J. Electrochem. Sci.*, 4: 717-729.
- Lamani, S.S., O. Kotresh, A. Phaniband, M. Shafi and C. Kadakol *et al.*, 2009. Synthesis, characterization and antimicrobial properties of Schiff bases derived from condensation of 8-formyl-7-hydroxy-4-methylcoumarin and substituted triazole derivatives. *J. Chem.*, 6: S239-S246.
- Lever, A.B.P., 1984. *Inorganic Electronic Spectroscopy*. 2nd Edn., Elsevier, Amsterdam, pp: 175.
- Mishra, A.P. and M. Soni, 2008. Synthesis, structural and biological studies of some Schiff bases and their metal complexes. *Metal Based Drugs*, 2008: 1-7.
- Mohamed, G.G., M.M. Omar and A.M. Hindy, 2006. Metal complexes of Schiff bases: Preparation, characterization and biological activity. *Turk. J. Chem.*, 30: 361-382.
- Nakamoto, K., 2009. *Infrared and Raman Spectra of Inorganic and Coordination Compounds: Theory and Applications in Inorganic Chemistry*. 6th Edn., John Wiley and Sons, USA., pp: 419.
- Nicholls, D., 1973. *The Chemistry of Iron, Cobalt and Nickel*. Pergamon Press, Oxford, UK., Pages: 1161.
- Onal, Z., H. Zengin and M. Sonmez, 2011. Synthesis, characterization and photoluminescence properties of Cu(II), Co(II), Ni(II) and Zn(II) complexes of N-aminopyrimidine-2-thione. *Turkish J. Chem.*, 35: 905-914.
- Parikh, V.M., 1974. *Absorption Spectroscopy of Organic Molecules*. Addison-Wesley, Boston, Massachusetts, USA., Pages: 325.
- Porter, Q.N., 1985. *Mass Spectrometry of Heterocyclic Compounds*. 2nd Edn., John Wiley & Sons, Hoboken, New Jersey, USA., ISBN:9780471099017, Pages: 966.
- Pouralimardan, O., A.C. Chamayou, C. Janiak and H. Hosseini-Monfared, 2007. Hydrazone schiff base-manganese (II) complexes: Synthesis, crystal structure and catalytic reactivity. *Inorg. Chim. Acta*, 360: 1599-1608.
- Rasheed, E.M., S.M. Saleh and S.M. Mohammed, 2014. Synthesis and spectral study of mixed ligand complexes of N, N-bis (4-nitro phenyl) malonyl dihydrazide and 1, 10-Phenanthroline with Cr (III), Co (II), Ni (II), Cu (II), Zn (II) and Cd (II) Ions. *Diyala J. Pure Sci.*, 10: 108-123.

- Silverstein, R.M., G.C. Bassler and T.C. Morrill, 1981. Spectrometric Identification of Organic Compounds. 4th Edn., John Wiley and Sons, New York, USA.
- Singh, D.P., V. Malik, R. Kumar, K. Kumar and S.S. Dhiman, 2010. Synthesis, characterization and antibacterial and antifungal studies of some tetraazamacrocyclic complexes. *J. Serb. Chem. Soc.*, 75: 1369-1380.
- Sommez, M. and M. Sekerci, 2002. Synthesis and characterization of Cu(II), Co(II), Ni(II) and Zn(II) Schiff base complexes from 1-amino-5-benzoyl-4-phenyl-1H-pyrimidine-2-one with salicylaldehyde. *Polish J. Chem.*, 76: 907-914.
- Srivastava, A.N., N.P. Singh and C.K. Shrivastaw, 2014. Synthesis and characterization of bioactive binuclear transition metal complexes of Schiff base ligand derived from 4-amino-pyrimidine-2-one, diacetyl and glycine. *J. Serb. Chem. Soc.*, 79: 421-433.
- Tharmaraj, P., D. Kodimunthiri, C.D. Sheela and S.C.S. Priya, 2009. Synthesis, spectral studies and antibacterial activity of Cu (II), Co (II) and Ni (II) complexes of 1-(2-hydroxyphenyl)-3-phenyl-2-propen-1-one, N2-[(3, 5-dimethyl-1H-pyrazol--1-yl) methyl] hydrazone. *J. Serb. Chem. Soc.*, 74: 927-938.
- Yaul, S.R., A.R. Yaul, G.B. Pethe and A.S. Aswar, 2009. Synthesis and characterization of transition metal complexes with N, O-chelating hydrazone Schiff base ligand. *Am. Eurasian J. Sci. Res.*, 4: 229-234.