

Studies on Shoot Regeneration of *Brassica juncea* Var. Poorbiraya Under Salt Stress Conditions

Athar Hussain, Mubashrah Munir and Sumaira Abbas
Department of Botany, Government College University Lahore, Pakistan

Abstract: Different salts NaCl, CaCl₂.2H₂O, MgSO₄.7H₂O and their combination, NaCl + CaCl₂ + MgSO₄.7H₂O were employed to establish different equimolar strengths i.e. 50mM, 100mM, 150mM and 200mM to study their effect on *Brassica Juncea* shoot regeneration from leaf explant. Best shoot regeneration in percentage (80%), number (8), length (10mm) was recorded in 50mM of NaCl+MgSO₄.7H₂O.+CaCl₂. 2H₂O. CaCl₂ proved to be more deleterious in comparable concentration as that of NaCl and MgSO₄.7H₂O. No age and position factor was observed for shoot regeneration however it was comparatively more towards lower halves of leaf lamina.

Key-words: Regeneration, *Brassica juncea* and stress conditions

Introduction

Brassica oil seed crop occupy over 11 million hectares of world agricultural land and provides 8% of the major oil crops under varied climatic conditions (Downey, 1983). Rapidly cycling nature of *Brassica* species have shaped them as a model plants for research in genetics, molecular biology and physiology (William and Hills, 1986).

Salinity and drought are the two major environmental stresses that are world wide in distribution and limiting the growth and productivity of plants community (except halophytes) including *Brassica* (Kavi kishore *et al.*, 1995; Rudulier *et al.*, 1984; Skirver and Mundy, 1990). The rich agricultural lands of Pakistan have no exception in facing the great danger of salinity. Therefore it is desirable to obtain high frequency of plant generation, which can grow under these stress conditions. *Brassica* can grow under saline conditions, which includes many important oil yielding vegetables and forage crop (Gangopadhyay *et al.*, 1997).

A great number of factors are reported to be involved in plant regeneration of *Brassica* spp. from cotyledons (George and Rao, 1980) peduncle (Stringham, 1977) and leaf disc explants (Eapen and George, 1996; Naqvi *et al.*, 1995).

Regeneration system characterized by a high frequency of shoot formation would be useful in allowing the study of biochemical events associated with shoot induction and wide spread application of gene transfer techniques for crop improvement. In spite of being a major oil seed crop of Indo-Pak subcontinent, little work has been done on *Brassica juncea* in reference to osmotic stress (Kumar, 1984).

It has been shown that *Brassica Juncea* controls glucose metabolism and reduces lipid peroxidation as well as level of oxygen radicals, ameliorating the damage caused by oxidative stress in diabetes (Young *et al.*, 2003). Further more *Brassica juncea* have shown

varying degree of hypoglycaemic and antiglycemic activity (Grover *et al.*, 2002). These abilities have enhanced studies in *Brassica Juncea*.

Materials and Methods

The seeds collected from NARC (National Agricultural Research Council) Islamabad were aseptically grown in MS medium (Murashige and Skoog, 1962) supplemented with 3.0 mg/l NAA and 1.0 mg/l BAP. Plumules were grown on the same medium for three weeks to get leaf explants. Sugar used was 3 % in all experiments and all the media were solidified with 1.25% agar. Routinely 30 ml of solidified media were dispensed into 250 ml culture flasks plugged with non-absorbent cotton covered in one layer of muslin cloth. All cultures were maintained at 26 ± 1 °C fewer than 3 k lux cool florescent light for fifteen hours photoperiod. Callus was sub-cultured after 5 weeks. All the MS media were supplemented with 50, 100, 150 and 200 mM of NaCl, MgSO₄.7H₂O, CaCl₂.2H₂O either separately or equimolar combinations. A method of constant salt concentration instead of stepwise approach was used to get salt adapted calli. The explant was treated in five replicate per treatment. The electroconductivity of saline solutions used is given in Table 1.

Result

Leaf explants, 5 to 6 days old showed curling and callusing at the points touching the media resulting compact and non-watery calli which were less vigorous and healthy than the control one. Browning of the callus was observed with an increase in age and salt concentration of the medium. Regenerated shoots were appeared when callus was 3-5 weeks old.. Regenerated shoots from all the media had rammo-colour (off-white to bright green), however yellowing and whitening of the leaves (older) from the regenerated shoots did

Table 1: The range of Electroconductivities of saline solutions

Medium	50mM	100mM	150mM	200mM
NaCl	5.5 x 10 ⁴	5.5 x 10 ⁴	6.855 x 10 ⁴	6.98 x 10 ⁴
CaCl ₂ .2H ₂ O	6.38 x 10 ⁴	6.81 x 10 ⁴	7.72 x 10 ⁴	6.55 x 10 ⁴
MgSO ₄ .7H ₂ O	5.93 x 10 ⁴	6.19 x 10 ⁴	6.54 x 10 ⁴	8.06 x 10 ⁴
NaCl + CaCl ₂	6.86 x 10 ⁴	6.95 x 10 ⁴	8.39 x 10 ⁴	11.47 x 10 ⁴

Table 2: Shoot regeneration response under different salt stress condition

Character studied	Concentration															
	NaCl (mM)				CaCl ₂ .2H ₂ O (mM)				MgSO ₄ .7H ₂ O (mM)				NaCl + CaCl ₂ . 2H ₂ O + MgSO ₄ .7H ₂ O (mM)			
	50	100	150	200	50	100	150	200	50	100	150	200	50	100	150	200
Callusing	+++	+++	++	++	++	++	+	+	+++	++	++	++	+++	+++	++	++
% of shoot regeneration	80	65	60	40	15	10	5	5	55	45	45	40	80	80	40	40
Mean number of regenerated shoots	4	3	3	2	3	1	1	1	4	3	2	2	8	4	2	1
Mean length of regenerated shoot (mm)	6	4	4	3	8	5	3	3	5	2	2	2	10	5	3	2

+ / + + / + + + = Slow, moderate and rapid callus growth respectively
 3 mg/l NAA and 1 mg/l BAP were supplemented in all these media

appear in the saline media of higher concentration (150-200mM) 200mM equimolar concentration of NaCl + MgSO₄.7H₂O + CaCl₂.2H₂O, gave highest percentage of shoot regeneration (80% in two cultures) and mean number of regenerated shoots (8) among all the same concentrations of these salts separately. CaCl₂ proved to be more deleterious in comparable concentration as that of NaCl and MgSO₄.7H₂O (Table 2). No age and position factor was observed for shoot regeneration however it was comparatively more towards lower halves of leaf lamina.

Discussion

Decline in the callus growth due to salt concentration (150 - 200 mM) observed in present study is a usual phenomenon in *Brassica juncea*. (Sharma *et al.*, 1991) and in other plants as well (Cushman *et al.*, 1990, Greenway and Munns, 1980; Reddy and Vaidyanath, 1986). This retardation of growth may be due to the fact that certain amount of the total energy available for tissue metabolism is channelled to resist the stress (Cushman *et al.*, 1990). Several reports of plant regeneration from seedling explants of Brassica species have been published during the past decade. However there is a considerable variation on the observation of different groups even when dealing with the same species (Lazzeri and Dunwell, 1986 and Murata and Orton, 1987). Equimolar concentration of NaCl + CaCl₂ 2H₂O + MgSO₄.7H₂O were applied as a source of salinity. It is believed that responses to this combination may be more representative of a generalized plant response to calamity (He and Cramer, 1992). Maximum shoot regeneration at this combination may be attributed to the stimulatory effect

of sulphate in the medium (Eapon and George, 1996) and particularly to some regulatory substance (s) emanating from lamina (Sharma *et al.*, 1991) and greater leaf turgor pressure (Wright *et al.*, 1995) in *Brassica juncea*. This study found salt stress, beneficial for regeneration which is in accordance with previous reports (Yoshida *et al.*, 1983, Reddy and Vaidyanath, 1986) as salt tolerance is considered nature of embryonic cells that tolerate a higher level of salt stress than the other cell types (Binh and Heszky, 1990). Proline is another factor, which can account for stimulating regeneration upto 80% (Rout *et al.*, 1995) because it has been found significantly due to an increase in salt concentration (Chandler and Thrope 1987, Jains *et al* 1991b, Kumar and Sharma 1989, Watad *et al.*, 1983, Gangopadhyay *et al.*, 1997). These results indicate beyond the genetic effect as shoot formation give rise to the question of cellular totipotency and of correlation influence. This system for efficient regeneration of *Brassica juncea* var. poorbiraya from leaf explant can be applied to improve cultivar selection for environmental stress at the cell levels.

References

Binh, D.O. and L.E. Heszky, 1990. Restoration of the regeneration potential of long term cell culture in rice (*Oryza sativa* L.) by salt pre-treatment. J.Plant physiol, 136: 336-340.
 Chandler, S.F. and T.A. Thrope, 1987.Characterization of growth water relations and proline accumulation in sodium sulphate tolerant callus of *Brassica napus* L. cv. Westar (Canola) Plant.Physiol, 84: 106 - 111.

- Cushman, J.C., E.J. De Rocher and H.J. Bohnent, 1990. Gene expression during adaptation to salt stress. IN: Katterman F. (ed) Environmental injury to plants (pp.173-203). Academic press. Inc. San Deigo.
- Downy, R. K., 1983. The origin and description of the Brassica oil seed crops. "In high and low crucic acid rape seed oils." Academic press Canada (pp.1-20).
- Eapen, S. and L. George, 1996. Enhancement in shoot regeneration from leaf discs of *Brassica juncea* L. by silver nitrate and silver thiosulphate. *Physiol.Mol. Biol. Plants*. 2: 83-86.
- Gangopadhyay, G., S. Baso and S. Gupta, 1997. In vitro selection and physiological characterization of NaCl and mannitol adapted callus lines in *Brassica juncea*. *Plant Cell Tiss.Org. Cult.*, 50:161-169.
- George, L. and P.S. Rao, 1980. In vitro regeneration of mustard plants (*Brassica juncea* Var. Rai 5) on cotyledon explants from non-irradiated and mutagen treated seed. *Ann. Bot.* 46: 107-112.
- Greenway, H. and R. Mnns, 1980. Mechanism of salt tolerance in nonhalophytes *Ann.Rev. Plant physiol.* 31:19-190.
- Grover, J.K., S.Yadav and V.Vats, 2002. Medicinal plant of India with anti diabetic potential. *J.Ethopharmacol* , 81: 81-100
- He, T. and G.R. Cramer, 1992. Growth and mineral nutrition of six rapid cycling *Brassica spp.* In response to sea water salinity. *Plant soil*, 139:285-294.
- Jains, S., H.S. Nainawata, R.K. Jains and J.B. Chowdhory, 1991b. Proline status of genetically stable salt tolerant *Brassica juncea* L. Somaclones and their parents cv. Prakash. *Plant Cell Rep.* 9:684-687.
- Kavi-Kishore, P.B., Z. Hang, G.H. Miao, C.A.A. Hu and D.P.S. Verma, 1995. Over expression of 1-pyrrolone-5-carboxylate synthetase increase proline production and confers osmo-tolerance in transgenic plants. *Plant.Physiol.* 108:1387-1394.
- Kumar, V., 1984. The value of certain parameters as an index for salt tolerance in Indian mustard (*Brassica juncea* L.) *Plant soil.*, 79:261-271.
- Kumar, V. and D.R.Sharma, 1989. Isolation and characterization of NaCl resistant callus of *Vigna radiata* L. Wilezek Var radiata. *J. Expt.Bot.*, 40: 143-147.
- Lazzeri, P.A. and J.M. Dunwell, 1986. *In vitro* regeneration from seedling organs of *Brassica Oleracea* var. Italic Plenck Cv. Green Comet. I. Effect of plant growth regulators. *Ann. Bot.* 58:689-697.
- Murashige, T. and F. Skoog, 1962. A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiologia Plantarum*, 15: 473-497.
- Murata, M. and T. J. Orton, 1987. Callus initiation and regeneration capacities in *Brassica spp.* *Plant cell Tiss. Org. Cult.* 11:111-123.
- Naqvi, Z., A. Hussain and S.Abbas, 1995. Callus induction capabilities of various tissues of *in vitro* grown *Brassica juncea* var. Poorbirays. *Biologia*, 41:99-102.
- Reddy, P.J. and K. Vaidyanath, 1986. In vitro Characterization of salt tolerance stress effects and the selection of salt tolerant plants in rice (*Oryza sativa* L.) *Theor. Appl. Genet.* 71:757-760.
- Rout, G.R., S. Samantary and P. Das, 1995. Somatic embryogenesis and plant regeneration from callus culture of *Acacia catechua* multipurpose leguminous tree. *Plant Cell Tiss. Org. Cult.* 42:283-285.
- Rudulier, D.L., A.R. Storm, A.M. Dondekar, L.T. Smith and R.C. Valentine, 1984. Molecular biology of osmoregulation. *Sci.*, 224:1064-1068.
- Sharma, K.K., S.S. Bhajawani and T.A. Thrope, 1991. The role of Shoots and roots from cotyledon explants of *Brassica juncea* L. *Czern Plant Cell Tiss. Org. Cult.* 24:55-59.
- Skirver, K. and J. Mundy, 1990. Gene expression in response to abscisic acid and osmotic stress. *Plant cell* 2:503-512.
- Stringham, G.R., 1977. Regeneration in stem explant of haploid rape seed (*Brassica napus* L.) *Plant Sci. Lett.* 9: 115-119.
- Watad, A.E.A., L. Reinhold and H.R. Lerner, 1983. Comparison between a stable NaCl-selected *Nicotiana* Cell line and the wild type *Plant physiol.* 7:624-629.
- William, P.H. and C.B. Hill, 1986. Rapid cycling population of *Brassica*. *Sci.*, 232:1385-1389.
- Wright, P.R., J.M. Morgin., R.S. Jessop, A.A.Cass, 1995. Comparative adaptation of Canola (*Brassica napus*) and Indian mustard (*B. juncea*) to soil water deficits, yield and yield components, *Field Crops Research* 42:1-13.
- Yoshida, S., M. Ogawa, K. Suenaga and H.C. Ye, 1983. Indication and selection of salt tolerant mutant rice by tissue culture. Recent progress at IRRI. In: *Cell Tissue Culture techniques for cereal crop improvement.* pp:237-254. Science press, Beijing.
- Young, K.H., T.Yokozawa, E.J.Cho, H.C.Silk, J.C.Sue and H. C. Young, 2003. *In vitro* antioxidant effects of mustard leaf (*Brassica Juncea*). *Phytophton*, 17: 465-71.