

Investigation into the Chemical Composition of the Dry Fruit of *Tetrapleura tetraptera* (Ubukirihu)

T.A. Abii and Elegalam Amarachi

Department of Chemistry, Michael Okpara University of Agriculture, Umudike,
P.M.B 7267 Umuahia Abia State, Nigeria

Abstract: The proximate, mineral and phytochemical composition of *tetrapleura tetraptera* were investigated using standard methods. The result of the phytochemical analysis showed the presence of alkaloids (1%), flavonoids (2.5%) saponins (20%) and tannins (0.12%). The proximate analysis gave ash (9%), fibre (45%) oil (4%), moisture (3%) crude protein (5.6), while the mineral analysis revealed in mg 100 g⁻¹ Ca (1.68), Fe (12.02), K (7.74), Zn (10.69), Na (4.74), Mg (2.4), P (1.3), Cd (BDL) Pb (BDL). The relevance of these constituents to the native use of this fruit was discussed.

Key words: Investigation, chemical composition, dry fruit, *tetrapleura tetraptera*, grave ash, alkaloids

INTRODUCTION

Tetrapleura tetraptera locally known as (uhiokirihu) in South Eastern Nigeria belongs to the mimosaceae family. The plant is found in most deciduous forest type and has a very distinctive four winged fruit in loops. The shiny glabrous dark, brown fruit hangs persistently at the ends of branches on sprout stalks. The plant is hard with heavy wood which is used for firewood, building poles, pestle, tool handle and carvings. The fruit is used extensively in traditional medicine for the management of an array of human ailments including arthritis, asthma, diabetes mellitus, hypertension, epilepsy, schistosomiasis etc. John *et al.* (2004). It is claimed to be therapeutically useful in the treatment of leprosy and rheumatoid pains (Daiziel, 1984). The dry fruit has a characteristic pleasant aroma which makes it a popular seasoning spice in southern and Eastern Nigeria (Essien *et al.*, 1994; Adesina, 1982; Okwu, 2004) while Ghanaians use it as vitamin. It is used extensively in soups of nursing mothers to prevent post partum contractions (Nwawu and Alah, 1986) and gastro-intestinal disorders especially stomach ulceration (Noamesi *et al.*, 1992). In cold weather it is used to prepare pepper soup and the aroma is believed to drive away snakes. The dry powdered fruit has been formulated into soap to increase the antimicrobial activity and improve the foaming and hardness of soaps (Adebayo *et al.*, 2000). The fruit is said to contain caffeic acid (Adesina, 1982) which is reported to have antitumor activity, anti-inflammatory properties and also inhibit HIV replication. The flavouring property of the fruit has been used to improve the flavour of cassava fufu by eliminating the pungent odour and inhibiting fungal

growth (Okwu, 2004). Although much research has been done on this plant *tetrapleura tetraptera* (Adegunmi *et al.*, 1989, 1991; Daiziel, 1984; Millard *et al.*, 1989) (Adebayo *et al.*, 2000; Aka and Nwabie, 1993; Salak *et al.*, 1990; Adesina, 1982) but little has been reported on the fruit and its metallic and chemical contents. This research examines specifically the metal content of the dry fruit and their correlation with the potential traditional medical uses so far reported.

MATERIALS AND METHODS

The fruits of *tetrapleura tetraptera* were bought from different markets in Abia state of Nigeria and identified by a taxonomist at the botany department of Michael Okpara University of Agriculture Umudike Nigeria. The dry fruits were manually washed with distilled water and residual moisture was dried in an oven at temperature of 60°C. The dried fruits were ground into fine powder using mortar and pestle and sieved through 20 mesh sieve and stored in air tight bottles.

Mineral analysis: Two grams of the ground powder was digested with concentrated nitric acid. The resulting solution was evaporated to dryness and redissolved in 100 mL deionised water. The solution was analysed for the minerals K, Fe, Zn, Ca, Na, Pb and Cd using atomic absorption spectrophotometer. Phosphorus was analysed colorimetrically using phosphor-vanadomolybdate method, sodium content was determined using flame photometer while magnesium was analyzed complexometrically (AOAC, 1980).

Proximate analysis: The ash, oil, moisture, fibre and crude protein contents were determined by standard method (AOAC, 1980). Ash content was determined by incineration of 2 g of the sample powder placed in a muffle furnace maintained at 600°C for 8 h. The % residue weighed was expressed as ash content. Oil was determined by exhaustively extracting 2 g sample with petroleum ether using the soxhlet method. Moisture was determined by drying 2 g sample in an oven at 105°C for 24 h. The difference in wt gave the moisture content. Nitrogen was determined using the kjeldahl method and crude protein was calculated by multiplying the percentage nitrogen content by the conversion factor of 6.25 (AOAC, 1980).

Phytochemicals: The phytochemicals were assayed using standard methods. Flavonoids and alkaloids were determined using the method of Trease and Evans (1989). Saponins and tanins were analysed by the method of Herborne (1978).

RESULTS AND DISCUSSION

Results of the analysis is as presented in Table 1-3. Table 1 shows the mineral content, Table 2 is the phytochemicals while Tproximate analysis. Table 1 shows the presence of mineral elements that are essential in human nutrition. Iron which tops the list with a concentration of 410 mg 100 g⁻¹ sample performs functions of vital importance in the body. It is found in the blood being part of the hemoglobin, which is responsible for the red colour and allows the transportation of oxygen from the heart to all the cells. It is essential for the oxidation of carbohydrates, proteins and fats (Adeyeye and Otokiti, 1999). From the result *tetrapleura tetraptera* fruit has higher iron content than some dry fruits pepper, garlic, fig, chenu with concentration in mg 100 g⁻¹ sample of 1.2, 1.7, 2.23, 1.01, respectively (George *et al.*, 2004). Its iron content is however lower than those of some herbal plant found in Nigeria (30-59) mg 100 g⁻¹ (Isong and Idiong, 1997). Despite the effect of antinutritional factors against bioavailability of iron (Ladian *et al.*, 1996) this amount of iron is more than adequate since the human body requires only about 1.00 mg per day of iron to balance intake and excretion (Bothwell *et al.*, 1989). This high concentration of iron in *tetrapleura tetraptera* fruit is probably the reason for its use by lactating mothers to regenerate lost blood. Another element of high concentration is zinc (10.5 mg 100 g⁻¹). Zn is a very useful medicinal trace element in human body. It provides a natural protective mechanism against virus especially those causing

Table 1: Mineral content of tetraptera of tetrapleura tetraptera

Mineral	Concentration in mg 100 g ⁻¹
Calcium	0.0168
Iron	0.1202
Sodium	0.0474
Magnesium	0.0024
Lead	BDL
Potassium	0.774
Zinc	0.1069
Cadmium	BDL
Phosphorus	0.13

Table 2: Phytochemical contents of tetrapleura tetraptera

Phytochemical	% determined
Alkaloid	1
Saponis	20
Flavonoid	2.5
Tannins	0.12

Table 3: Proximate composition of tetrapleura tetraptera

Constituent	% determined
Ash	9
Fibre	45
Oil	4
Moisture	3
Crude protein	5.6

respiratory track infections (Rowland, 2004). It is used extensively in the war against HIV and is believed to have a way in which it delays the integration of the HIV virus in the blood. Mochegiari *et al.* (1995), Sadler (2004) and Raynolds (2003). This reinforces the belief of traditional healers in using the fruit of terapleura tetraptera to cure HIV, cold, etc Zn plays an important role in the development and functioning of the pituitary gland, the gonards and the reproductive organs (George, 2004). This is in agreement with the use of the fruit for nursing mothers and pregnant ones. The deficiency of Zn causes diarrhea, apathy immunosensitivity and mental depression (Strausel and Saltman, 2000; WHO, 1996). This supports the native use of the fruit to treat diarrhea and mental fatigue. The element manganese present with a concentration of 2.4 mg 100 g⁻¹ is another important trace element involved in calcium metabolism in bones and management of circulatory disease like Ischemic heart disease Magnesium serves as a catalyst in energy producing reactions within the cells and facilitates, muscles relaxation (George *et al.*, 2004). The concentration of mg is lower than that in some fruits-orange (10 mg), banana (29), apricot (8 mg), fig (17 mg), apple (7 mg) but higher than those reported for some medicinal plants (1.152-1.824) mg 100 g⁻¹ (Edeoga *et al.*, 2006). The presence of mg reinforces the use of the fruit to treat heart disease. Calcium present with a concentration of 1.68 mg 100 g⁻¹ fall within the range reported for some medicinal plants (1.034-1.946) mg 100 g⁻¹. (Edeogu *et al.*, 2006) but lower than those of some fruits-orange 40 mg 100 g⁻¹, cabbage. (George *et al.*, 2004). Calcium together with

phosphorus are involved in bone and teeth formation, blood clotting and transmission of nerve impulses. Phosphorus with a concentration of 1.13 mg 100 g⁻¹ is used in the maintenance of healthy bones and teeth, energy metabolism and acid base balance in the body (George *et al.*, 2004). In cold weather the use of this fruit to prepare hot pepper soup, which invigorates and activates the muscles and bones is not unconnected with the presence of mg, Ca and P. The absence of lead and cadmium in *T. tetraptera* make it very good for human consumption.

Phytochemicals: Table 2 shows the phytochemical analysis showing high concentration of saponins (20%), followed by flavonoids 2.5% and low Alkaloid 1% while tannin is 0.12%. The presence of phytochemical which are antioxidants partly explains the preventive and healing action of the fruit. Saponins can foam and can also hemolyse red blood cells (Hill, 1952). Flavonoids protect the heart and arteries (Rimm *et al.*, 1996). They prevent stroke and other vascular accidents (Keli *et al.*, 1996). These supports the traditional use of the fruit to manage heart related diseases, Alkaloids are involved in growth regulation with dramatic physiological activities hence their wide use in medicine (Harborne, 1973) (Wattenberg, 1990).

Proximate analysis: The proximate analysis result in table 111 shows fibre 45%, ash 9%, crude protein, 5.6% moisture 3% and fat 4%. The R.D.A of fibre for children, adult and pregnant woman are 19-25, 21-38 and 78%, respectively. Thus the fruit could be a good source of dietary fibre.

The RDA of fibre for children, adult and pregnant women are 19-25, 21-38 and 28%. Thus this T.T could be a good source of dietary fibre. The ash content, which signifies the index of mineral content in biota, agrees with the high mineral contents observed. The research has shown that the fruit of *tetrapleura tetraptera* is a reservoir of medicinal constituents with a wide range of application potential. It should be used as a raw material in formulating drugs and cosmetics. A study of the antinutritional and toxicological factors is however, necessary.

REFERENCES

Adebayo, A.S., I.A. Gbadomosi and C.O. Adewunni, 2000. Formulation of antimicrobial dried powdered herbs in soap bases in: Phytomedicines in malaria and sexually transmitted diseases; challenges for the new millennium C.O. Adewunni and S.K. Adesina, Adv. Obafemi Awolowo University Ilse-Ife, pp: 97.

Adesina, S.K., 1982. Int. J. Crude Drug Res., 20: 93-100.

Adewunmi, C.O., P. Furu and H. Madsen, 1989. Phytother. Res., 3: 81-84.

Adewunmi, C.O. and P. Furu, 1989. J. Ethiopharmacol., 27: 277-283.

Adewunmi *et al.*, 1991. Toxicological and environmental chemistry, 30: 69-74.

Adeyeye, E.I. and M.K.O. Otokiti, 1999. Proximate Composition and some Metritionally valuable minerals of two varieties of capsicum annum (Bell and Cherry papers) Discovery and Innovation 11: 75 -81.

Aka, P.A. and A.I. Nwabie, 1993. Use of *T. tetraptera* as anti-culant, Fitoterapia, 64: 42.

AOAC, 1980. Official Method of Analysis (13th Edn.), Washington DC. Association of Official Analytical Chemists.

Daiziel, J.M., 1984. The useful plants of tropical West Africa: Crown Agents for overseas Governments and Administration London, pp: 223-244.

Edeoga, H.O. *et al.*, 2006. Chemical Composition of *hyptis sua veolens* and *Ocimum gratissimum* hybrids from Nigeria: Afr. J. Biotech., 5: 892-895.

Essien, E.U., B.C. Izunwane, C.Y. Aremu and O.U. Eka, 1994. Plant Foods for Human Nut., 45: 47-51.

George, D. Pamplana-Roger Mid, 2004. Encyclopedia of foods and their healing Power: Review and herald Publishing Mary Land USA., 1: 412.

Harborne, J.B., 1978. Textbook of Phytochemical Methods Champion and Hall. Ltd. London, pp: 110-113.

Harborne, J.B., 1993. Phytochemical methods A guide to modern technique in plant analysis: Chapman and Hill New York.

Hill, A.F., 1952. Economci Botany (2nd Edn.), Mc graw. Hill Book Company Inc., pp: 15.

Isong, E.U. and U.I. Idiong, 1997. Comparative Studies on the Nutritional and Toxic composition of three varieties of *Lesianthera Africana* Plant Food for Human Nutr., 51: 79 -84.

John, A.O. Ojewole and O.A. Clement, 2004. Anti-inflammatory and Hypoglycemic effects of *tetrapleura teraptera*: J. Ethnopharmacol., 95: 177-182.

Keli, S.O., M.G. Hertog and E.J. Feskens, 1996. Dietary flavonoids, antioxidant vitamins and incidence of strock; The zutphen study. Arch. Int. Med., 156: 637-642.

Ladian, J. *et al.*, 1996. Nutrient composition of some green leafy vegetables consumed in Sokoto Nigeria; J. Basic and Applied Sci., 5: 39-44.

- Mochegiari, E., S. Veccia, F.K. Ancarani, G. Scalise and N. Fabris, 1995. Benefit of oral zinc supply, mentalin as an Adjust to Zidovadine (AZI) therapy against oportunist infections in Aid; *Int-immunopharmac*, 17: 719-727.
- Noamesi, B.K., J.F. Mensah, E. Dagne and M. Bogale, 1992. Effect of tetrapleura tetraptera; Seventh symposium on medicinal plants, spices and other natural product: Abstract No Wp -20 Manila.
- Nwawu, J.I. and P.A. Alah, 1986. Effect of tetraplaura teraptera; *J. Ethnopharmacol.*, 18: 103-107.
- Millard, M. *et al.*, 1989. *Helvetica Chimica Acta*, 72: 668-672.
- Okwu, D.O., 2001. Evaluation of the chemical composition of indigenous spices and flavouring agents: *Global J. Pure and Applied Sci.*, 7: 445-459.
- Rowland, T.C., 2004. Zinc ions and Sars Zn news: *Iza*, 63: 82-91.
- Rowlands, T.C., 2004. *Weuropsychopharmacol. Biopsy-Chaiatry*, 28: 181-190.
- Rimm, E.B., M.B. Katan and A. Ascherio *et al.*, 1996. Relation between intake of flavonoids and risk for coronary heart disease in male health professionals. *Ann. Int. Med.*, 125: 384 -389.
- Salako, Q., U.E. Akpau, E.I. Ette, E.E. Essien and O. Ipeaiyeda, 1990. *Fitoterapia*, 61: 169-171.
- Sadler, K.F., 2004. *Neuropsychopharmacol Biopsy-Chiatry*, 28: 181-190.
- Strausel and P. Saltman, 2000. Trace mineral interactions. Department of Biology, University of California at San Diego.
- Trease, G.E. and W. Evans, 1989. *Pharmacognosy*. (11th Edn.), Braillier Tiridel and Macmillan Publishers.
- Wattenberg, L.W., 1990. Inhibition of cercinogenesis by minor nutrient constituents of the diet; *Proc. Nutr. Soc.*, 49: 173-183.
- WHO, 1996. Trace element in Human Nutrition and Health; Geneva, pp: 3-5, 72: 105, 209.