

## Performance Comparison of Carbofuran and Bio-Nematicidal Potentials of the Extracts from Rattle Weed and Nitta Plants on Root Knot Nematode Pest of Pepper

<sup>1</sup>T.I. Olabiyi and <sup>2</sup>E.E.A. Oyedunmade

<sup>1</sup>Faculty of Business, Environment and Society, 1 Priory Street, Coventry University, CV1 5FB, Coventry, United Kingdom

<sup>2</sup>Department of Crop Protection, Faculty of Agriculture, University of Ilorin, Ilorin, Nigeria

**Abstract:** Two years field trial were carried out to compare the effectiveness of carbofuran and plant extracts from the roots and leaves of rattle weed and nitta plant in root knot nematode control on pepper. Carbofuran was applied at 1.5 and 2.5 kg a.i ha<sup>-1</sup>, while plant extracts were applied at 25, 50, 75 and 100% concentrations. An experimental plot where neither carbofuran nor plant extract was applied served as the control treatment. Distilled water and methanol served as media of extraction from rattle weed and nitta plant. Both carbofuran and plant extracts significantly, reduced the soil population of root-knot nematode and galling on the root of pepper and also improved growth and yield of pepper. Pepper plants treated with carbofuran and 100% concentration of the plant extracts gave the best control of the root-knot nematode.

**Key words:** Carbofuran, rattle and nitta plants, control, root-knot nematode, pepper, Nigeria

### INTRODUCTION

Pepper, *Capsicum annum*, is an important crop of high economic and nutritional importance in Nigeria. On the basis of research that was done by Erinle (1989), Nigeria is the largest producer of pepper in Africa (50% production) and was rated second in the world (1979). Yamaguchi (1982) wrote that pepper is an important soup ingredient that supplies essential nutrients that may not be present in staple food items.

On the basis of research that were done by Waliullah (1992), Yousef and S'Jacob (1994), Olabiyi and Ndana (2003) and Olabiyi (2004), root knot nematodes (*Meloidogyne* sp.) was been identified as economic pathogens of vegetables. Richard and Nicola (1990) said root knot nematodes are recognised on plants by the presence of galls on the plant roots. Above ground symptoms include stunting, chlorosis, wilting and early senescence.

In the time years, chemical nematicides (carbofuran, vertimec, vydate oxamy, ethoprop, cadusafos, fenamiphos and aldicard) had been in use by numerous farmers. Restriction on the importation of chemical through the international market made chemical nematicide to be scarce and expensive, even beyond the reach of poor resource farmers in Nigeria. Moreover, the hazard effects

of chemical nematicide on both target and non-target organisms and also on the environment (as pollutant) had brought much discouragement to many Nigerian farmers.

Between 1999 and 2002, severe root knot nematode infections, which caused complete crop (100%) loss on some pepper plants, were observed in the Horticultural garden of the Osun State College of Education, Ila-Orangun, Nigeria. As a result of the field root knot nematode population reaching an epidemic level, an attempt was made to control the root knot nematode disease on pepper in 2003 and 2004 cropping seasons using botanicals. Moreover, several investigators like Pandey (1990), Abid and Maqbool (1991), D' Abbaddo and Sasanelli (1996) and Olabiyi and Ndana (2003) have carried out studies on the use botanicals as alternative control measure against plant parasitic nematodes but not on the root and leaf extracts of rattle weed and nitta plants.

In Nigeria, efforts are being geared toward the search for plant materials that could be used in the control of root knot nematode pathogenic organism on the economic food crops. In this study, attempts were made to compare the performances of carbofuran with that of extracts from the roots and leaves of rattle weed and nitta plants in the control of root knot nematode disease on pepper.

## MATERIALS AND METHODS

Two consecutive years' field experiments were carried out during the 2003 and 2004 cropping seasons at the Horticultural garden of the Osun state College of Education, Ila-Orangun, Nigeria. The experimental site soil type was sand-loam (55 sand, 16 clay and 29% silt) with pH 5.8. Two levels (1.5 and 2.5 kg a.i ha<sup>-1</sup>) of carbofuran (Furadan) and 4 concentrations (25, 50, 75 and 100%) each of aqueous and methanol extracts of rattle weed and nitta root were tested for the control of root-knot nematode, *Meloidogne incognita*, on pepper, *Capsicum annum*. An experimental plot where neither carbofuran nor plant extracts (0% treatment concentration) was applied served as control. The experiment was laid out as a randomized complete block design. There were 4 blocks with 1 m alleyway left in-between the blocks. Each block was subdivided into 35 treatment plots with each plot having a size of 5 m (length) by 1 m (breadth) and alleyway of 0.5 m in-between 1 plot and the other. Each treatment was replicated 4 times.

Pre-planting soil samples (zigzag form) were taken and assessed for nematode occurrence and abundance following the Southey (1986). Pepper seeds (cv.DT95/422) were bought from National Horticultural Research Institute (NIHORT), Ibadan, Nigeria. Five hundred pepper seedlings were raised in the nursery and transplanted at 4 weeks old at a spacing of 60 by 60 cm apart. The synthetic nematicide, carbofuran and plant extracts were applied close to pepper roots, 1 cm away from plant base and 5 cm into the soil, at 2 weeks after transplanting. Fertilizer N:P:K (15:15:15) was applied at the rate of 5 g per plant at 3 weeks after transplanting. Manual weeding was done at 4 weeks intervals during the period of the field experiment.

Plant height, number of leaves per plant, number of fruits per plant and fruit weight per plant were recorded. Post-harvest soil samples (zigzag form) were taken for nematode population assessment. Root gall indices were scored as used by Sasser *et al.* (1984) on a scale of 0-5, where 0 = no galls; 1 = 1-2% galls; 2 = 3-10% galls; 4 = 31-100% galls and 5 = more than 100% galls. The mean data of the 2 years were calculated and subjected to analysis of variance and means were partitioned using Tukey-Kramer's highest significant difference at the 5% probability level (S.A.S, 1997).

## RESULTS AND DISCUSSION

The effects of different treatments of carbofuran and plant extracts on the plant height of pepper are presented in Table 1. It was observed that plant height at 10 weeks

after transplanting (WAT) did not vary significantly ( $p < 0.05$ ) among the treatments. Pepper plants treated with 100% concentrations of the test plants and carbofuran (1.5 and 2.5 kg a.i ha<sup>-1</sup>) grew significant taller at 12 and 14 WAT than pepper plants treated with lower test plant extract concentrations and the control experiment.

Comparative effects of carbofuran and plant extracts on the average number of leaves of pepper infected with *M. incognita* are presented in Table 1. No significant differences between the number of leaves on pepper plants occurred at 10 WAT. However, significant differences became prominent on number of leaves at 12 and 14 WAT. It was observed that the number of leaves of pepper plant at either 12 or 14 WAT was significantly higher in pepper plants treated with either carbofuran or 100% concentration of the plant extracts than in pepper plants treated with lower test plant extract concentration. The control treatment had the least number of leaves per pepper plant.

The results of this research are in agreement with earlier investigations of Abid and Maqbool (1991), Olabiyi (2004) and Oyedunmade *et al.* (2001). These researchers reported that carbofuran and plant extracts significantly increased vegetative growth (plant height and number of leaves per plant) of tomato, egg plant, okra and soybeans.

Number of fruit and fruit weight per plant were significantly higher in pepper plants that were treated with carbofuran than other treatments (Table 2). It was observed that the 100% plant extracts in most cases resulted in higher number of fruit and fruit weight per plant than pepper plants treated with lower test plant extract concentrations. Pepper plants that did not receive any treatment had the least number of fruit and lowest fruit weight amongst all the treatments. This is in accordance with Oyedunmade *et al.* (2001) saying that 100% neem leaf extract and carbofuran, singly and in combination, significantly increased number of fruits and fruit weight per okra plant.

Table 2 also shows that carbofuran and 100% concentration of the plant extract significantly reduced the soil root knot nematode population and consequently roots galls as compared with lower plant extracts and the control. Olabiyi (2004) said that the roots of rattle weed and nitta plants contained saponins which are nematotoxic. In an *in vitro* experiment carried out with the use of saponins on eggs and juveniles of root-knot nematode, it was shown that saponins were inhibitory to root knot nematodes (Olabiyi, 2004). On the basis of research that was done by Oyedunmade *et al.* (2001) of Nigeria showed that there were significant reduction in root gall index and soil root knot nematode population as a result of a Vertimec (R) and plant material applied to root knot nematode infested soil grown with soybean.

Table 1: Comparative effects of carbofuran and plant extracts on the mean plant height (cm) and number of leaves of pepper infected with *Meloidogyne incognita* at 10,12, 14 weeks after transplanting (WAT)

Treatment	Concentrations (%)	Mean plant height (cm)			Mean number of leaves		
		10 WAT	12 WAT	14 WAT	10 WAT	12 WAT	14 WAT
Aqueous extract of Rattle weed leaf	25	5.3	7.3b	10.0c	5.5	9.5bc	15.8c
	50	5.5	7.5b	10.7c	6.8	11.0b	17.3c
	75	5.5	8.4b	12.7c	6.3	11.3b	18.5c
	100	5.6	10.6a	17.2a	7.3	13.3a	23.3b
Methanol extract of Rattle weed leaf	25	5.2	7.3b	10.1c	5.3	10.0c	15.0c
	50	5.3	7.7b	10.7c	6.0	11.5b	17.5c
	75	5.5	8.3b	12.5c	6.8	12.8b	19.3bc
	100	5.6	10.7a	18.0a	7.8	14.8a	22.3b
Aqueous extract of Rattle weed root	25	5.3	7.2b	10.1c	5.5	10.0c	14.3c
	50	5.4	7.9b	10.6c	6.0	12.0b	17.8c
	75	5.3	10.4b	12.5c	7.0	13.3a	19.8bc
	100	5.8	17.4a	18.2a	7.8	14.5a	22.8b
Methanol extract of Rattle weed root	25	5.3	7.2b	10.1c	5.3	10.3c	15.8c
	50	5.5	7.7b	11.4c	5.5	11.5b	17.5c
	75	5.5	9.4b	12.2c	5.8	11.8b	18.5c
	100	5.6	10.6a	18.2a	6.8	13.0a	23.5b
Aqueous extract of Nitta leaf	25	5.3	7.2b	10.3c	6.0	8.9c	16.5c
	50	5.3	7.1b	10.6c	7.0	10.3c	17.0c
	75	5.4	9.5b	13.1c	7.0	10.3c	21.0bc
	100	5.5	11.0a	17.2a	7.3	12.0b	21.5bc
Methanol extract of Nitta leaf	25	5.3	7.4b	11.1c	5.5	8.8c	15.8c
	50	5.4	7.8b	12.5c	5.5	9.8bc	17.0c
	75	5.3	7.8b	13.6c	6.3	10.5c	19.0bc
	100	5.3	10.6a	16.3ab	7.3	12.0b	28.3a
Aqueous extract of Nitta root	25	5.2	7.2b	10.7c	6.5	9.3bc	15.5c
	50	5.5	7.3b	11.0c	5.0	9.8bc	18.0c
	75	5.5	8.3b	12.3c	5.5	11.0b	19.5bc
	100	5.7	10.7a	16.2ab	6.3	12.0b	24.5b
Methanol extract of Nitta root	25	5.2	7.8b	11.7c	6.3	9.3bc	17.5c
	50	5.5	8.3b	12.1c	5.5	10.3c	18.8c
	75	5.6	8.9b	13.5c	6.0	11.0b	20.0bc
	100	5.5	10.9a	15.0a	6.0	11.0b	20.0bc
Carbofuran	1.5kg a.i ha <sup>-1</sup>	5.6	11.2a	17.5a	7.3	14.8a	30.1a
	2.5 kg a.i ha <sup>-1</sup>	5.5	11.6a	19.1a	7.8	14.9a	30.6a
Control	0	5.2	5.8c	9.2d	5.0	7.0d	10.8d
		NS			NS		

WAT = Weeks after transplanting; NS = Not significant; Numbers with same letter in the column are not significantly different. The means were partitioned using Tukey-Kramer Highest Significant Difference at 5% probability level; WAT = Weeks after transplanting; NS = Not significant; Numbers with same letter in the column are not significantly different. The means were partitioned using Tukey-Kramer Highest Significant Difference at 5% probability level

Table 2: Comparative effects of carbofuran and plant extracts on the mean number of fruit and fresh fruit weight (g) per plant, and soil nematode population (200 g soil) and root gall index of pepper infected with *Meloidogyne incognita*

Treatment	Concentration (%)	Number of fruit/plant	Fresh fruit weight (g)/plant	Initial <i>M.incognita</i> population	Final <i>M.incognita</i> population	Gall index
Aqueous extract of Rattle weed leaf	25	18.8d	62.3d	1298	600d	2.4c
	50	18.8d	68.0d	1300	565d	1.6c
	75	22.0c	96.8c	1310	388ab	0.4a
	100	22.0c	108.5c	1310	335a	0.1a
Methanol extract of Rattle weed leaf	25	18.0d	66.0d	1298	520c	2.3c
	50	20.0d	68.8d	1299	511c	1.4b
	75	20.8d	96.5c	1310	330a	0.5a
	100	24.8c	108.0b	1308	298a	0.3a
Aqueous extract of Rattle weed root	25	18.8d	64.5d	1298	521c	2.3c
	50	20.8c	66.8d	1298	488bc	1.3b
	75	22.8c	96.8c	1305	315a	0.1a
	100	24.0c	110.8b	1300	280a	0.1a
Methanol extract of Rattle weed root	25	18.5d	66.3d	1302	525c	2.4c
	50	18.8d	66.5d	1305	490bc	0.8b
	75	22.5c	94.8c	1298	337a	0.2a
	100	26.3b	108.5b	1298	341a	0.1a
Aqueous extract of Nitta leaf	25	20.3d	64.0d	1310	489bc	3.0d
	50	22.0c	76.5d	1305	497bc	1.3b
	75	22.8c	110.0b	1307	348a	0.5a
	100	26.8b	130.3a	1308	332a	0.2a

Table 2: Continue

Treatment	Concentration (%)	Number of fruit/plant.	Fresh fruit weight (g)/plant.	Initial <i>M.incognita</i> population	Final <i>M.incognita</i> population	Gall index
Methanol extract of nitta leaf	25	20.0d	62.9d	1304	570d	2.3c
	50	20.5d	72.3d	1305	497bc	1.1b
	75	22.3c	108.0b	1299	405b	0.6ab
	100	28.0a	128.0a	1298	328a	0.3a
Aqueous extract of nitta root	25	18.8d	64.0d	1310	558d	2.3c
	50	20.3d	66.3d	1308	507c	1.5b
	75	22.3c	96.5c	1309	340a	0.2a
	100	22.5c	108.8b	1305	331a	0.1a
Methanol extract of nitta root	25	18.8d	65.8d	1303	564d	2.2c
	50	18.5d	67.1d	1299	511c	0.9b
	75	22.0c	94.3c	1306	510c	1.1b
	100	26.5a	106.3b	1305	337a	0.2a
Carbofuran	1.5kg a.i ha <sup>-1</sup>	32.5a	130.9a	1298	315a	0.3a
	2.5kg a.i ha <sup>-1</sup>	34.5a	131.3a	1306	248a	0.0a
Control	0	14.3e	54.9e	1305	8529e	4.6e
NS						

Numbers with same letter in the column are not significantly different. The means were partitioned using Tukey-Kramer Highest Significant Difference at 5% probability level

### CONCLUSION

It could be concluded from this study that roots of rattle weed and nitta appear to hold promise as nematicidal plants and their effectiveness on the control of root-knot nematode at 100% concentration are at par with carbofuran.

### REFERENCES

Abid, A. and M.A. Maqbool, 1991. Effect of bare-root dip treatment in oil-cakes and neem leaf extracts on the root-knot development and growth in tomato and egg plant. *Pak. J. Nematol.*, 9 (1): 13-16.

D'Abbaddo, T. and N. Sasanelli, 1999. Effect of olive pomace soil amendment on *Meloidogyne incognita*. *Nematol. Medit.*, 24: 91-94.

Erinle, I.D., 1989. Present Status and Prospects for Increased Production of Tomato and Pepper in Northern Nigeria. In AVROC (Ed.). *Proc. Int. Symp. Integrated Manag. Prac.*, pp: 536-547.

Olabiya, T.I. and R.W. Ndana, 2003. Comparative effects of the water extracts of 4 plant materials on the root-knot nematode infection on cock's comb (*Celosia argentea* L.). *Zuma J. Pure. Applied Sci.*, 5 (2): 167-171.

Olabiya, T.I., 2004. Assessment of the nematicidal properties of extracts from tagetes erecta, *Ocimum gratissimum*, *Hyptis suaveolens* and *Crotalaria retusa*. Ph.D. Thesis submitted to University of Ilorin, Ilorin, Nigeria, pp: 177.

Oyedunmade, E.E.A., T.I. Olabiya and E.F. Okafor, 2001. Effect of neem (*Azadirachta indica*) leaf extract and carbofuran, singly and in combination, on growth and yield of okra on root-knot nematode infested soil. *NISEB J.*, 3: 223-227.

Oyedunmade, E.E.A., O.F. Ayo-Oke and T.I. Olabiya, 2001. Comparative efficacies of soil amendments and a synthetic nematicide on growth, yield and food components of a root-knot nematode infected soyabean (*Glycine max* L. Merrill). *NISEB J.*, 1 (3): 367-371.

Pandey, R., 1990. Studies on the phytonematotoxic properties in the extract of some medicinal and aromatic plants. *Int. Nemat. Network Newslett.*, 7: 19-20.

Richard, A.S. and G. Nicola, 1990. Nematode Parasites of Food Legumes. *Plant Parasitic Nematodes in Sub-Tropical and Tropical Agriculture*. Luc, M., R.A. Sikora and J. Bridge (Eds.). C.A.B. Int. pp: 197-198.

S.A.S. Institute, 1997. *SAS User's Guide Statistics*. Version 6.09 (Eds.). S.A.S. Institute, Cary NC, USA.

Sasser, J.N., C.C. Carter and K.M. Hartman, 1984. Standardization of host suitability studies and reporting of resistance to root-knot nematodes. *Coop. Pub. Dept. Plant Path. N.C. State University and USDA, Raleigh, N. C.*, pp: 7.

Southey, J.F., 1986. *Laboratory methods for work with plant and soil nematodes*. Ministry of Agriculture and fisheries, HMSO, London U.K., pp: 202.

Waliullah, M.I.S., 1992. Nematodes associated with vegetable crops in the Kashmir valley, India. *Nematol. Medit.*, 20: 47-48.

Yamaguchi, M., 1982. *World vegetables: Principles, production and nutritive values*. Cllis Horwood Chichester, England, pp: 145.

Yousef, D.M. and S' J.J. Jacob, 1994. A nematode survey of vegetable crops and some orchards in the Ghor of Jordan. *Nematol. Medit.*, 22: 11-15.