



Phytochemical, Pharmacognostic, Anti-Radical, Anti-Arthritic and Antibacterial Potential of the Leaf Extract of *Dialium guineense*

¹Ololade Zacchaeus S., ²Anuoluwa Iyadunni A and ³Ishaku Paul

¹Department of Chemistry, University of Medical Sciences, Ondo, Nigeria

²Department of Biological Sciences, University of Medical Sciences, Ondo, Nigeria

³Department of Chemical Evaluation and Research, National Agency for Food and Drug Administration and Control (NAFDAC), Lagos, Nigeria

Key words: *Dialium guineense*, leaf extract, secondary metabolite, pharmacological activities, evaluates

Corresponding Author:

Ololade Zacchaeus S.

Department of Chemistry, University of Medical Sciences, Ondo, Nigeria

Page No.: 192-203

Volume: 16, Issue 5, 2021

ISSN: 1815-932x

Research Journal of Applied Sciences

Copy Right: Medwell Publications

Abstract: This research evaluates the extract of the leaf of *Dialium guineense* for its phytochemical and pharmacognostic potential. GC-MS, UV-Visible Spectrophotometer and other established biochemical methods assays were used for the study. GC-MS analysis showed the presence of 37 phytochemicals. The main component was: 4-O-methylmannose (49.0%). The β -carotene, lycopene, total carotenoid, chlorophyll a, b, total chlorophyll, TPC, TFC, TTC, TAA values were 0.20 mgg^{-1} , 0.10 mgg^{-1} , 12.68 mgg^{-1} , 4.40 mgg^{-1} , 4.70 mgg^{-1} , 7.80 mgg^{-1} , $2,668.17 \pm 0.01 \text{ } \mu\text{gmg}^{-1}$, $17.80 \pm 0.00 \text{ } \mu\text{gmg}^{-1}$, $152.65 \pm 0.01 \text{ } \mu\text{gmg}^{-1}$, $43.75 \pm 0.04 \text{ } \mu\text{gmg}^{-1}$, respectively while Galvinoxyl and DPPH antioxidant IC_{50} were 125.0 and $6.0 \text{ } \mu\text{g mL}^{-1}$, respectively. The TAC value was $279.49 \pm 0.01 \text{ } \mu\text{gmg}^{-1}$. The highest anti-arthritic value was 96.00%. The extract inhibited the growth of all the screener bacteria. These results showed that the leaf can be used in the treatment of various human and animal's related aliment or diseases and more importantly in formulation food additives and preservative.

INTRODUCTION

Africa nations are naturally blessed with enormous biodiversity of edible plants which are used in consumption as food supplements and therapeutic purposes. Globally, medicinal plants including vegetables, herbs and trees; have been serve as a major and natural sources for human being to meet their daily health care and nutritional needs^[1-5]. Natural products especially plants represent an inexhaustible reservoir of novel molecules for new drug discovery. In many parts of rural African

rural community, medicinal plants are the most easily accessible and affordable health resource available to the local community and at times the only therapy that subsists. Nonetheless, there is still a paucity of updated comprehensive compilation of promising medicinal plants from the African continent^[3, 6-7]. Medicinal plants and vegetables grown in Africa are used for therapeutic purposes and as precursors for chemo pharmaceutical semi-synthesis^[9-12]. The phytochemicals present in the plant with pharmacological potentials could serve as lead sources for development of novel drugs for human and

animal uses. There has been tremendous increase in the interest of scientists in the field of pharmacognostic, food chemistry, pharmacy etc in tracking the health benefits of vegetables and other medicinal plants for their useful secondary metabolites and treatment of various diseases^[3, 4, 14]. Green vegetables contain varieties of metabolites that may act singly or synergistically to improve health of man and animal^[3, 15, 6, 16]. Many plants contain diverse of polyphenols that are excellent antioxidants to scavenge free radicals and inhibit lipid peroxidation^[18, 19, 6, 20]. Nowadays, many people prefer uses of natural antioxidants due to low resistance to high temperatures and the side effects such as carcinogenicity of some of the synthetic compounds^[21-23].

Dialium guineense (Velvet tamarind) is an evergreen fruit bearing tree in the family of *Fabaceae*. It belongs to the subfamily of *Dialioideae*^[24-25]. The plant is rich in micronutrients. It is a potential food supplement, rich sources of secondary metabolites used in traditional medicine. It has sweet edible pulp can be eaten raw or can be soaked in clean water for the production of fresh local juices^[24-31]. The plant exhibits natural pharmacological response against infectious and non-infectious diseases. It is used in the treatment of diseases such as heart disease, diarrhea, severe cough, bronchitis, stomach aches, malaria, fever, jaundice, hypertension and haemorrhoids, tooth decay and dental plaque^[32]. It has also been observed to possess some analgesic, wound healing, antioxidant, antiulcer and anti-vibrio properties^[31-35].

To the best of our knowledge, the phytochemistry, pharmacognostic and pharmacognostical activities of the leaf of this plant grown in Nigeria have not been extensively investigated despite the traditional uses of the green herb. Therefore, the present study was undertaken with the aim of looking into the phytochemical, carotenoid, chlorophyll, ascorbic acid, phenolic, flavonoid, tannin, contents, antioxidant, anti-arthritis and antibacterial potential of the leaf of the plant investigated.

Practical applications: This study revealed that the leaves of *D. guineense* have abundant bioactive secondary metabolites that may be applied in the formulation food or drug supplement and preservative. With its high medicinal potential and it could be alternative sources of natural therapeutic agents. The high antioxidant capability of the leaf extract shows that it can be used to prevent diverse forms of diseases such as: microbial, respiratory digestive, various chronic infectious, degenerative, respiratory, foodborne, deficiency, Reactive Oxygen Species (ROS) related diseases. Moreover, the antibacterial and anti-arthritis activities indicate the capability of the leaf as a source of natural anti-radical, antibiotic or anti-inflammatory agent with possible

brilliant health benefits for human and animals. Therefore, the results showcase the possible application in the food, pharmaceutical and veterinary industries/sectors.

MATERIALS AND METHODS

Identification of the plant: The medicinal plant investigated was collected Atowode, Ota, Ogun State, Nigeria and it was identified as *Dialium guineense* Wild (*Fabaceae*) at the Department of Biological Science, University of Medical Sciences, Ondo, Nigeria by Dr. Oghale Ovuakporie-Ova.

Preparation of extract: The solvent extraction of the air dried leaves was carried out according to method described by Azwanida^[36] with slight modification. The pulverized sample material was extracted for at least 3 days by maceration using methanol and ethyl acetate (2:1) at room temperature with periodical shaking. After filtration, the solvent was evaporated using Uniscope water bath. The concentrated extract was refrigerated until used. The extraction and dilution procedures were protected from contamination.

Phytochemical analysis using gas chromatography-mass spectrometry: The qualitative and quantitative analysis of the extract was carried out by using GC-MS QP2010 Plus (Shimadzu, Kyoto, Japan) system at the Shimadzu Training Centre for Analytical Instruments (STC) Lagos, Nigeria. The analytical specifications of the GC-MS was used as in our previously study^[37].

Determination of lycopene and β -carotene contents: Evaluation of the lycopene and β -Carotene were quantitatively determined according to the method previously used by Ogunmola *et al.*^[37] and Lakra *et al.*^[38]. It is a simultaneous assay for determination of plant pigments quantitatively. 0.2 g of the fresh leaves sample was homogenized and extracted with 10.0 mL of mixture of 40% acetone and 60% hexane solution. The liquid phase was centrifuged at 4000 x g for 15 min using K-1000 Laboratory Centrifuge. The aliquot was taken from the upper solution (supernatant) and the absorbance were measured at 663, 645, 505 and 453 nm using Uniscope SM 7504 UV Spectrophotometer. The results were expressed as mean \pm SD for triplicate values. Lycopene and β -carotene contents in mgg^{-1} fw were determined using the following expression:

$$\text{Lycopene} = -0.0458A_{663} + 0.204A_{645} + 0.372A_{505} - 0.0806A_{453}$$

$$\beta\text{-Carotene} = 0.216A_{663} - 1.22A_{645} - 0.304A_{505} + 0.452A_{453}$$

where, A absorbance recorded at specific wavelengths.

Total chlorophyll and carotenoid: Total chlorophyll and carotenoid were quantitatively determined according to the assay previously used by Silva-Beltran *et al.*^[39]. Briefly, 0.20 g of the fresh leaf sample was extracted thrice using 80% acetone:water (ratio 4:1) till the green pigments absolutely extracted and then filtered. The upper layer of the aliquot for the three extractions up were added together and the total volume was increased to 25.0 mL. The absorbance was measured at 663, 652, 645 and 470 nm using UniscopeSM 7504 UV Spectrophotometer. The chlorophyll contents were determined as mg g^{-1} of extract (ge) based on the following expression:

$$\begin{aligned} \text{Chlorophyll a} &= (12.7A_{663} - 2.7A_{645}) \\ \text{Chlorophyll b} &= (22.9A_{645} - 2.7A_{663}) \\ \text{Total Chlorophyll a} &= (27.8A_{652}) \\ \text{Total chlorophyll} &= \frac{(1000A_{470} - 1.63 \text{ Chlorophyll a})}{221} \times 104.96 \text{ Chlorophyll b} \end{aligned}$$

Evaluation of Polyphenol Content (PC): The quantitative amount of phenolic compounds was evaluated using Folin-Ciocalteu reagent as described by Nahla *et al.*^[40] using Gallic acid was used as standard for the preparation the calibration curve. 1000 $\mu\text{g mL}^{-1}$ of the extract solution was added to 1.0 mL Folin-Ciocalteu reagent (1:10 v/v in distilled H_2O), 4 mL of 7.5% Na_2CO_3 was added to the mixture after 3 min in order to neutralize the solution. The mixture was allowed to incubate at room temperature for 3 hrs with periodical shaking; the absorbance at 760 nm was measured using UniscopeSM 7504 UV Spectrophotometer. The readings were done in triplicate. The amount of phenolic compounds was expressed in term of Gallic Acid Equivalent (GAE) in μg per mg of extract^[41].

Evaluation of total flavonoid content: The quantitative amount of flavonoid compounds of the extract was evaluated using AlCl_3 solution as described in the previous method used by Shafii *et al.*^[42]. Briefly, 0.10 mL AlCl_3 (1:10 v/v in distilled H_2O), 0.10 mL of CH_3COONa (1 M) and 2.80 mL of distilled H_2O was added to 1.0 mL of the extract. The absorbance was read at 415 nm using UniscopeSM 7504 UV Spectrophotometer after 40 min of incubation at room temperature. To evaluate the total amount of flavonoids in the extract, a calibrated curve was prepared using quercetin as standard. The readings were done in triplicate. The total amount of flavonoid compounds was expressed in μg Quercetin Equivalent (QE) per mg of extract^[43].

Evaluation of tannin content: The tannin content of the sample was evaluated using the iron(III) chloride/gelatin test described by Alsiede *et al.*^[44]. 0.10 g of the air dried

leaves was boiled in 50 mL distilled H_2O for 30 min. 500 mL of distilled H_2O was then added to the filtrate of the solution. 0.5 mL of the aliquot was added to 1 mL 1% $\text{K}_3\text{Fe}(\text{CN})_6$ along with 1 mL of 1% FeCl_3 and volume of the mixture was adjusted to 10 mL by adding deionized water. The absorbance of solution was measured at 720 nm using UniscopeSM 7504 UV Spectrophotometer after 5 min a calibrated curve was prepared using tannic acid as standard in order to evaluate the amount of tannin in the extract. The total amount of tannin compounds in the leaves was expressed in μg Tannic Acid Equivalent (TAE) per mg ^[45].

Evaluation of total ascorbic acid: 1.0 mL 2,4-dinitrophenylhydrazine solution was added to 0.1 mL of the leaf extract (1000 $\mu\text{g mL}^{-1}$). The absorbance of the solution was taken at 515 nm using UniscopeSM 7504 UV Spectrophotometer after the solution was incubated in the dark for 30 min. Ascorbic acid was used as a standard for calibration curve; result was expressed in mg per g of ascorbic acid equivalent^[46].

Antioxidant activities; evaluation of *in vitro* galvinoxyl antioxidant potential: The *in vitro* Galvinoxyl antioxidant activity was carried out according to the method previously described by Amira *et al.*^[47] with slight modification. Briefly, 1.0 mL of various concentrations (62.5-1000 $\mu\text{g mL}^{-1}$) of the extract was added to 1.0 mL of 0.1 mM Galvinoxyl in methanol. The absorbance of the solutions was measured using UniscopeSM 7504 UV Spectrophotometer at 429 nm after 3 h of incubation in the dark. Ascorbic acid was used as positive control. The percentage of radical scavenging was calculated using the expression:

$$\% \text{ Galvinoxyl} = \frac{A_{\text{blank}} - A_{\text{ext}}}{A_{\text{blank}}} \times 100$$

Where:

A_{blank} = Equal the absorbency of blank solution (Galvinoxyl in methanol)

A_{ext} = Equal the absorbance of the extract solution

The dose-response curve was plotted and IC_{50} value for the leaf extract and the reference compound were determined.

Evaluation of *in vitro* DPPH antioxidant activity: DPPH antiradical potential of the extract was determined as described by Foe *et al.*^[48] with slight modification; this was based on the ability of the leaf extract solution to neutralize DPPH radical. Different concentrations of the test sample (1000-62.5 $\mu\text{g mL}^{-1}$) were allowed to react with 0.004% DPPH in methanol for 30 min for proper incubation in the dark. The absorbance was measured at 517 nm using UniscopeSM 7504 UV Spectrophotometer. Percentage radical scavenging activity of the sample was

determined using ascorbic acid as positive control. The antioxidant ability of the extract was calculated using the following expression:

$$1\% \text{ DPPH} = \frac{A_{\text{blank}} - A_{\text{ext}}}{A_{\text{blank}}} \times 100$$

Where:

A_{blank} = Equal the absorbance of blank solutionie (DPPH in methanol)

A_{ext} = Equal the absorbance of the extract solution

The dose-response curve was plotted and IC_{50} value for the leaf extract and the reference compound were determined.

Antioxidant activity index: The Antioxidant Activity Index (AAI) was calculated as:

$$AAI = \frac{\text{DPPH}^{\square} \text{ Initial Concentration}}{IC_{50}}$$

AAI was classified as weak when $AAI < 0.5$, moderate, when AAI ranged between 0.5-1.0, strong, when AAI ranged between 1.0-2.0 and very strong when $AAI > 2.0$ ^[48].

Phosphomolybdate total antioxidant capacity: The total antioxidant of the extract solution was investigated by phosphomolybdate method described by Afsar *et al.*^[49] with slight modification. 1 mL of the extract solution ($1000 \mu\text{g mL}^{-1}$) in methanol was combined a with mixture of 1.0 mL of reagent solution (0.6 M Sulfuric acid, 0.028 M sodium phosphate and 0.004 M ammonium molybdate). The mixture was incubated for 1½ hr at 95 °C in a water bath. The absorbance of the solution was measured at 695 nm using UniscopeSM 7504 UV Spectrophotometer after cooling at room temperature. The antioxidant capacity was expressed mg per g of ascorbic acidequivalent.

Anti-arthritis protein denaturation: *In vitro* anti-arthritis inhibition of protein denaturation was evaluated by the method previously used by Alrasheid *et al.*^[50] with slight modification using fresh egg albumin. 5.0 mL of reaction mixture comprised of egg albumin (0.2 mL) plus phosphate buffer at pH of 6.4 (2.8 mL) and varying concentration of extract (2.0 mL each). Distilled H₂O of the same volume served as control. The solution was incubated at 37°C in a Bio-Oxygen Demand (BOD) laboratory incubator for about 15 min and then heated in water bath at 70°C for 5 min. The absorbance was measured at 660 nm after cooling at room temperature. Diclofenac sodium (100 mg) was used as reference drug. The percentage anti-arthritis potential was evaluated using the following expression:

$$I\% = 100 \times \frac{V_e}{V_c} - 1$$

Where:

V_e = The absorbance of leaf extract

V_c = The absorbance of the control

Evaluation of *in vitro* antibacterial properties: The antibacterial property of varying concentrations of the leaf extract was investigated by Agar-well diffusion method as previously described by Ogunmola *et al.*^[37]; Yunana *et al.*^[5] with slight modification. Fifteen clinical isolates made up of six Gram positive and nine Gram negative bacteria were employed in the antibacterial test. The Gram-positive bacteria were: *Bacillus spp*, *Enterococcus faecalis*, *Micrococcus varians*, *Streptococcus agalactiae* and *Staphylococcus aureus* and *Staphylococcus saprophyticus* while the Gram-negative bacteria were: *Acinetobacter spp*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Providencia stuartii*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Serratia marcescens* and *Shigella dysenteriae*.

Antibacterial Activity Index (AI): The activity index of the extract was determined using the formula below:

$$\text{Antibacterial Index (AI)} = \frac{\text{Zone of inhibition of Extract against each Bacteria}}{\text{Zone of the Reference Drug against each Bacteria}}$$

The Relative Percentage Inhibition (RPI): The Relative Percentage Inhibition (RPI) of the test extract with respect to the positive control was calculated by using the method described by Hepsibah and Jothi^[52]:

$$RPI\% = \frac{I_{\text{ext}} - I_{\text{solv}}}{I_{\text{std}} - I_{\text{solv}}} \times 100$$

Where:

I_{ext} = Total area of inhibition of the test extract

I_{solv} = Total area of inhibition of the solvent

I_{std} = Total area of inhibition of the standard drug

The total area of the inhibition was calculated by using $\text{area} = \pi r^2$. r = radius of the zone of inhibition.

RESULTS AND DISCUSSION

Chemical constituent of the leaf extract of *D. guineense*: From the GC-MS phytochemical analysis, (37) pharmacologically active secondary metabolites were identified in the leaf methanolic/ethylacetate extract

of *D. guineense*, accounting for 99.2% of the total chemical composition of the leaf extract (Table 1) and the most abundant constituents identified were: 4-O-methylmannose (49.0%), D-(-)-quinic acid (7.0%), methyl 8-[2-((2-[(2-ethylcyclopropyl)-methyl]-cyclopropyl)methyl)-cyclopropyl]-octanoate (6.7%), 3-O-methyl-d-glucose (6.0%), methyl- α -d-mannofuranoside (6.0%) and palmitic acid (5.8%). Most of the metabolites in the leaf extract belong to the family of saccharides. Some phenolic compounds such as p-vinylguaiacol, 3',5'-dimethoxyacetophenone and methyl salicylate were also present in a trace amounts in the leaf extract of the plant studied. Comparatively, the previously study done by Osman *et al.*^[53] on the characterization of the seed and endocarp of the fruit of *D. indum* from Malaysia for their chemical composition using GC-MS showed that a total of thirty-eight (38) metabolites were detected in derivatized exocarp dichloromethane (DCM) fraction and seed methanol fraction (SMF). The principal secondary metabolites identified in the seed methanol fraction

(SMF) were: sucrose (18.86%), β -D-glucopyranose (3.89%) and β -D-galactofuranose (2.43%) while the most abundant phytochemical identified in the *D. indum* derivatized exocarp dichloromethane (DCM) fraction were: palmitic acid (9.71%), oleic acid (7.17%) and vanillin (3.48%) (Osman *et al.*, 2018). It was also reported that some other classes of phytochemicals such as derivatives of phenolic compounds and amino acids were also detected in the analysis. Additionally, the Seed Methanol Fraction (SMF) still contained some other derivatives of amino acids, saccharides, polyol and sesquiterpene which were not present in the derivatized exocarp dichloromethane (DCM) fraction. Surprisingly, some of the therapeutically active organic compounds detected in our work on *D. guineense* grown in Nigeria were also present in the sample of *D. indum* from Malaysia as reported by Osman *et al.*^[53].

β -carotene and lycopene contents: The quantitative amount of β -carotene and lycopene contents of the leaf extract were 0.2 and 0.1 mgg⁻¹, respectively (Table 2).

Table 1: Phytochemical composition of leaf extract of *D. guineense*

Compound	Retention index	Composition (%)
Ethyl-2,3-dimethylbutanoate	856	0.3
1-ethynyl-4-fluorobenzene	866	0.1
Coumaran	1036	0.1
E-3-methyl-2-hexenoic acid	1059	0.1
Thymine	1118	0.1
5-hydroxymethylfurfuran	1163	0.2
Ethyl linalool	1181	0.1
N-(3-butenyl)-N-methylcyclohexanamine	1217	0.1
1-(octylsulfanyl)ethylene	1256	0.3
3,5-dihydroxy-6-methyl-2,3-dihydro-4H-pyran-4-one	1269	0.6
Dimethyl d-tartrate	1277	0.7
Methyl salicylate	1281	0.2
p-vinylguaiacol	1293	0.7
n-capric acid	1372	0.1
3',5'-dimethoxyacetophenone (phenolic)	1407	1.0
4-methylpentylcyclohexanecarboxylate	1480	0.2
3,4-diketo- α -methylglutaric acid	1538	0.1
3-O-methyl-d-glucose	1647	6.0
Methyl- α -d-mannofuranoside	1667	6.0
5-dimethylsilyloxytetradecane	1677	0.2
4-O-methylmannose	1714	49.0
D-(-)-quinic acid	1852	7.0
n-hexadecan-1-ol	1854	0.1
Palmitic acid, methyl ester	1878	3.6
Methyl-(7E)-7-hexadecenoate	1886	0.1
n-heptadecanol-1	1954	0.1
Palmitic acid	1968	5.8
Trans-phytol	2045	0.2
Methyl stearate	2077	1.0
Oleic acid, methyl ester	2085	0.1
Methyl linolenate	2101	4.5
Methyl petroselinate	2104	0.1
14-hydroxy-15-methyl-15-hexadecenoic acid	2197	0.1
Methyl 8-[2-((2-[(2-ethylcyclopropyl)methyl]cyclopropyl)methyl)cyclopropyl]octanoate	2266	6.7
Z,Z,Z-9,12,15-octadecatrienoic acid, 2-[(trimethylsilyloxy)-1-[[trimethylsilyloxy]methyl]ethyl ester	2804	1.0
Farnesyl- β -D-mannofuranoside	3014	1.6
Adenosine, N6-phenylacetic acid	3731	0.3
Percentage total		98.5

Table 2: Pigment and their contents in the leaf extract of *D. guineense*

Pigment	Content (mgg ⁻¹)
Lycopene	0.1
β-Carotene	0.2
Total carotenoid	12.68
Chlorophyll a	4.4
Chlorophyll b	4.7
Total chlorophyll	7.8

This showed that the β-carotene content is twice the lycopene content in the leaf of *D. guineense*. Lycopene has several health benefits such as antioxidative, anti-inflammatory, anti-atherogenic, anti-atherosclerotic, cardioprotective and blood pressure control^[54-60].

Total pigment: Chlorophyll a, Chlorophyll b, Total Chlorophyll and carotenoid contents. Quantitatively, the amount of chlorophyll a, chlorophyll b, and total chlorophyll content in the leaf extract investigated were 4.4, 4.7 and 7.8, respectively (Table 2). It was reported that chlorophyll and its derivatives as fat-soluble compounds play many therapeutic roles such as anti-mutagenic and anti-carcinogenic properties^[61-63]. Plant pigments play important roles in human and animal health. According to Lehto *et al.*^[64]; Sen *et al.*^[65] carotenoids such as lycopene, β-carotene, xanthophylls, lutein, capsanthin, astaxanthin, cantaxanthin, cryptoxanthin and zeaxanthin etc found in natural green vegetables are natural food additives, colouring agents and preservatives. They have key roles playing in maintaining growth, good health, prevention and cure of diseases. They are good sources of natural provitamin A and non-provitamin A^[66-68].

Evaluation of polyphenol content; total phenolic content:

The total amount polyphenols in the leaf extract investigated was 2,668.17 μgmg⁻¹ GAE (Table 2). The presence of some active phenolic secondary metabolite such as p-vinylguaiacol, 3',5'-dimethoxyacetophenone and methyl salicylate in the leaf extract. Comparatively, the phenolic content evaluated in the leaf extract studied was approximately 14 times higher than those reported in other *Fabaceae* family such as the methanolic extract of *D. ovoideum* grown in Sri Lanka with TPC of 192.90 mg PGE/g^[70]. Likewise, Muthukumaran *et al.*^[71] reported that the phenolic content of the leaf and flower ethanolic extracts of *Peltophorum pterocarpum* (*Fabaceae*) from India were 33.17 mg and 24.91 mg of GAE/g, respectively, this is about 89% lower than the phenolic proportion in the leaf extract investigated in this study.

Flavonoid content: The quantitative estimate of the flavonoid content in the leaf extract was 17.80 μgmg⁻¹ (Table 3), this is comparable to the amount of flavonoid in the ethanolic leaf extract of *P. pterocarpum* (*Fabaceae*) with the value of 1.44±0.01 mgQE/g^[71].

Table 3: Polyphenol and TAA of the leaf extract of *D. guineense*

TPC	TFC	TTC	TAA
2,668.17±0.01	17.80±0.00	152.65±0.01	43.75±0.04
μgmg ⁻¹ GAE	μgmg ⁻¹ QE	μgmg ⁻¹ TAE	μgmg ⁻¹ AAE

Data are shown as the mean value±S.D of triplicate

Tannin content: The leaf extract of *D. guineense* studied had high amount of tannin 152.65 μgmg⁻¹ TAE and its derivatives (Table 2). Comparatively, another report showed that the tannin content of ethanolic extract of flower and aqueous methanolic extracts of leaves of *P. pterocarpum* (*Fabaceae*) from India have highest content of tannin of 844±10.38 mg TAE/g and 776.32±35.01 mg TAE/g, respectively^[71]. These classes of polyphenols considered in this study have a lot of potential health benefits; they are good antioxidants that protect human body from heart diseases and cancer by preventing cellular damage^[72]. Lin *et al.*^[73]; Panche *et al.*^[74]; De-Silva and Alcorn^[75] stated that polyphenols are an important class of natural products widely available in green vegetables, fruits and some natural juices. Vinha *et al.*^[76]; Panche *et al.*^[74]; Kozłowska and Szostak-Węgierek^[77]; Forni *et al.*^[78]; Liu *et al.*^[79] reported that polyphenols have the ability to interfere and combat the pathogenesis of degenerative and neurodegenerative diseases such as cancer, cystic fibrosis, parkinson's and huntington's, Alzheimer's Disease (AD), atherosclerosis etc. Some polyphenol have ability to inhibit the replication of HIV while some also lower blood glucose levels and have insulin-like problems^[74, 80, 81].

Ascorbic acid (vitamin c) content: The amount of ascorbic acid in the leaf extract was evaluated as 43.75 μgmg⁻¹ AAE (Table 3). Ascorbic acid is an essential dietary nutrient both at infant, adolescence and adult stages of life. It has series of metabolic, physiological and therapeutic functions. Aghajanian *et al.*^[82] reported that vitamin C is an important phytochemical which is good antioxidant and cofactor playing a key role in the development, regulation, function and maintenance of several cells in human and animals. Ascorbic acid and its derivatives enhance the absorption of iron in the gut by reducing ferric to ferrous state and also aid the conversion of cholesterol into bile acids, in order to reduce the blood cholesterol. It also facilitates tissue and wound healing. Moreover, it is a fundamental in the lipid metabolism, atherogenesis and management of neurodegenerative disorders, cancer, diabetes and male infertility problems^[83]. Moreover, Young *et al.*^[84]; Tan *et al.*^[60] also reported that vitamin C plays a serious role in biosynthesis of collagen by facilitating the hydroxylation of proline and lysine residues, therefore, helps in proper intracellular folding of pro-collagen for export and deposition as mature collagen. Grosso *et al.*^[85]; Aghajanian *et al.*^[82] stated that deficiencies in vitamin C in human and animal lead to symptoms such as scurvy, Gingivitis, bone pain, impaired wound healing etc.

Table 4: Galvinoxyl, DPPH and TAC antioxidant properties of the leaf extract of *D. guineense*

Extract	IC ₅₀ µg mL ⁻¹	AAE µg mg ⁻¹
GALV	125.0	-
DPPH•	6.0	-
TAC	-	279.49±0.01

Anti-radical and antioxidant potential; galvinoxyl activities: The galvinoxyl percentage radical scavenging of by the extract at varying concentrations (1000, 500, 250, 125 and 62.5 µg mL⁻¹) were 77.92, 60.73, 59.13, 25.35 and 20.40%, respectively. The leaf extract of *D. guineense* having IC₅₀ value of 125.0 µg mL⁻¹. This was lower than that of ascorbic acid which had an IC₅₀ value of 15.0 µg mL⁻¹ (Table 4).

DPPH activity: The radical inhibitions of the extract at varying concentrations (1000, 750, 500, 250 and 62.5 µg mL⁻¹) were 91.92, 88.53, 86.22, 85.48 and 85.25% respectively. The methanolic leaf extract of *D. guineense* with IC₅₀ and AAI values of 6.0 µg mL⁻¹ and 7.0, respectively (Table 4). This is similar to the IC₅₀ and AAI of ascorbic acid with value of 9.0 µg mL⁻¹ and 4.4, respectively. This study showed that the leaf extract investigated in this study had more anti-radical potential than other plant sample in the family such as the seed crude methanolic extract of *D. indium* from Malaysia with DPPH percentage free radical scavenging and IC₅₀ values of 90.99±0.03% and 99.95±0.98 µg mL⁻¹, respectively^[53]. In the same manner, methanolic extract of *D. ovoidium* from Sri Lanka had DPPH percentage antiradical activity of 92.90±0.4%^[70].

Total antioxidant capacity: The evaluated amount of total antiradical capacity of leaf extract of *D. guineense* grown in Nigeria was 279.49±0.00 µg mg⁻¹ AAE (Table 4). The result shows that the leaf extract used in this study had more free radical scavenging potential compared to other related species such as the seed crude methanolic extract of *D. indium* from Malaysia with TAC value of 222.72±16.03^[53]. Medicinal plants have significant high phenolic contents and other phytochemicals with high antioxidant properties compared to the conventional antioxidants that in some cases have side effects^[86]. Moreover, medicinal plants present abundant sources of natural antioxidants and find enormous application in human nutrition, not only as flavouring spices but also as natural remedy. Datta *et al.*^[87]; Da-Oliveira *et al.*^[88] stated that food enrichment and fortification with natural products such as polyphenols from naturally grown vegetables is a new trend to improve the nutritional and therapeutic values of food at home and industries. Plants are rich in minerals, vitamins, flavoring agents and natural antioxidants. It was reported that natural product from medicinal plants added to food substances increased the total phenolic and

Table 5: Anti-arthritis activity (Egg albumin assay) of the leaf extract of *D. guineense*

Concentration µg mL ⁻¹	Inhibition (%)	EC ₅₀ µg mL ⁻¹
1000	96.00±0.00	
500	94.19±0.01	
250	90.05±0.00	5.00
125	88.76±0.00	
62.5	83.33±0.00	

Data are presented as the mean value±S.D. of triplicate

antioxidant values of raw material and final food product^[89, 90]. Some of these medicinal plants have been shown to have both chemo-preventive or therapeutic potential on breast and other form of cancer^[91-92].

Anti-arthritis and anti-inflammatory potential: The anti-arthritis/anti-inflammatory efficacy of the investigated leaf extract was studied using egg albumin protein denaturation method. The extract of *D. guineense* showed high percentage inhibition of anti-arthritis activities ranging between 83.33-96.00% as compared to the Diclofenac sodium (reference drug) with percentage inhibition of 80.00% (Table 5). In support of the research on the mechanism of the anti-arthritis activity of natural products from plants, ability of the leaf sample used in this study to inhibit arthritis and inflammation was investigated. Rheumatoid arthritis is joint-related, immune-based inflammatory disorder that affects almost 1% of the world's population^[93-96]. Fu *et al.*^[97]; Rahman *et al.*^[98]; Walling and Kim^[99] reported that pathogenesis is observed in the T-cell activation that leads to clusters of differentiation in the T-cells and concurrently increases various inflammatory indicators. These include matrix metalloproteinase, cytokines, chemokines, NF-κβ, adhesion molecules, lipoxigenase, nitric oxide and arachidonic acid. They destroy bone and cartilage by changing the structure and shape which lead to degradation. Based on this, incorporation of phytochemicals such as polyphenols, terpenoids and other secondary metabolites have high efficacy against arthritis via targeted action on inflammatory biomarkers^[100-102]. Obaseki *et al.*^[103]; Sangeetha and Vidhya^[104] stated that different anti-arthritis and anti-inflammatory drugs have shown dose dependent ability to inhibit thermally induced protein denaturation. In this study, the extract of *D. guineense* was able to significantly inhibited heat induced protein denaturation and therefore, shows strong anti-arthritis activity.

Antibacterial activities: The effects of the leaf extract of *D. guineense* at a 1000-250 µg mL⁻¹ concentration on the growth of the 15 clinically isolated bacteria strains tested are presented in Table 6. The zones of inhibition of the leaf extract (20.0-08.0 mm) showed high antimicrobial activities as compared to gentamicin antibiotic (10 µg per disc). Greater or similar activities were observed as concentration of the extract 1000-250 µg mL⁻¹ by the

Table 6: Antibacterial potential (Zones of Inhibition in mm) of the leaf extract of *D. guineense*

Conc. ($\mu\text{g mL}^{-1}$)/ Organisms	ZI of the leaf extract			Gentamicin		
	1000	500	250	10 μg	AI	RPI
<i>Acinetobacter</i> spp (-)	10	10	10	-	10	10.00
<i>Bacillus</i> spp (+)	15	10	10	20	0.8	56.23
<i>E. coli</i> (-)	20	20	08	15	1.3	177.85
<i>E. faecalis</i> (+)	15	11	10	14	1.1	114.80
<i>K. pneumoniae</i> (-)	16	12	10	15	1.1	113.80
<i>M. varians</i> (+)	15	-	-	19	0.8	63.33
<i>P. aeruginosa</i> (-)	10	08	08	20	0.5	25.00
<i>P. mirabilis</i> (-)	16	16	10	18	0.9	79.02
<i>P. stuartii</i> (-)	14	14	-	16	0.9	76.55
<i>S. agalactiae</i> (+)	10	10	10	20	0.5	25.00
<i>S. aureus</i> (+)	20	20	10	20	1.0	100.00
<i>S. dysenteriae</i> (-)	10	10	10	20	0.5	25.00
<i>S. marcescens</i> (-)	15	15	14	17	0.9	77.85
<i>S. typhimurium</i> (-)	15	-	-	20	0.8	56.23
<i>S. saprophyticus</i> (+)	20	08	08	17	1.2	138.50

Resistant (-), not sensitive (<8 mm), sensitive (9-14 mm), very sensitive (15-19 mm) and ultrasensitive (>20 mm)

extract than gentamicin (standard antibiotic). The Activity Index (A.I) was greater than or equal to 1.0 against the bacteria isolates. No significant reduction in activities was observed as the extract concentrations were reduced gradually from 1000-250 $\mu\text{g mL}^{-1}$. Comparatively, the extract investigated in this study had a similar antibacterial potential as reported by Ijoma *et al.*^[105] on their study on *Dialium indum* leaves fractions with average diameter zones of inhibition ranging between 14-36 mm on the following bacteria *B. typhi*, *E. aerogenes*, *E. coli*, *K. pneumonia*, *P. aeruginosa*, *P. vulgaris*, *S. typhi*, *S. albus*, *S. aureus*, *S. muteus*. Moreover, previous work on the extract of a related species such as the ethanol extract of the aerial (leaf and bark) of *Butea monosperma* (*Fabaceae*) grown in India was found to be only effective against *B. subtilis* and *S. aureus* (13 mm at 100 mg mL^{-1}) among the bacteria isolates used, the petroleum ether extract did not inhibit *P. aeruginosa* while *E. coli* was not inhibited by any type of extract^[106]. From the results obtained in this study, the leaf extract exhibited a notable and promising antimicrobial inhibitory activity against all the tested pathogenic bacteria. In fact, the presence of various polyphenols such as p-vinylguaiaicol, 3',5'-dimethoxyacetophenone, methyl salicylate etc in the leaf studied contributed tremendously to the antibacterial activities. In addition, synergistic activity of all secondary metabolite in the extract also played central roles to the antibacterial activities observed in this study^[107, 108]. Therefore, the leaves of the plant can be used in developing antibacterial drug in combating multidrug resistant bacteria.

CONCLUSION

The study demonstrated the therapeutic potential of *D. guineense* grown in Nigeria. The medicinal properties

of the leaf may be related to its polyphenols and other active metabolites. This study indicated that the leaf of the plant can be used for development of important drugs for human and animals. This observation is very significant because of the possibility of developing therapeutic drug that will be active against multi-drug resistant organisms and easily accessible natural antioxidant substances that may help to modulate oxidative stress related disorders. Further study is therefore recommended on the pharmacology properties of the leaves of the plant which are not covered by this study.

REFERENCES

- Kiringe, J.W., 2005. Ecological and anthropological threats to ethno-medicinal plant resources and their utilization in Maasai communal ranches in the Amboseli region of Kenya. *Ethnobot. Res. Applic.*, 3: 231-241.
- Falodun, A., 2010. Herbal medicine in africa-distribution, standardization and prospects. *Res. J. Phytochem.*, 4: 154-161.
- Mahomoodally, M.F., 2013. Traditional medicines in Africa: An appraisal of ten potent African medicinal plants. *Evidence-Based Complement. Altern. Med.*, 2013. 10.1155/2013/617459
- Lawal, B., O.K. Shittu, F.I. Oibiokpa, E.B. Berinyuy and H. Mohammed, 2017. African natural products with potential antioxidants and hepatoprotectives properties: A review. *Clin. Phytosci.*, 2: 1-66.
- Genet, A., 2018. Indigenous herbal medicinal knowledge among the Shinasha. *Global Sci. J.*, 6: 63-83.
- Altemimi, A., N. Lakhssassi, A. Baharlouei, D.G. Watson and D.A. Lightfoot, 2017. Phytochemicals: Extraction, isolation and identification of bioactive compounds from plant extracts. *Plants*, Vol. 6, No. 4. 10.3390/plants6040042
- Ekeopara, C.A. and A.M. Ugoha, 2017. The contributions of African traditional medicine to Nigeria's health care delivery system. *J. Hum. Social Sci.*, 22: 32-43.
- Abebe, D. and B. Garedew, 2019. Ethnobotanical survey of plants traditionally used for malaria prevention and treatment in indigenous villages of Tepi town South West Ethiopia. *J. Pharmacogn. Phytother.*, 11: 9-16.
- Shah, M., Y.R. Behara and B. Jagadeesh, 2012. Phytochemical screening and *in vitro* antioxidant activity of aqueous and hydroalcoholic extract of *Bacopa monnieri* linn. *Int. J. Pharma. Sci. Res.*, 3: 3418-3424.
- Karunamoorthi, K., K. Jegajeevanram, J. Vijayalakshmi and E. Mengistie, 2013. Traditional medicinal plants: A source of phytotherapeutic modality in resource-constrained health care settings. *J. Evidence-Based Complementary Altern. Med.*, 18: 67-74.

11. Pan, S.Y., S.F. Zhou, S.H. Gao, Z.L. Yu and S.F. Zhang *et al.*, 2013. New perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics. Evidence-Based Complementary Alternative Med., 2013: 1-25.
12. Kutama, R.M., R.S. Adbulkadi, S.A. Kwalli and G. Chiroma, 2018. Phytochemical compositions in some Nigerian medicinal plants and their pharma-cological properties: A review. J. Anesthesiology, 6: 15-25.
13. Kumar, S., S. Paul, Y.K. Walia, A. Kumar and P. Singhal, 2015. Therapeutic potential of medicinal plants: A review. J. Biol. Chem. Chron., 1: 46-54.
14. Xu, D.P., Y. Li, X. Meng, T. Zhou, Y. Zhou, J. Zheng, J.J. Zhang and H.B. Li, 2017. Natural antioxidants in foods and medicinal plants: Extraction, assessment and resources. Int. J. Mol. Sci., 18: 1-32.
15. Taghvaei, M. and S.M. Jafari, 2015. Application and stability of natural antioxidants in edible oils in order to substitute synthetic additives. J. Food Sci. Technol., 52: 1272-1282.
16. Ohwofasa, I.K. and J.O. Pon, 2018. Synergistic effect of the extracts of *Vernonia amygdalina* and *Solenostemon monostachyus* on Gram-negative bacteria. Pak. J. Biol. Sci., 21: 369-375.
17. Masarkar, N., S. Mukherjee, S.K. Goel and R. Nema, 2019. Naturally Derived formulations and prospects towards cancer. Health, 11: 971-997.
18. Asif, M., 2015. Chemistry and antioxidant activity of plants containing some phenolic compounds. Chem. Int., 1: 35-52.
19. Zhang, Y.J., R.Y. Gan, S. Li, Y. Zhou, A.N. Li, D.P. Xu and H.B. Li, 2015. Antioxidant phytochemicals for the prevention and treatment of chronic diseases. Molecules, 20: 21138-21156.
20. Huyut, Z., S. Beydemir and I. Gülçin, 2017. Antioxidant and antiradical properties of selected flavonoids and phenolic compounds. Biochem. Res. Int., Vol. 2017. 10.1155/2017/7616791
21. Kozarski, M., A. Klaus, D. Jakovljevic, N. Todorovic and J. Vunduk *et al.*, 2015. Antioxidants of edible mushrooms. Molecules, 20: 19489-19525.
22. Yuan, H., Q. Ma, L. Ye and G. Piao, 2016. The traditional medicine and modern medicine from natural products. Molecules, Vol. 21. 10.3390/molecules21050559
23. Alizadeh, M., A. Safarzadeh, M. Bahmani, F. Beyranvand and M. Mohammadi *et al.*, 2018. Brucellosis: Pathophysiology and new promising treatments with medicinal plants and natural antioxidants. Asian Pac. J. Trop. Med., 11: 597-608.
24. Okudu, H.O., E.J. Umoh, M.C. Ojinnaka and O.F. Chianakwalam, 2017. Nutritional, functional and sensory attributes of jam from velvet tamarind pulp. Afr. J. Food Sci., 11: 44-49.
25. Gnansounou, M.S., S. Iskandar, M. Robin, J.M. Brunel, E. Dahouenon and P. Piccerelle, 2018. *Dialium guineense* Willd. *Parkia biglobosa* (Jacq.) R. Br. Ex Benth. and *Tamarindus indica* L.: Review of known and synergetic bioactive compounds. J. Med. Plants Stud., 6: 103-111.
26. Ogu, G.I., J. Ezeadila and J.M. Ehiobu, 2013. Antioxidant and antimicrobial activities of *Dialium guineense* (Willd) leaf extract. Pharm. Pharmacol. Res., 1: 1-7.
27. Osanaiye, F.G., M.A. Alabi, R.M. Sunday, T. Olowokere, E.T. Salami, T.A. Otunla and S.C. Odiaka, 2013. Proximate composition of whole seeds and pulp of african black velvet tamarind (*Dialium guineense*). IOSR J. Agric. Vet. Sci., 5: 49-52.
28. Asoiro, F.U., S.L. Ezeoha, G.I. Ezenne and C.B. Ugwu, 2017. Chemical and mechanical properties of velvet tamarind fruit (*Dialium guineense*). Niger. J. Technol., 36: 252-260.
29. Davies R.M. and D.D. Yusuf, 2017. Studies of physical and mechanical properties of velvet tamarind. Mayfeb J. Agric. Sci., 2: 36-43.
30. Utubaku, A.B., O.E. Yakubu and D.U. Okwara, 2017. Comparative phytochemical analysis of fermented and unfermented seeds of *Dialium giuneense*. J. Tradit. Med. Clin. Natur., Vol. 6.
31. Eze, E.M., P.I. Orjiakor, J.C. Okolo and G.I. Ogu, 2018. *In vitro* antibacterial efficacy of velvet tamarind (*Dialium guineense* Willd) root extract. Int. J. Adv. Acad. Res., 4: 45-58.
32. Besong, E.E., M.E. Balogun, S.F.A. Djobissie, D.C. Obu and J.N. Obimma, 2016. Medicinal and economic value of *Dialium guineense*. Afr. J. Biomed. Res., 19: 163-170.
33. Lokonon, B.E., W.N. Bonou, B. Kassa, A.F. Azihou, E. Assogbadjo and R. Glelekakai, 2013. Structural and ethnobotanical characterization of velvet tamarind (*Dialium guineense* willd), a multipurpose tree species. Agronomie Afr., 25: 121-131.
34. Ajayi, A.I., O.V. Adeniyi and W. Akpan, 2018. Use of traditional and modern contraceptives among childbearing women: Findings from a mixed methods study in two South Western Nigerian states. BMC Public Health, 18: 1-9.
35. Nkanu, E.E., G. Ujong, V. Okon and I.E. Bassey, 2018. *In vivo* efficacy of *Dialium guineense* fruit pulp on hemeoxygenase-1 and angiotensin converting enzyme in experimental diabetes. J. Med. Plants Res., 12: 483-492.
36. Azwanida, N.N., 2015. A review on the extraction methods use in medicinal plants, principle, strength and limitation. Med. Aromat. Plants, Vol. 4. 10.4172/2167-0412.1000196
37. Ogunmola, O.O., S.E. Kuyooro and O.O. Abiona, 2017. *Stachytarpheta jamaicensis* leaf extract: Chemical composition, antioxidant, anti-arthritic, anti-inflammatory and bactericidal potentials. J. Sci. Innovative Res., 6: 119-125.

38. Lakra, A., J. Trivedi and S. Mishra, 2018. Studies on biochemical composition of various tomato (*Solanum lycopersicum* L.) genotypes. Int. J. Curr. Microb. Applied Sci., 7: 977-987.
39. Silva-Beltran, N.P., S. Ruiz-Cruz, L.A. Cira-Chavez, M.I. Estrada-Alvarado and J.J. Ornelas-Paz *et al.*, 2015. Total phenolic, flavonoid, tomatine and tomatidine contents and antioxidant and antimicrobial activities of extracts of tomato plant. Int. J. Anal. Chem., 15: 1-10.
40. Nahla, T.K., S.U. Wisam and N.M. Tariq, 2018. Antioxidant activities of beetroot (*Beta vulgaris* L.) extracts. Pak. J. Nutr., 17: 500-505.
41. Achouri, N., N. Smichi, N. Kharrat, F. Rmili, Y. Gargouri, N. Miled and A. Fendri, 2015. Characterization of liver oils from three species of sharks collected in Tunisian coasts: *In vitro* digestibility by pancreatic lipase. J. Food Biochem., Vol. 42, 10.1111/jfbc.12453
42. Shafii, Z.A., M. Basri, E.A. Malek and M. Ismail, 2017. Phytochemical and antioxidant properties of *Manilkara zapota* (L.) P roen fruit extracts and its formulations for cosmeuetical application. Asian J. Plant Sci. Res., 7: 29-41.
43. Orak, H.H., H. Yagar and S.S. Isbilir, 2012. Comparison of antioxidant activities of juice, peel, and seed of pomegranate (*Punica granatum* L.) and inter-relationships with total phenolic, Tannin, anthocyanin and flavonoid contents. Food Sci. Biotechnol., 21: 373-387.
44. Alsiede, M.M.S.A., M.A. Abdrahman and A.E.M. Saeed, 2015. Phytochemical screening, total phenolics content and antioxidants activity of cassia singueana. J. Med. Plants, 3: 160-165.
45. Ahmed, S., F.B. Rahman, S. Ahmed and A.M. Shohaël, 2018. Insights into the bioactive compounds, antioxidant potential and TLC profiling of different extracts of Tomato plants. Jahangirnagar Univ. J. Biol. Sci., 7: 65-77.
46. Benites, R.S.R., A.S.N. Formagio, E.J.S. Argandona, C.R.F. Volobuff, L.N.F. Trevizan, M.C. Vieira and M.S. Silva, 2015. Contents of constituents and antioxidant activity of seed and pulp extracts of *Annona coriacea* and *Annona sylvatica*. Braz. J. Biol., 75: 685-691.
47. Amira, S., M. Dade, G. Schinella and J.L. Rios, 2012. Anti-inflammatory, anti-oxidant and apoptotic activities of four plant species used in folk medicine in the *Mediterranean basin*. Pak. J. Pharm. Sci., 25: 65-72.
48. Foe, F.M.C.N., T.F.K. Tchingang, A.M. Nyegue, J.P. Abdou and A.J. Yaya *et al.*, 2016. Chemical composition, *in vitro* antioxidant and anti-inflammatory properties of essential oils of four dietary and medicinal plants from Cameroon. BMC Complementary Altern. Med., 16: 1-12.
49. Afsar, T., S. Razak, M. Shabbir and M.R. Khan, 2018. Antioxidant activity of polyphenolic compounds isolated from ethyl-acetate fraction of *Acacia hydaspica* R. Parker. Chem. Central J., Vol. 12. 10.1186/s13065-018-0373-x
50. Alrasheid, A.A., A.A. Mohamed, E.G. Mohieldin, K.I. Eldein and L.F. Yassin *et al.*, 2019. Phytochemical investigation and assessment of antimicrobial, anti-inflammatory and antioxidant activities of Sudanese *Citrus paradise* peel extract. J. Pharmacog. Phytother., 11: 1-8.
51. Yunana, B.T., J.C. Guiyi and B.B. Bakar, 2018. *In vitro* and *in vivo* evaluation of antibacterial activity of *Bridelia ferrugine* extracts on some clinical isolates. J. Phytopharmacology, 7: 392-398.
52. Hepsibah, A.H. and G.J. Jothi, 2016. A comparative study on the effect of solvents on the phytochemical profile and biological potential of *Ormocarpum cochinchinense* Auct. Non (Lour.) Merrill. Int. J. Pharm. Pharm. Sci., 9: 67-72.
53. Osman, M.F., M.N. Hassan, A. Khatib and S.M. Tolos, 2018. Antioxidant activities of *Dialium indum* L. fruit and Gas Chromatography-Mass Spectrometry (GC-MS) of the active fractions. Antioxidants, Vol. 7, No. 11. 10.3390/antiox7110154
54. Sahabi, D.M., R.A. Shehu, Y. Saidu and A.S. Abdullahi, 2012. Screening for total carotenoids and β -carotene in some widely consumed vegetables in Nigeria. Nigerian J. Basic Applied Sci., 20: 225-227.
55. Nimse, S.B. and D. Pal, 2015. Free radicals, natural antioxidants and their reaction mechanisms. RSC Adv., 5: 27986-28006.
56. Upadhyay, S. and M. Dixit, 2015. Role of polyphenols and other phytochemicals on molecular signaling. Oxid. Med. Cell. Longev., Vol. 2015. 10.1155/2015/504253
57. Shastri, A., R. Srivastava, B. Jyoti and M. Gupta, 2016. The antioxidants-scavengers of free radicals for immunity boosting and human health/overall well being. Int. J. Contemp. Med. Res., 3: 2918-2923.
58. Casas, R., S. Castro-Barquero, R. Estruch and E. Sacanella, 2018. Nutrition and cardiovascular health. Int. J. Mol. Sci., Vol. 19, 10.3390/ijms19123988
59. Mozos, I., D. Stoian, A. Caraba, C. Malainer, J.O. Horbanczuk and A.G. Atanasov, 2018. Lycopene and vascular health. Front. Pharmacol., Vol. 9, 10.3389/fphar.2018.00521
60. Tan, B.L., M.E. Norhaizan, W.P.P. Liew and H.S. Rahman, 2018. Antioxidant and oxidative stress: A mutual interplay in age-related diseases. Front. Pharmacol., Vol. 9. 10.3389/fphar.2018.01162
61. Kumar, G., P. Sajeera and K.K. Sivadasan, 2017. A quantitative study on the selected phytochemicals of six indigenous medicinal plants of Mahe, UT of Puducherry, India. Int. J. Herbal Med., 5: 1-4.

62. Vankova, K., I. Markova, J. Jasprova, A. Dvorak and I. Subhanova *et al.*, 2018. Chlorophyll-mediated changes in the redox status of pancreatic cancer cells are associated with its anticancer effects. *Oxidative Med. Cell. Longevity*, Vol. 2018, 10.1155/2018/4069167
63. Zeng, Y., X. Pu, J. Yang, J. Du and X. Yang *et al.*, 2018. Preventive and therapeutic role of functional ingredients of barley grass for chronic diseases in human beings. *Oxid. Med. Cell. longevity*, Vol. 2018, 10.1155/2018/3232080
64. Lehto, S., M. Buchweitz, A. Klimm, R. Strabburger, C. Bechtold and F. Ulberth, 2017. Comparison of food colour regulations in the EU and the US: A review of current provisions. *Food Additives Contaminants Part A*, 34: 335-355.
65. Sen, T., C.J. Barrow and S.K. Deshmukh, 2019. Microbial pigments in the food industry—challenges and the way forward. *Front. Nutr.*, 10.3389/fnut.2019.00007
66. Paiva, S.A. and R.M. Russell, 1999. β -carotene and other carotenoids as antioxidants. *J. Am. Coll. Nutr.*, 18: 426-433.
67. Mezzomo, N. and S.R. Ferreira, 2016. Carotenoids functionality, sources and processing by supercritical technology: A review. *J. Chem.*, Vol. 2016, 10.1155/2016/3164312
68. Torregrosa-Crespo, J., Z. Montero, J.L. Fuentes, R.M. Garcia-Galbis, I. Garbayo, C. Vilchez and R.M. Martinez-Espinosa, 2018. Exploring the valuable carotenoids for the large-scale production by marine microorganisms. *Marine Drugs*, Vol. 16, 10.3390/md16060203
69. Toti, E., C.Y.O. Chen, M. Palmery, V.D. Valencia and I. Peluso, 2018. Non-provitamin A and provitamin A carotenoids as immunomodulators: Recommended dietary allowance, therapeutic index or personalized nutrition?. *Oxid. Med. Cell. Longevity*, Vol. 2018, 10.1155/2018/4637861
70. Weerasinghe, W.P.N.W. and S.A. Deraniyagala, 2016. Antioxidant of some Sri Lanka edemic medicinal plants. *Pharm. J. Sri Lanka*, 6: 9-14.
71. Muthukumar, P., N. Saraswathy, V. Aswitha, R. Balan, V.B. Gokhul, P. Indumathi and S. Yuvapriya, 2016. Assessment of total phenolic, flavonoid, tannin content and phytochemical screening of leaf and flower extracts from *Peltophorum pterocarpum* (DC.) Backer ex K. Heyne: A comparative study. *Pharmacog. J.*, 8: 140-143.
72. Cory, H., S. Passarelli, J. Szeto, M. Tamez and J. Mattei, 2018. The role of polyphenols in human health and food systems: A mini-review. *Front. Nutr.*, Vol. 5. 10.3389/fnut.2018.00087
73. Lin, D., M. Xiao, J. Zhao, Z. Li and B. Xing *et al.*, 2016. An overview of plant phenolic compounds and their importance in human nutrition and management of type 2 diabetes. *Molecules*, Vol. 21. 10.3390/molecules21101374
74. Panche, A.N., A.D. Diwan and S.R. Chandra, 2016. Flavonoids: An overview. *J. Nutr. Sci.*, Vol. 5, 10.1017/jns.2016.41
75. De Silva, S.F. and J. Alcorn, 2019. Flaxseed lignans as important dietary polyphenols for cancer prevention and treatment: Chemistry, pharmacokinetics and molecular targets. *Pharmaceuticals*, Vol. 12, 10.3390/ph12020068
76. Vinha, A.F., A.S.G. Costa, J.C.M. Barreira, R. Pacheco and M.B.P.P. Oliveira, 2016. Chemical and antioxidant profiles of acorn tissues from *Quercus* spp.: Potential as new industrial raw materials. *Ind. Crops Prod.*, 94: 143-151.
77. Kozłowska, A. and D. Szostak-Wegierek, 2018. Flavonoids-Food Sources, Health Benefits and Mechanisms Involved. In: *Bioactive Molecules in Food*, Merillon, J.M. and K. Ramawat (Eds.). , Springer, Cham, Switzerland, pp: 1-27.
78. Forni, C., F. Facchiano, M. Bartoli, S. Pieretti and A. Facchiano *et al.*, 2019. Beneficial role of phytochemicals on oxidative stress and age-related diseases. *BioMed Res. Int.*, Vol. 2019, 10.1155/2019/8748253
79. Liu, W., X. Bai, A. Zhang, J. Huang, S. Xu and J. Zhang, 2019. Role of exosomes in central nervous system diseases. *Frontiers Mol. Neurosci.*, Vol. 12, 10.3389/fnmol.2019.00240
80. Mojzer, B.E., K.M. Hrnčić, M. Skerget, Z. Knez and U. Bren, 2016. Polyphenols: Extraction methods, antioxidative action, bioavailability and anticarcinogenic effects. *Molecules*, Vol. 21, 10.3390/molecules21070901
81. Saeed, F., M. Afzaal, B. Niaz, M.U. Arshad, T. Tufail, M.B. Hussain and A. Javed, 2018. Bitter melon (*Momordica charantia*): A natural healthy vegetable. *Int. J. Food Prop.*, 21: 1270-1290.
82. Aghajanian, P., S. Hall, M.D. Wongworawat and S. Mohan, 2015. The roles and mechanisms of actions of vitamin C in bone: New developments. *J. Bone Mineral Res.*, 30: 1945-1955.
83. Chambial, S., S. Dwivedi, K.K. Shukla, P.J. John and P. Sharma, 2013. Vitamin C in disease prevention and cure: An overview. *Indian J. Clin. Biochem.*, 28: 314-328.
84. Young, J.I., S. Zuchner and G. Wang, 2015. Regulation of the epigenome by vitamin C. *Annual Rev. Nutr.*, 35: 545-564.
85. Grosso, G., R. Bei, A. Mistretta, S. Marventano and G. Calabrese *et al.*, 2013. Effects of vitamin C on health: A review of evidence. *Front Biosci.*, 18: 1017-1029.
86. Singh, A., S. Singh and S.M. Prasad, 2016. Role of medicinal plants for health perspective: Special reference to antioxidant potential. *J. Chem. Biol. Ther.*, 1: 1-5.
87. Datta, M. and M.Z. Vitolins, 2016. Food fortification and supplement use-are there health implications?. *Crit. Rev. Food Sci. Nutr.*, 56: 2149-2159.

88. De Oliveira, V.S., F.S. Ferreira, M.C.R. Cople, T.D.S. Labre, I.M. Augusta, O.D. Gamallo and T. Saldanha, 2018. Use of natural antioxidants in the inhibition of cholesterol oxidation: A review. *Compr. Rev. Food Sci. Food Safety*, 17: 1465-1483.
89. Dimov, I., N. Petkova, G. Nakov, I. Taneva, I. Ivanov and V. Stamatovska, 2018. Improvement of antioxidant potential of wheat flours and breads by addition of medicinal plants. *Ukrainian Food J.*, 7: 671-681.
90. Datta, S., B.K. Sinha, S. Bhattacharjee and T. Seal, 2019. Nutritional composition, mineral content, antioxidant activity and quantitative estimation of water soluble vitamins and phenolics by RP-HPLC in some lesser used wild edible plants. *Heliyon*, Vol. 5, No. 3. 10.1016/j.heliyon.2019.e01431
91. Richard, T.S., A.H.N. Kamdje and F. Mukhtar, 2015. Medicinal plants in breast cancer therapy. *J. Dis. Med. Plants*, 1: 19-23.
92. Kuruppu, A.I., P. Paranagama and C.L. Goonasekara, 2019. Medicinal plants commonly used against cancer in traditional medicine formulae in Sri Lanka. *Saudi Pharm. J.*, 27: 565-573.
93. Katla, V.R., R. Syed, C.S. Kuruva, H.K. Kuntrapakam and N.R. Chamarthi, 2013. Synthesis of novel phosphorylated guanidine derivatives from cyanamide and their anti-inflammatory activity. *Chem. Pharm. Bull.*, 61: 25-32.
94. Elisha, I.L., J.P. Dzoyem, L.J. McGaw, F.S. Botha and J.N. Eloff, 2016. The anti-arthritis, anti-inflammatory, antioxidant activity and relationships with total phenolics and total flavonoids of nine South African plants used traditionally to treat arthritis. *BMC Complement. Altern. Med.*, Vol. 16. 10.1186/s12906-016-1301-z
95. Kamble, A.A. and D.K. Nazia, 2017. *In vitro* anti-arthritis activity of *Vitex negundo* and *Punica granatum*. *Res. J. Pharm.*, 6: 5-7.
96. Kiranmayi, G.V.N., 2018. Preliminary phytochemical screening and *in vitro* evaluation of anti-inflammatory, antiarthritic and thrombolytic activities of ethanolic leaf extract of *Bauhinia purpurea*. *Int. J. Green Pharm. (IJGP)*, 12: S241-S247.
97. Fu, H., E.J. Ward and F.M. Marelli-Berg, 2016. Mechanisms of T cell organotropism. *Cell. Mol. Life Sci.*, 73: 3009-3033.
98. Rahman, M., S. Beg, A. Verma, F.A. Al Abbasi and F. Anwar *et al.*, 2017. Phytoconstituents as pharmacotherapeutics in rheumatoid arthritis: Challenges and scope of nano/submicromedicine in its effective delivery. *J. Pharm. Pharmacol.*, 69: 1-14.
99. Walling, B.L. and M. Kim, 2018. LFA-1 in T cell migration and differentiation. *Frontiers Immunol.*, 9: 1-10.
100. Mellado, M., L. Martinez-Munoz, G. Cascio, P. Lucas, J.L. Pablos and J.M. Rodriguez-Frade, 2015. T cell migration in rheumatoid arthritis. *Frontiers Immunol.*, Vol. 6, 10.3389/fimmu.2015.00384
101. Khairutdinov, V.R., A.F. Mikhailichenko, I.E. Belousova, E.S. Kuligina, A.V. Samtsov and E.N. Imyanitov, 2017. The role of intradermal proliferation of T-cells in the pathogenesis of psoriasis. *Anais Brasileiros Dermatologia*, 92: 41-44.
102. Tai, Y., Q. Wang, H. Korner, L. Zhang and W. Wei, 2018. Molecular mechanisms of T cells activation by dendritic cells in autoimmune diseases. *Front. Pharmacol.*, Vol. 9, 10.3389/fphar.2018.00642
103. Obaseki, O.E., O.I. Adesegun, G.N. Anyasor and O.O. Abebawo, 2016. Evaluation of the anti-inflammatory properties of the hexane extract of *Hydrocotyle bonariensis* Comm. Ex Lam. leaves. *Afr. J. Biotechnol.*, 15: 2759-2771.
104. Sangeetha, G. and R. Vidhya, 2016. *In vitro* anti-inflammatory activity of different parts of *Pedaliium murex* (L.). *Inflammation*, 4: 31-36.
105. Ijoma, K.I., V.I.E. Ajiwe and C.I. Awuzie, 2016. Antimicrobial analysis and structural elucidation of active compounds of *Dialium indum* leaves extract (Velvet Tamarind). *Int. J. Pharm. Chem.*, 6: 237-241.
106. Dhale, D.A., D.R. Chamle and V.H. Panchal, 2010. Evaluations of phytochemical constituents and antimicrobial activity of *Butea monosperma* (Fabaceae). *J. Phytol.*, 2: 17-21.
107. Chomel, M., M. Guittonny Larcheveque, C. Fernandez, C. Gallet and A. DesRochers *et al.*, 2016. Plant secondary metabolites: A key driver of litter decomposition and soil nutrient cycling. *J. Ecol.*, 104: 1527-1541.
108. Ali, M., B.H. Abbasi, N. Ahmad, H. Khan and G.S. Ali, 2017. Strategies to enhance biologically active-secondary metabolites in cell cultures of *Artemisia*-current trends. *Crit. Rev. Biotechnol.*, 37: 833-851.