

Epidemiologic Pattern of Skin Malignancies in Semnan, Iran Between 1999 and 2007 and Comparing it with Meta-Analysis of Published Papers in World Between 2000 and 2008

¹Semnani Vahid, ¹A. Toussy Jafar, ²Soltany Setareh and ³Toghraie Mirmohammad
¹Department of Pathology, ²Department of Surgery,
³Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran

Abstract: The skin is the body's largest organ. Melanoma and NMSC are now the most common types of cancer in white populations and both show an increasing incidence rate worldwide. The goal of this study was answering questions about skin malignancies epidemiologic pattern in Semnan. This study carried out in Semnan, Iran, from 1999-2007. Results of data from Semnan compared with meta-analysis results. Frequency of skin malignancies was 18.40%. Totally, the skin cancers were most prevalent malignancy. The most prevalent skin malignancies were BCC 65.1, SCC 28.5 and melanoma 1.07%. Male sex ratio was 61.87% for all cases, 56.9% for BCC and 72.2% for SCC cases ($p = 0.02$). Mean age was 64.92 years for all skin malignancies, 65.74 in males and 63.74 years for females ($p > 0.05$). This was 63.42 and 68.25 years for BCC and SCC, respectively ($p = 0.026$). The most prevalent sites were nose or other parts of face with 40.21% and scalp and neck with 22.78%. The most prevalent site for BCC and SCC was nose or other parts of face to with 47.5 and 26.6%, respectively ($p = 0.002$). Present results showed relatively constant trends by time for incidence of skin malignancies, including BCC and SCC and for sex and age. Epidemiologic pattern of skin cancers in Semnan was roughly comparable with the corresponding indexes in world and Iran especially for NMSC (BCC and SCC) comparing with other parts of Iran and Middle Asia, but the incidence of melanoma was lower.

Key words: Epidemiology, cancer, malignancy, skin, BCC, SCC, melanoma, meta-analysis

INTRODUCTION

The skin is the body's largest organ. The epidermis is made up of squamous cells, basal cells and contains melanocytes. Cancer may develop in any of the cell types: Squamous Cell Carcinoma (SCC), Basal Cell Carcinoma (BCC) and melanoma. Basal cell and squamous cell carcinomas are >10 times as common as melanoma but account for less morbidity and mortality. Nonmelanoma Skin Cancers (NMSC) are more prevalent in sun-exposed areas such as scalp, neck and face, old ages and mostly in male sex. Skin melanoma usually begins in a mole. It can occur on any skin surface in men, melanoma is often found on the trunk or the head and neck in women, it often develops on the lower legs (Johnson, 2004).

Melanoma and NMSC are now the most common types of cancer in white populations. Both tumor entities show an increasing incidence rate worldwide but a stable or decreasing mortality rate. Skin cancer is the most common form of cancer in the United States (Leiter and Garbe, 2008).

The rising incidence rates of NMSC are probably caused by a combination of increased sun exposure or

exposure to Ultraviolet (UV) light, increased outdoor activities, changes in clothing style, increased longevity, ozone depletion, genetics and in some cases, immune suppression. A dose-dependent increase in the risk of Squamous Cell Carcinoma (SCC) of the skin was found associated with exposure to Psoralen and UVA irradiation. An intensive UV exposure in childhood and adolescence was causative for the development of Basal Cell Carcinoma (BCC), whereas for the aetiology of SCC a chronic UV exposure in the earlier decades accused (Leiter and Garbe, 2008). Some experts say the rise in incidence (trend) of melanoma reflects a true increase in the disease, while others contend it is an artifact of more intensive recent surveillance. Others suggest that the rise in melanoma incidence may in part reflect longer life expectancy as well as efforts to detect melanoma earlier. The incidence of thin invasive lesions is increasing faster than that of thick ones, which reflects earlier detection by physicians and greater public awareness of warning signs of skin cancer. Among the reasons for these trends, increased exposure to UV radiation because of lifestyle changes is generally recognized as an important factor (Johnson, 2004).

In Iran and many underdeveloped or developing countries, including many Asian, African and other countries, well established registration systems, proper hard or soft databases and data validation and supervision don't exist, so many mistakes are seen in the data entry and output generation from these data. The aim of this study was correcting these data in Semnan pathology registration systems by pathology and database management experts, then creating valid outputs and corresponding world and Iran ASRs. The final goal was answering questions about skin malignancies epidemiologic pattern in Semnan (as a representative sample for Iranian population), for example trends in rate, changes in sex and age and many others; these questions in many instances previously answered based on incorrect data or personal nonscientific observations.

MATERIALS AND METHODS

This study carried out in Semnan, Iran, from 1999-2007. All of the pathology reports of three pathology departments in this city during these years were collected (this data collected for Cancer Registry of Semnan or archived routinely). Data entered into the software specially designed for this purpose manually for years 1999-2001 and by converting electronic data from DBF (Fox pro) format to MDB (Access) format for other years; then all cases reviewed and corrected if necessary according to the ICD-O codes (International Classification of Diseases for Oncology) by two pathologists. Corrected data exported to SPSS format and used for statistical analysis.

Analytical methods: The software for entering data was designed by Access and VBA programming language (Microsoft Access, Microsoft Office 2007, Microsoft Corporation). Analytical tests performed with SPSS (release 11.5.0, ©SPSS Inc.) and EpiInfo (version 6.4, WHO and CDC) statistical packages. Calculating corresponding Age Standardized incidence Rates (ASR) and Standardized Incidence Ratios (SIR) for Iran and world done with software designed for this purpose by Excel and VBA (Microsoft Excel, Microsoft Office, 2007, Microsoft Corporation). Population data used for these calculations are shown in Table 1. Populations and age distribution data of years 1990, 1996 and 2007 for Semnan, Iran and world, respectively used as reference for calculating incidences and ASRs.

Meta analysis methods: We used Pubmed (<http://www.pubmed.com> or <http://www.ncbi.nlm.nih.gov/pubmed>) and SID (<http://www.sid.ir>) search engines for finding

Table 1: Population of Semnan, Iran and world used for calculating ASRs

Years	Semnan*	Iran*	World**
1999	154422	62736002	5948042000
2000	159249	63663942	6034542000
2001	164108	64528162	6122541000
2002	168937	65466600	6211579000
2003	173632	66365200	6301911000
2004	178527	67263800	6393556000
2005	183448	68162400	6486792000
2006	188211	69061000	6581126000
2007	193038	69959600	6676832000

*<http://amar.sci.org.ir>; **<http://www.ibiblio.org/lunarbin/worldpop>

relative researchers. Search keyword used for the Pubmed was skin and (cancer or cancers or malignancy or malignancies) and prevalence and incidence and registry (Title/Abstract) without any limitations that returned 687 results. All abstracts of these researchers reviewed and appropriate ones selected. For SID skin cancer and equivalent Persian search keywords used that returned 9 results. All results entered into database designed by SPSS and analyzed by it.

RESULTS AND DISCUSSION

Totally, 1528 malignancies registered, 281 cases were skin malignancies (18.40 and CI95%: 16.51-20.39), 61.78% (CI95%: 56.05-67.44) of them were male and 38.12% (CI95%: 32.56-43.95) female with male/female sex ratio of 1.62 (CI95%: 1.46-1.85). Totally, the skin cancers were most prevalent malignancy. The stomach was the second prevalent site totally (13.48 and CI95%: 11.84-15.26).

The most prevalent skin malignancies were BCC 65.1% (CI95%: 59.41-70.53), SCC 28.5% (CI95%: 23.42-33.96) and melanoma 1.07% (CI95%: 0.27-2.88). Incidences and corresponding ASRs for Iran and world are shown in Table 2. As shown in Table 2, total incidence of skin malignancies is relatively constant by year but that of males is decreased and that of females is increased; of course no one of these trends are significant ($p > 0.05$ by linear regression analysis). Prevalence of other morphologies and incidence of cases of BCC, SCC and melanoma are shown in Table 3.

Male sex ratio was 61.87% (CI95%: 56.05-67.44) for all cases, 56.9% for BCC and 72.2% for SCC cases ($p = 0.02$, Pearson Chi-square (χ^2) = 5.401). The trend for sex by time (year) showed no significant change ($p > 0.05$ by linear and logistic regression analysis). This was true for BCC and SCC too.

Mean age was 64.92 years (SEM: 1.02, SD: 14.68) for all skin malignancies, 65.74 in males and 63.74 years for females ($p > 0.05$ with t-test). This was 63.42 and 68.25 years for BCC and SCC, respectively ($p = 0.026$, $t = -2.241$). The trend for age by time (year) showed no significant change ($p > 0.05$ by linear and logistic regression analysis). This was true for BCC and SCC too.

Table 2: Incidence and corresponding ASRs of skin malignancies according to year and sex in Semnan, Iran (1999-2007)

Years	Frequency	%	Incidence (100,000)			Iran ASR (100,000)			World ASR (100,000)		
			Male	Female	Total	Male	Female	Total	Male	Female	Total
1999	28	10.0	17.82	18.46	18.13	17.14	17.27	17.06	22.38	23.70	23.37
2000	27	9.6	24.68	8.95	16.95	23.75	8.37	15.86	31.01	11.49	22.18
2001	32	11.4	22.75	16.13	19.50	22.01	14.33	18.24	27.96	22.85	25.45
2002	39	13.9	37.23	8.44	23.09	35.72	7.89	21.41	47.24	10.83	30.79
2003	30	10.7	21.51	12.90	17.28	20.72	11.67	16.11	26.90	17.71	22.75
2004	31	11.0	14.31	20.53	17.36	13.66	18.18	15.91	18.54	29.28	23.80
2005	29	10.3	22.50	8.88	15.81	21.76	7.96	14.84	27.71	12.40	20.50
2006	30	10.7	15.66	16.23	15.94	15.09	14.64	14.87	19.57	22.38	20.99
2007	35	12.5	21.38	14.76	18.13	20.43	14.21	16.93	27.59	17.84	23.81
Total ¹	281	100.0	197.84	125.28	162.19	190.28	114.52	151.23	248.90	168.48	213.64
Mean	-	-	21.98	13.92	18.02	21.14	12.72	16.80	27.66	18.72	23.74
SEM	-	-	2.23	1.47	0.74	2.14	1.32	0.68	2.82	2.12	1.02
SIR ²	-	-	-	-	-	0.96	0.91	0.92	1.26	1.34	1.32

ASR: Age Standardized incidence Rate, SIR: Standardized Incidence Ratio, ¹Total (sum) of all incidences and ASRs is a rough estimate of 9 years prevalence (cumulative incidence), ²Calculated for mean of ASRs (Iran or world) relative to Semnan

The most prevalent sites were nose or other parts of face with 40.21% (CI95%: 34.59-46.03) and scalp and neck with 22.78% (CI95%: 18.16-27.95). The most prevalent site for BCC and SCC was nose or other parts of face too with 47.5 and 26.6%, respectively (p = 0.002, Pearson Chi-square (χ^2) = 10.004). The less prevalent site was skin of anus with 0.7% (2 cases of SCC). SCC was more prevalent in skin of lip (13.9%) and trunk-limbs (15.2%) relative to BCC (1.6 and 3.2%, respectively, p = 0.002 for lip and <0.001 for trunk-limbs with Chi-square (χ^2) test). Frequencies of other topographies are shown in Table 4.

Meta-analysis results: From 687 papers returned from the Pubmed search engine (with publication date between 1970 and 2008), 327 (47.6%) results were published between years 2000 and 2008 that 158 (48.32% of results between 2000 and 2008) of them selected and downloaded. After evaluating these papers, 73 papers (22.32% of results between 2000 and 2008) containing data about prevalence, incidence, ASR, or trends of skin cancers selected for meta-analysis (Schuz *et al.*, 2000; Plesko *et al.*, 2000; Holme *et al.*, 2000, 2001; Fidaner *et al.*, 2001; Kittler *et al.*, 2001; Garbe and Blum, 2001; Ocaña-Riola *et al.*, 2001; Ishihara *et al.*, 2001; Stang *et al.*, 2001, 2003, 2006; Wassberg *et al.*, 2001; Liao *et al.*, 2001; Hemminki *et al.*, 2001, 2003; Micheli *et al.*, 2002; Nola *et al.*, 2002; Verdecchia *et al.*, 2002; Crocetti *et al.*, 2002; Milan *et al.*, 2002; Lutz *et al.*, 2003; Forman *et al.*, 2003; Boi *et al.*, 2003, 2008; Sadjadi *et al.*, 2003; Barro-Traore *et al.*, 2003; Katalinic *et al.*, 2003; Al-Aboud *et al.*, 2003; Brisson *et al.*, 2003; Ulmer *et al.*, 2003; Martin *et al.*, 2004; Al-Maghrabi *et al.*, 2004; Uhoda *et al.*, 2004; De Vries *et al.*, 2004, 2005; Quatresooz *et al.*, 2004; San Sebastian and Hurtig, 2004; Asulin *et al.*, 2004; Andersen *et al.*, 2005; Paavilainen *et al.*, 2005; Vaktskjold *et al.*, 2005;

Babaei *et al.*, 2005; Bhurgri *et al.*, 2005; Stang *et al.*, 2006; Boniol *et al.*, 2006; Omari *et al.*, 2006; Lasithiotakis *et al.*, 2006a, 6; Downing *et al.*, 2006, 2008; AIRT Working Group, 2006; Jamal *et al.*, 2006; Piñeros *et al.*, 2006; El-Mistiri *et al.*, 2003; Hayes *et al.*, 2007; Hui *et al.*, 2007; Golger *et al.*, 2007; Brewster *et al.*, 2007; Hoey *et al.*, 2007; Brathwaite *et al.*, 2007; Youl *et al.*, 2007; Nasseri *et al.*, 2007; Valentin *et al.*, 2007; Maalej *et al.*, 2007; Noorbala and Kafaie, 2007; Moan *et al.*, 2008; Siesling *et al.*, 2008; Pellacani *et al.*, 2008; Dal *et al.*, 2008; Karim-Kos *et al.*, 2008; Zeb *et al.*, 2008; Soerjomataram *et al.*, 2008; Yavari *et al.*, 2008). From the 9 papers returned from the SID search engine, 6 papers (66.67%) selected for meta-analysis (Naimi, 2001; Golchai *et al.*, 2000; Tousi *et al.*, 2004; Shahbazi, 2006; Amouzgar *et al.*, 2006; Nour Bala, 2007).

Pubmed results: Year 2007 with 12 papers (16.4%) had the maximum frequency. Europe with 44 papers (60.3%) and Central America with one paper (1.4%) had the highest and lowest frequencies. Middle Asia had published 10 papers (13.7%). In Europe Germany and Italy with 6 (13.6) and 5 (11.4%) papers had the highest frequency. About melanoma, 26 papers published that Germany with 4 papers (15.4%) had the highest frequency. Four published papers (5.48%) from Iran listed in Pubmed that one of them was from Semnan. Sixteen papers (21.9) studied all skin cancers, 26 papers (35.6%) studied only melanoma and 3 papers (4.1%) studied only NMSC; other papers contained data about other cancers plus skin or studied BCC or SCC restrictively. The most common morphology was BCC (all of the papers) and the most common sites were nose or other parts of face in half the papers and scalp and neck in other half. The most common topography for melanoma was skin, lower limbs and hip totally and in females and skin, trunk for males in all papers.

Table 3: Prevalence and incidence of skin malignancies, Semnan city, Iran (1999-2007)

Malignancy	Frequency	%	Incidence* (100,000)			Iran ASR* (100,000)			World ASR* (100,000)		
			Male	Female	Total	Male	Female	Total	Male	Female	Total
BCC	183	65.10	12.22	10.55	10.88	11.71	10.23	10.25	15.56	12.51	13.98
SCC	80	28.50	10.18	4.22	7.25	9.71	3.98	6.68	13.21	5.34	9.83
Melanoma	3	1.07	1.02	2.11	1.55	1.00	1.89	1.47	1.17	2.95	1.97
MFH and DFSP	4	1.40	-	-	-	-	-	-	-	-	-
Vascular	2	0.80	-	-	-	-	-	-	-	-	-
Others	9	3.20	-	-	-	-	-	-	-	-	-

*BCC and SCC for year 2007, melanoma for all years (due to low incidence)

Table 4: Topographies of skin malignancies, Semnan, Iran (1999-2007)

Site	Frequency	%
Skin, nose or other parts of face	113	40.2
Skin, scalp and neck	64	22.8
Skin, eyelid	20	7.1
Skin, lip	14	5.0
Skin, external ear	13	4.6
Skin, trunk	11	3.9
Skin, lower limbs and hip	8	2.8
Skin, upper limb and shoulder	5	1.8
Skin, anus	2	0.7
Skin, nose	31	11.0
Total	281	100.0

About 83.3% of papers that evaluated trends in incidence of skin cancers reported an increase in this trend; this was true for all papers about BCC and SCC too. In the case of melanoma, 80.9% of papers reported increase, 14.3% constant trend and 4.8% decrease in the trend.

Six papers published by countries of Middle East excluding Iran, 3 by Pakistan, 2 by Saudi Arabia and 1 by Jordan that the last two countries have hot and dry climate like Semnan and placed near or in the desert. Most common morphology was BCC, most common topography scalp and neck for all skin cancers and BCC, nose or other parts of face for SCC. In the case of melanoma, one paper reported skin, lower limbs and hip as the most common topography. Only one paper evaluated the trend of all skin cancers and reported increase of it.

Papers from Iran (Pubmed or SID search engines): Ten papers published from Iran (4 listed in Pubmed and other 6 listed in SID). Half of these papers reported skin cancers as the most prevalent malignancy of the body, all of them noted BCC as the most common skin cancer and nose or other parts of face as the most common topography in skin cancers.

Other data from all papers (Pubmed or SID search engines) are shown in Table 5. According to these results as shown in Table 5, epidemiologic pattern of skin cancers in Semnan is roughly comparable with the corresponding indexes in world and Iran, except that the incidence of melanoma was lower; this is especially, true for NMSC (BCC and SCC) comparing with other parts of Iran and Middle Asia. Relative frequency and incidence of skin

cancers in Semnan are relatively higher than world but similar to Middle Asia; this may be due to similar climates in Semnan and other middle Asia countries especially, Saudi Arabia and Jordan that have hot and dry climates. Similar clothing and body covering customs and social behaviors such as tanning, may be proposed as the cause for this equality. Totally, age and sex patterns are very similar in world, Middle Asia, Iran and Semnan. Relative frequencies of BCC and SCC in skin cancers, age and sex have relatively same distribution in world, Middle Asia and Iran in comparison with Semnan. Relative frequency and incidence of melanoma is significantly lower than world and Middle Asia in Semnan and this is more similar with Iran; different skin type, climates, genetic predisposition, life expectancy and screening-treatment programs may be the cause. This is in contrast with developed countries, notably those with fair-skinned populations (Stang *et al.*, 2001) that have more susceptible skin type and also better screening-treatment programs for skin and other cancers and so longer life expectancies. Clothing and skin type of the individual can play protective role (Al-Aboud *et al.*, 2003; Noorbala and Kafaie, 2007). Different exposure to UV radiation because of different lifestyles is an important factor too (Johnson, 2004). Both hereditary and environmental factors are implicated in the etiology of cutaneous neoplasms. Studies of twins suggests that environmental and not hereditary effects are most important in the causation of malignant skin cancers in a white population with low levels of sun exposure (Milan *et al.*, 2002) but we can't exclude it totally.

As seen in these results epidemiologic patterns for morphology and topography of skin cancers, including BCC, SCC and melanoma in Semnan show total similarity to world, Middle Asia and Iran patterns.

The results showed relatively constant trends by time for incidence of skin malignancies, including BCC and SCC and for sex and age. This is in contrast to nearly all results of other researches (Johnson, 2004; Stang *et al.*, 2001; Wassberg *et al.*, 2001; Nola *et al.*, 2002; Lutz *et al.*, 2003; Ulmer *et al.*, 2003; De Vries *et al.*, 2004, 2005; Quatresooz *et al.*, 2004; Downing *et al.*, 2006; Lasithiotakis *et al.*, 2006a, b; Hayes *et al.*, 2007).

Table 5: Results of meta-analysis data of skin cancers from published papers between years 2000 and 2008 in comparison with Semnan

Index*	World	Middle East**	Iran	Semnan
Relative Freq. (%)	14.58		19.58	18.4
Incidence (100,000)	13.50 (M: 12.4, F: 9.4)	16	11 (M: 12.4, F: 9.4)	18.02 (M: 21.98, F: 13.92)
World ASR (100,000)	07.80 (M: 6.7, F: 7.46)		M: 10.2, F: 7.46	23.74 (M: 27.66, F: 18.72)
Male relative Freq. (%)	63.82	65.39	61.47	61.87
Age (years)	75.00 (M: 61, F: 60)	75	64.49	64.92
BCC relative Freq. (%)	54.41	48.3	71.26	65.10
BCC world ASR (100,000)	31.70 (M: 35, F: 26.87)	M: 23.2, F: 19.7		13.98 (M: 15.56, F: 12.51)
BCC male relative Freq. (%)	57.98			56.90
SCC relative Freq. (%)	23.77		21.32	28.50
SCC world ASR (100,000)	04.40 (M: 10.7, F: 4.79)			9.83 (M: 13.21, F: 5.34)
SCC male relative Freq. (%)	71.83			72.20
Melanoma relative Freq. (%)	08.16	11.94	3.28	1.07
Melanoma incidence (100,000)	09.25 (M: 8.83, F: 9.95)			1.55 (M: 1.17, F: 2.95)
Melanoma male relative Freq. (%)	49.93	54.55	47.5	

*Mean of values of all papers (empty cells are due to inadequate data), **Excluding Iran, M: Value for males, F: Value for females

Incidence of cancer may vary within a country and over time because of previous differences in exposure to risk factors or interventions for early detection (screening) (Siesling *et al.*, 2008). This may reflect a true increase, or may be an artifact of more intensive recent surveillance. Longer life expectancy as well as efforts to earlier detection may be the cause too; for example, greater public awareness of warning signs of skin cancer (Johnson, 2004).

Increased exposure to UV radiation because of lifestyle changes (Johnson, 2004; Boniol *et al.*, 2006), increased accumulated sun exposure and increasing incidence among the elderly (Wassberg *et al.*, 2001; De Vries *et al.*, 2005), effectiveness of the health system on cancer control (Crocetti *et al.*, 2002), paradoxical effect of better treatments and thereby survival in well-developed economies (Lutz *et al.*, 2003) and belonging to white populations (Ulmer *et al.*, 2003) are generally recognized as important factors. Trend of the increase in incidence of melanoma is more impressive than that of SCC and BCC; this trend is particularly obvious for melanoma affecting young adults (Uhoda *et al.*, 2004). Incidence of BCC in the Netherlands is increasing rapidly, especially at body sites that are not chronically exposed to sunlight; the most likely explanation is an increased intermittent overexposure to UV radiation. More sensitive detection using the combination of clinical dermoscopy and laboratory immunohistochemistry are among these factors too (Quatresooz *et al.*, 2004). The relative increase of SCC of the trunk and upper limbs is a plausible reflection of intentional tanning (Dal *et al.*, 2008).

Small proportion of papers reported decrease in incidence or constant trend of skin cancers (Martin *et al.*, 2004; Boi *et al.*, 2008).

CONCLUSION

In the end, comparison of the data with the only study that studied cancer prevalence in Semnan

(Babaei *et al.*, 2005) seems helpful. The study reported gastric cancer as the most prevalent malignancy (in contrast to this study that reported skin cancer as the most prevalent and gastric cancer as the second one), incidence rate of 12.4 for males and 9.4/100,000 for females (this study: 21.98 and 13.92, respectively) and world ASR of 10.2 for males and 7.46/100,000 for females (This study: 27.66 and 18.72, respectively). According to using same archive and electronic data from Cancer Registry of Semnan, this discrepancy can show the reason for the goal that mentioned in objectives: the aim of this study was correcting these data in Semnan pathology registration systems by pathology and database management experts, then creating valid outputs. We think different approaches in validation of data and role of experts in pathology and informatics can explain this difference.

REFERENCES

- AIRT Working Group, 2006. Italian cancer figures-report 1. Incidence, mortality and estimates. *Epidemiol Prev.*, 30 (1 Suppl. 2): 8-10, 12-28, 30-101 Passim. PMID: 16826725.
- Al-Aboud, K.M., K.A. Al-Hawsawi, M.A. Bhat, V. Ramesh and S.M. Ali, 2003. Skin cancers in Western Saudi Arabia. *Saudi Med. J.*, 24 (12): 1381-1387. PMID: 14710289.
- Al-Maghrabi, J.A., A.S. Al-Ghamdi and H.A. El-Hakeem, 2004. Pattern of skin cancer in Southwestern Saudi Arabia. *Saudi Med. J.*, 25 (6): 776-779. PMID: 15195210.
- Amouzgar, M.H., Y.P.M. Javad and M.E. Rad, 2006. Frequency of different skin cancers in Qaem hospital, Mashhad, from 1975-1995: A cross-sectional study. *Iranian J. Dermatol.*, 9(1 (35): 28-34. <http://sid.ir/en/ViewPaper.asp?ID=48172>.

- Andersen, S.L., D.F. Terry, M.A. Wilcox, T. Babineau, K. Malek and T.T. Perls, 2005. Cancer in the oldest old. *Mech. Ageing Dev.*, 126 (2): 263-367. PMID: 15621206.
- Asulin, Y., T.J. McCann, C.W. McCarty, R.W. Hage, P.J. Rooney and C.N. Macpherson, 2004. Cancer incidence and mortality in Grenada 1990-2000. *West Indian Med. J.*, 53 (6): 368-373. PMID: 15816262.
- Babaei, M., S. Mousavi, M. Malek, G. Tosi, Z. Masoumeh, N. Danaei and G. Gafar, 2005. Cancer occurrence in Semnan, Iran: Results of a population based cancer registry. *Asian Pac. J. Cancer Prev.*, 6 (2): 159-164. PMID: 16101326.
- Barro-Traore, F., A. Traore, I. Konate, S.S. Traore, N.O. Sawadogo, I. Sanou, B.R. Soudre, E. Heid and E. Grosshans, 2003. Epidemiological features of tumors of the skin and mucosal membranes in the Department of Dermatology at the Yalgado Ouedraogo National Hospital, Ouagadougou, Burkina Faso. *Sante*, 13 (2): 101-104. PMID: 14530122.
- Bhurgri, Y., A. Bhurgri, S. Pervez, M. Bhurgri, N. Kayani, R. Ahmed, A. Usman and S.H. Hasan, 2005. Cancer profile of hyderabad, Pakistan 1998-2002. *Asian Pac. J. Cancer Prev.*, 6 (4): 474-480. PMID: 16435995.
- Boi, S., M. Cristofolini, R. Micciolo, E. Polla and P. Dalla Palma, 2003. Epidemiology of skin tumors: Data from the cutaneous cancer registry in Trentino, Italy. *J. Cutan Med. Surg.*, 7 (4): 300-305. PMID: 12879331.
- Boi, S., M. Cristofolini, R. Micciolo, E. Polla, S. Piffer and M. Gentilini, 2008. Incidence and mortality data from cutaneous melanoma in Trentino: Registry based study. *J. Cutan Med. Surg.*, 12 (2): 59-63. PMID: 18346401.
- Boniol, M., B.K. Armstrong and J.F. Dore, 2006. Variation in incidence and fatality of melanoma by season of diagnosis in new South Wales, Australia. *Cancer Epidemiol Biomarkers Prev.*, 15 (3): 524-526. PMID: 16537711.
- Brathwaite, A.F., N. Brathwaite and A. Del Riego, 2007. Epidemiological profile of cancer for grand Bahama residents: 1988-2002. *West Indian Med. J.*, 56 (1): 26-33. PMID: 17621841.
- Brewster, D.H., L.A. Bhatti, J.H. Inglis, E.R. Nairn and V.R. Doherty, 2007. Recent trends in incidence of nonmelanoma skin cancers in the East of Scotland, 1992-2003. *Br. J. Dermatol.*, 156 (6): 1295-300. PMID: 17535229.
- Brisson, C., C. Reynaud-Hautin and E. Bure, 2003. Prospective epidemiological study on cutaneous melanomas in the Vendee area. *Ann. Dermatol. Venereol.*, 130 (12 Pt 1): 1121-1124. PMID: 14724514.
- Crocetti, E., E. Paci, G. Miccinesi, A.S. Costantini and M. Zappa, 2002. Time trends in cancer incidence and mortality in central Italy. *Eur. J. Cancer Prev.*, 11 (4): 387-395. PMID: 12195166.
- Dal, H., C. Boldemann and B. Lindelöf, 2008. Trends during a half century in relative squamous cell carcinoma distribution by body site in the Swedish population: Support for accumulated sun exposure as the main risk factor. *J. Dermatol.*, 35 (2): 55-62. PMID: 18271799.
- De Vries, E., M. Louwman, M. Bastiaens, F. De Gruijl and J.W. Coebergh, 2004. Rapid and continuous increases in incidence rates of basal cell carcinoma in the Southeast Netherlands since 1973. *J. Invest. Dermatol.*, 123 (4): 634-638. PMID: 15373766.
- De Vries, E., L.V. Van de Poll-Franse, W.J. Louwman, F.R. De Gruijl and J.W. Coebergh, 2005. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br. J. Dermatol.*, 152 (3): 481-488. PMID: 15787817.
- Downing, A., J.A. Newton-Bishop and D. Forman, 2006. Recent trends in cutaneous malignant melanoma in the Yorkshire, England; incidence, mortality and survival in relation to stage of disease (1993-2003). *Br. J. Cancer*, 95 (1): 91-95. PMID: 16755289.
- Downing, A., X.Q. Yu, J. Newton-Bishop and D. Forman, 2008. Trends in prognostic factors and survival from cutaneous melanoma in Yorkshire, UK and New South Wales, Australia between 1993 and 2003. *Int. J. Cancer*, 123 (4): 861-866. PMID: 18500743.
- El-Mistiri, M., A. Verdecchia, I. Rashid, N. El-Sahli, M. El-Mangush and M. Federico, 2003. Cancer incidence in Eastern Libya: The first report from the Benghazi cancer registry. *Int. J. Cancer*, 120 (2): 392-397. PMID: 17066425.
- Fidaner, C., S.Y. Eser and D.M. Parkin, 2001. Incidence in Izmir in 1993-1994: First results from Izmir cancer registry. *Eur. J. Cancer*, 37 (1): 83-92. PMID: 11165134.
- Forman, D., D. Stockton, H. Möller, M. Quinn, P. Babb, R. De Angelis and A. Micheli, 2003. Cancer prevalence in the UK: Results from the Europrevall study. *Ann. Oncol.*, 14 (4): 648-654. PMID: 12649115.
- Garbe, C. and A. Blum, 2001. Epidemiology of cutaneous melanoma in Germany and worldwide. *Skin Pharmacol Appl. Skin Physiol.*, 14 (5): 280-290. PMID: 11586069.
- Golchai, J., A. Sobhani, M.R. Feizikhah and Ch. Fathi, 2000. Evaluating of malignant skin tumors in Gilan, Iran. *Gilan. Med. Sci. J.*, 42: 1-5. <http://sid.ir/en/ViewPaper.asp?ID=57313814201>.
- Golger, A., D.S. Young, D. Ghazarian and P.C. Neligan, 2007. Epidemiological features and prognostic factors of cutaneous head and neck melanoma: A population-based study. *Arch. Otolaryngol. Head Neck. Surg.*, 133 (5): 442-427. PMID: 17515502.

- Hayes, R.C., S. Leonfellner, W. Pilgrim, J. Liu and D.N. Keeling, 2007. Incidence of nonmelanoma skin cancer in New Brunswick, Canada, 1992-2001. *J. Cutan. Med. Surg.*, 11 (2): 45-52. PMID: 17374314.
- Hemminki, K., X. Li, K. Pina, C. Granström and P. V. aittinen, 2001. The Nation-wide Swedish family-cancer database-updated structure and *Amiial* rates. *Acta Oncol.*, 40 (6): 772-777. PMID: 11765074.
- Hemminki, K., H. Zhang and K. Czene, 2003. Incidence trends and familial risks in invasive and *in situ* cutaneous melanoma by sun-exposed body sites. *Int. J. Cancer*, 104 (6): 764-771. PMID: 12640685.
- Hoey, S.E., C.E. Devereux, L. Murray, D. Catney, A. Gavin, S. Kumar, D. Donnelly and O.M. Dolan, 2007. Skin cancer trends in Northern Ireland and consequences for provision of dermatology services. *Br. J. Dermatol.*, 156 (6): 1301-1307. PMID: 17535230.
- Holme, S.A., K. Malinovsky and D.L. Roberts, 2001. Malignant melanoma in South Wales: Changing trends in presentation (1986-98). *Clin. Exp. Dermatol.*, 26 (6): 484-489. PMID: 11678870.
- Holme, S.A., K. Malinovsky and D.L. Roberts, 2000. Changing trends in non-melanoma skin cancer in South Wales (1988-98). *Br. J. Dermatol.*, 143 (6): 1224-1229. PMID: 11122025.
- Hui, S.K., W.Y. Tang, T.W. Wong, K.H. Lau, S. Lee, L.Y. Chong and K.K. Lo, 2007. Cutaneous melanoma: A population-based epidemiology report with 989 patients in Hong Kong. *Clin. Exp. Dermatol.*, 32 (3): 265-267. PMID: 17397350.
- Ishihara, K., T. Saida and A. Yamamoto, 2001. Japanese Skin Cancer Society Prognosis and Statistical Investigation Committee. Updated statistical data for malignant melanoma in Japan. *Int. J. Clin. Oncol.*, 6 (3): 109-116. PMID: 11706778.
- Jamal, S., S. Moghal, N. Mamoon, S. Mushtaq, M. Luqman and M. Anwar, 2006. The pattern of malignant tumours: Tumour registry data analysis, AFIP, Rawalpindi, Pakistan (1992-2001). *J. Pak. Med. Assoc.*, 56 (8): 359-362. PMID: 16967787.
- Johnson, C., 2004. State/National statistics: Basic epidemiology of skin cancer. Fourth annual conference on advancing cancer care skin cancer and management of treatment related Fatigue. McCleary Center Saint Alphonsus Regional Medical Center. <http://www.idcancer.org/Special/Melanoma.ppt>.
- Karim-Kos, H.E., E. De Vries, I. Soerjomataram, V. Lemmens, S. Siesling and J.W. Coebergh, 2008. Recent trends of cancer in Europe: A combined approach of incidence, survival and mortality for 17 cancer sites since the 1990s. *Eur. J. Cancer*, 44 (10): 1345-1389. PMID: 18280139.
- Katalinic, A., U. Kunze and T. Schäfer, 2003. Epidemiology of cutaneous melanoma and non-melanoma skin cancer in Schleswig-Holstein, Germany: Incidence, clinical subtypes, tumour stages and localization (epidemiology of skin cancer). *Br. J. Dermatol.*, 149 (6): 1200-1206. PMID: 14674897.
- Kittler, H., M. Binder, K. Wolff and H. Pehamberger, 2001. A 10-years analysis of demographic trends for cutaneous melanoma: Analysis of 2501 cases treated at the University Department of Dermatology in Vienna (1990-1999). *Wien Klin Wochenschr*, 113 (9): 321-327. PMID: 11388077.
- Lasithiotakis, K., U. Leiter, S. Kruger-Krasagakis, A. Tosca and C. Garbe, 2006. Comparative analysis of incidence and clinical features of cutaneous malignant melanoma in Crete (Greece) and Southern Germany (Central Baden-Württemberg). *Br. J. Dermatol.*, 154 (6): 1123-1127. PMID: 16704644.
- Lasithiotakis, K.G., U. Leiter, R. Gorkiewicz, T. Eigentler, H. Breuninger, G. Metzler, W. Strobel and C. Garbe, 2006. The incidence and mortality of cutaneous melanoma in Southern Germany: Trends by anatomic site and pathologic characteristics (1976-2003). *Cancer*, 107 (6): 1331-1339. PMID: 16909413.
- Leiter, U. and C. Garbe, 2008. Epidemiology of melanoma and nonmelanoma skin cancer the role of sunlight. *Adv. Exp. Med. Biol.*, 624: 89-103. PMID: 18348450.
- Liao, Y.H., K.H. Chen, M.P. Tseng and C.C. Sun, 2001. Pattern of skin diseases in a geriatric patient group in Taiwan: A 7-years survey from the outpatient clinic of a university medical center. *Dermatology*, 203 (4): 308-313. PMID: 11752818.
- Lutz, J.M., S. Francisci, E. Mugno, M. Usel, V. Pompe-Kirn, J.W. Coebergh and M. Bieslka-Lasota, 2003. Europrevall working group. Cancer prevalence in central Europe: The Europrevall study. *Ann. Oncol.*, 14 (2): 313-322. PMID: 12562661.
- Maalej, M., D. Hentati, M. Slimene, S. Kanoun, M. Hechiche, K. Ben Romdhane and M. Ben Abdallah, 2007. Skin cancer in Tunisia: A retrospective study: 1379 cases and risk factors. *Tunis Med.*, 85 (9): 728-733. PMID: 18254300.
- Martin, R.C. and E. Robinson, 2004. Cutaneous melanoma in Caucasian New Zealanders: 1995-1999. *ANZ J. Surg.*, 74 (4): 233-237. PMID: 15043734.
- Micheli, A., E. Mugno, V. Krogh, M.J. Quinn, M. Coleman, T. Hakulinen, G. Gatta, F. Berrino and R. Capocaccia, 2007. Europrevall working group. Cancer prevalence in European registry areas. *Ann. Oncol.*, 13 (6): 840-865. PMID: 12123330.
- Milan, T., P.K. Verkasalo, J. Kaprio, M. Koskenvuo and E. Pukkala, 2002. Malignant skin cancers in the finnish twin Cohort: A population-based study (1976-97). *Br. J. Dermatol.*, 147 (3): 509-512. PMID: 12207592.

- Moan, J., A.C. Porojnicu, A. Dahlback and R.B. Setlow, Addressing the health benefits and risks, involving vitamin D or skin cancer, of increased sun exposure. *Proc. Natl. Acad. Sci. USA*, 105 (2): 668-673. PMID: 18180454.
- Naimi, M., 2001-2002. Non melanotic skin cancer of head and neck review of 439 cases during last 5 years (1372-1376) from Omid and Chaem Hospital. *The Iranian J. Otorhinolaryngol.*, 13 (3-4 (28)): 30-36. <http://sid.ir/en/ViewPaper.asp?ID=47557>.
- Nasseri, K., P.K. Mills and M. Allan, 2004. Cancer incidence in the Middle Eastern population of California (1988-2004). *Asian Pac. J. Cancer Prev.*, 8 (3): 405-411. PMID: 18159978.
- Nola, I., B. Kruslin, D. Müller, L. Oremovic and M. Belicza, 2002. The rise in melanoma incidence in Croatia. *Acta Dermatovenerol. Croat.*, 10 (1): 3-7. PMID: 12137724.
- Noorbala, M.T. and P. Kafaie, 2007. Analysis of 15 years of skin cancer in central Iran (Yazd). *Dermatol. Online J.*, 13 (4): 1. PMID: 18318998.
- Nour Bala, M.T., 2007. Skin cancer in Yazd. *Iranian J. Dermatol.*, 10 (1 (39)): 13-19. <http://sid.ir/en/view-paper.asp?ID=80881>.
- Ocana-Riola, R., C. Martinez-Garcia, S. Serrano, A. Buendia-Eisman, C. Ruiz-Baena and J. Canela-Soler, 2001. Population-based study of cutaneous malignant melanoma in the Granada province (Spain) (1985-1992). *Eur. J. Epidemiol.*, 17 (2): 169-174. PMID: 11599692.
- Omari, A.K., M.R. Khammash and I. Matalka, 2006. Skin cancer trends in Northern Jordan. *Int. J. Dermatol.*, 45 (4): 384-388. PMID: 16650163.
- Paavilainen, V., J. Tuominen, E. Pukkala and K.M. Saari, 2005. Basal cell carcinoma of the eyelid in Finland during 1953-97. *Acta Ophthalmol Scand.*, 83 (2): 215-220. PMID: 15799736.
- Pellacani, G., G.L. Scocco, M. Vinceti, G. Albertini, A.A. Raccagni, L. Baldassari, S. Catrani, S. Donelli, P. Ghetti, A. Lanzoni, M. Leardini, M. Reggiani, M. Santini, I. Stanganelli, A. Virgili and S. Seidenari, 2008. Melanoma epidemic across the millennium: Time trends of cutaneous melanoma in Emilia-Romagna (Italy) from 1997-2004. *J. Eur. Acad. Dermatol. Venereol.*, 22(2): 213-218. PMID: 18211415.
- Piñeros, M., J. Ferlay and R. Murillo, 2006. Cancer incidence estimates at the national and district levels in Colombia. *Salud Publica Mex.*, 48 (6): 455-465. PMID: 17326341.
- Plesko, I., G. Severi, A. Obsitnikova and P. Boyle, 2000. Trends in the incidence of non-melanoma skin cancer in Slovakia (1978-1995). *Neoplasma*, 47 (3): 137-142. PMID: 11043834.
- Quatresooz, P., I. Uhoda, I. Fumal, C. Pierard-Franchimont and G.E. Pierard, 2004. Revisiting the gender-linked melanoma burden. *Dermatology*, 209 (3): 197-201. PMID: 15459532.
- Sadjadi, A., R. Malekzadeh, M.H. Derakhshan, A. Sepehr, M. Nouraie, M. Sotoudeh, A. Yazdanbod, B. Shokoochi, A. Mashayekhi, S. Arshi, A. Majidpour, M. Babaei, A. Mosavi, M.A. Mohagheghi and M. Alimohammadian, 2003. Cancer occurrence in Ardabil: Results of a population-based cancer registry from Iran. *Int. J. Cancer*, 107 (1): 113-118. PMID: 12925965.
- San Sebastian, M. and A.K. Hurtig, 2004. Cancer among indigenous people in the Amazon Basin of Ecuador (1985-2000). *Rev. Panam. Salud Publica*, 16 (5): 328-333. PMID: 15729982.
- Schüz, J., D. Schön, W. Batzler, C. Baumgardt-Elms, B. Eisinger, M. Lehnert and C. Stegmaier, 2000. Cancer registration in Germany: Current status, perspectives and trends in cancer incidence (1973-1993). *J. Epidemiol. Biostat.*, 5 (2): 99-107. PMID: 10890281.
- Shahbazi, A., 2006. Evaluating prevalence of radiation related cancers in Chaharmahal va Bakhtiari province for 5 years (1997-2001). *Shahre Kord Med. Sci. J.*, 6 (1): 7-13. <http://sid.ir/en/ViewPaper.asp?ID=62613830107>.
- Siesling, S., M.A. Van Der Aa, J.W. Coebergh and E. Pukkala, 2008. Working group of the Netherlands cancer registry. Time-space trends in cancer incidence in the Netherlands (1989-2003). *Int. J. Cancer*, 122 (9): 2106-2114. PMID: 18183593.
- Soerjomataram, I., W.J. Louwman, V.E. Lemmens, J.W. Coebergh and E. De Vries, 2008. Are patients with skin cancer at lower risk of developing colorectal or breast cancer? *Am. J. Epidemiol.*, 167 (12): 1421-1429. PMID: 18424428.
- Stang, A., E. Pukkala, R. Sankila, B. Soderman and T. Hakulinen, 2006. Time trend analysis of the skin melanoma incidence of Finland from 1953 through 2003 including 16,414 cases. *Int. J. Cancer*, 119 (2): 380-384. PMID: 16477634.
- Stang, A., K. Stang, C. Stegmaier, T. Hakulinen and K.H. Jockel, 2001. Skin melanoma in Saarland: incidence, survival and mortality (1970-1996). *Eur. J. Cancer Prev.*, 10 (5): 407-415. PMID: 11711755.
- Stang, A., C. Stegmaier and K.H. Jöckel, 2003. Nonmelanoma skin cancer in the Federal state of Saarland, Germany (1995-1999). *Br. J. Cancer*, 89 (7): 1205-1208. PMID: 14520447.

- Tousi, P., S.S. Kermani and A.S. Kebria, 2004. Epidemiologic patterns of skin cancers Loghman and the Booali Hospital (2001-2003). *Tehran Univ. Med. J.*, 62 (6): 509-517. <http://sid.ir/en/ViewPaper.asp?ID=13395>.
- Uhoda, I., P. Quatresooz, I. Fumal, A.F. Nikkels, C. Pierard-Franchimont and G.E. Pierard, 2004. Updating trends in cutaneous cancers in South-east Belgium. *Oncol. Rep.*, 12 (1): 111-114. PMID: 15201969.
- Ulmer, M.J., J.M. Tonita and P.R. Hull, 2003. Trends in invasive cutaneous melanoma in Saskatchewan (1970-1999). *J. Cutan. Med. Surg.*, 7 (6): 433-442. PMID: 15926213.
- Vaktskjold, A., J.A. Lebedintseva, D.S. Korotov, A.V. Tkatsjov, T.S. Podjakova and E. Lund, 2005. Cancer incidence in Arkhangelskaja oblast in Northwestern Russia. The Arkhangelsk cancer registry. *BMC Cancer*, 5 (1): 82. PMID: 16029510.
- Verdecchia, A., A. Micheli, M. Colonna, V. Moreno, M.I. Izarzugaza and E. Paci, 2002. Europreval working group. A comparative analysis of cancer prevalence in cancer registry areas of France, Italy and Spain. *Ann. Oncol.*, 13 (7): 1128-1139. PMID: 12176794.
- Valentin, S.M., J.L. Sanchez, L.D. Figueroa and C.M. Nazario, 2007. Epidemiology of melanoma in Puerto Rico (1987-2002). *PR Health Sci. J.*, 26 (4): 343-348. PMID: 18246962.
- Wassberg, C., M. Thörn, A.M. Johansson, R. Bergström, B. Berne and U. Ringborg, 2001. Increasing incidence rates of squamous cell carcinoma of the skin in Sweden. *Acta Derm. Venereol.*, 81 (4): 268-272. PMID: 11720174.
- Yavari, P., B. Sadrolhefazi, M. Mohagheghi, H. Madani, A. Mosavizadeh, A. Nahvijou, Y. Mehrabi and M. Pourhseingholi, 2008. An Epidemiological Analysis of cancer data in an Iranian hospital during the last three decades. *Asian Pac. J. Cancer Prev.*, 9 (1): 145-150. PMID: 18439094.
- Youl, P.H., P.D. Baade, M. Janda, C.B. Del Mar, D.C. Whiteman and J.F. Aitken, 2007. Diagnosing skin cancer in primary care: How do mainstream general practitioners compare with primary care skin cancer clinic doctors? *Med. J. Aust.*, 187 (4): 215-220. PMID: 17708723.
- Zeb, A., A. Rasool and S. Nasreen, 2008. Cancer incidence in the districts of Dir (North West Frontier Province), Pakistan: A preliminary study. *J. Chin. Med. Assoc.*, 71 (2): 62-65. PMID: 18290249.